described in ref 5. The workup included chromatography (3×30 , CH_2Cl_2) and yielded 1.87 g (82%) of **2a**: mp 59–62 °C (lit.²⁷ mp 65–66 °C).

N-(2-Chloroethyl)toluene-4-sulfonamide (3a). Authentic probe: **1a** was added to a mixture of AlCl_3 and CH_2Cl_2 . After 10 min the mixture was washed with water. Evaporation of the organic layer provided **3a**: mp 98 °C (lit.²⁸ mp 99 °C).

Acknowledgment. We thank the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen In-

dustrie for financial support and Professor Hans Suschitzky for the help in improving the English.

Registry No. **1a**, 3634-89-7; **2b**, 5450-75-9; **3a**, 6331-00-6; **4b**, 19871-46-6; **5b**, 117583-57-0; **6**, 1520-42-9; **7a**, 70-55-3; **7b**, 98-10-2; **7c**, 88-05-1; *cis*-**8b**, 110143-77-6; *trans*-**8b**, 110143-78-7; **9b**, 117583-58-1; **10**, 632-50-8; **11a**, 5048-64-6; **11b**, 5048-63-5; **11c**, 117583-63-8; **12aA**, 117583-64-9; **12aP**, 117583-61-6; **12aT**, 117583-69-4; **12bP**, 42801-56-9; **12cA**, 117583-68-3; **12cP**, 117583-59-2; **12cT**, 117583-72-9; **13A**, 117583-65-0; **13P**, 1985-96-2; **13T**, 117583-70-7; **14A**, 117583-66-1; **14P**, 1634-11-3; **14T**, 117583-71-8; **15a**, 2849-72-1; **15c**, 117583-60-5; **16a**, 1206-41-3; **16b**, 1203-15-2; **17a**, 110871-36-8; **18aP**, 110871-37-9; **18aT**, 117583-67-2; **18bP**, 117583-62-7; **PH**, 71-43-2; **AN**, 100-66-3; **TH**, 108-88-3.

(27) Johnson, T. B.; Guest, H. G. *Am. Chem. J.* **1909**, 42, 340-353.

(28) Peacock, D. H.; Dutta, U. C. *J. Chem. Soc.* **1934**, 1303-1305.

Stereochemical Aspects of the "tert-Amino Effect". 1. Regioselectivity in the Synthesis of Pyrrolo[1,2-a]quinolines and Benzo[c]quinolizines

Walter H. N. Nijhuis,[†] Willem Verboom,[†] A. Abu El-Fadl,^{‡,§} Sybolt Harkema,[†] and David N. Reinhoudt^{*,†}

Laboratories of Organic Chemistry and Chemical Physics, University of Twente, 7500 AE Enschede, The Netherlands

Received June 20, 1988

Substituted 2-vinyl-*N,N*-dialkylanilines cyclize in refluxing 1-butanol to give substituted pyrrolo[1,2-a]quinolines and benzo[c]quinolizines. This reaction proceeds via a 1,5-hydrogen transfer and subsequent C-C bond formation. When in the 2-vinyl-*N,N*-dialkylanilines **4**, $\text{R}^1 = \text{H}$ and $\text{R}^2 = \text{H}$ (**4a,d**), CH_3 (**4b,e**), or C_2H_5 (**4f**), the cyclization products **5a,b,d-f** are formed selectively, with the substituent R^2 at the bridgehead carbon atom. This regioselectivity is lost when $\text{R}^2 = \text{CH}_2\text{OCH}_3$ (**4c,g**), and a mixture of the regioisomers **5c,g**, **6c,g**, and **7c,g** is formed. Reaction of compounds **4h-n** ($\text{R}^1 = \text{CH}_3$) yields the pyrrolo[1,2-a]quinolines **5-7(h-j)** and benzo[c]quinolizines **5-7(k-n)** selectively, in which the substituent at the bridgehead carbon atom is at the same face of the molecule (*cis*) as the hydrogen atom at C-5 [**5-7(h-j)**] or at C-6 [**5-7(k-n)**]. The configuration of these compounds was determined by ^1H NOE difference spectroscopy and single-crystal X-ray analysis (**6n**). Heating of **4o-q** ($\text{R}^1 = 4\text{-C}_6\text{H}_4\text{CH}_3$) in refluxing 1-butanol gives mixtures of the *cis* [**5-7(o-q)**] and *trans* [**8-10(o-q)**] compounds. The mechanism of these cyclizations, which are further examples of the "tert-amino effect", and the effect of variation in substituents are discussed.

Introduction

In 1972 Meth-Cohn and Suschitzky reviewed the formation of heterocycles by ring closure of ortho-substituted tertiary anilines (the "tert-amino effect").¹ We have demonstrated that this type of reaction has a wider applicability, e.g. for the synthesis of pyrrolo- and pyrido[1,2-a]indoles, pyrrolo[1,2-a]quinolines, benzo[c]quinolizines, [1,4]oxazino[4,3-a]quinolines,² and pyrazinoquinolines.³ The synthesis of benzoxazines,⁴ benzothiazines, and a quinoxaline⁵ can also be regarded as examples of this type of reaction. The formation of all these compounds takes place via either a 1,5- or a 1,6-hydrogen shift, depending on the structure of the reactant. The two different types of dipolar species subsequently undergo cyclization to give 6- and 5-membered rings, respectively.

In the course of our investigations of the "tert-amino effect" we have studied the formation of pyrrolo[1,2-a]quinolines and benzo[c]quinolizines in more detail, in particular the regioselective aspects. When we could control the regioselectivity of the cyclization reaction this would greatly enhance the synthetic utility of the "tert-

amino effect" in heterocyclic synthesis.

In the present paper dealing with this regioselectivity we describe the thermal isomerization of 2-vinyl-*N,N*-dialkylanilines **4** ($\text{X} = -, \text{CH}_2$) with different substituents at the α -carbon atom of the vinyl moiety ($\text{R}^1 = \text{H}$, CH_3 , or $4\text{-C}_6\text{H}_4\text{CH}_3$) and at one of the two carbon atoms adjacent to nitrogen of the amine moiety ($\text{R}^2 = \text{H}$, CH_3 , CH_2CH_3 , or CH_2OCH_3). Firstly, we describe the effect of the nature (size, stabilizing effect) of the substituent R^2 when $\text{R}^1 = \text{H}$. Secondly, the influence of both the substituents $\text{R}^1 (\neq \text{H})$ and R^2 on the regioselectivity of the cyclization will be discussed.

Results

Synthesis of the Starting Materials 4. The starting compounds **4** for the thermal isomerization were conven-

(1) Meth-Cohn, O.; Suschitzky, H. *Adv. Heterocycl. Chem.* **1972**, 14, 211.

(2) Verboom, W.; Reinhoudt, D. N.; Visser, R.; Harkema, S. *J. Org. Chem.* **1984**, 49, 269.

(3) Nijhuis, W. H. N.; Verboom, W.; Reinhoudt, D. N. *Synthesis* **1987**, 641.

(4) Verboom, W.; van Dijk, B. G.; Reinhoudt, D. N. *Tetrahedron Lett.* **1983**, 24, 3923.

(5) Verboom, W.; Hamzink, M. R. J.; Reinhoudt, D. N.; Visser, R. *Tetrahedron Lett.* **1984**, 25, 4309.

[†] Laboratory of Organic Chemistry.

[‡] Laboratory of Chemical Physics.

[§] Present address: Physics Department, Faculty of Science, Assiut University, Assiut, Egypt.

Chart I

Table I. Yields and Reaction Times of the Products Obtained by Thermal Isomerization of the Condensation Compounds 4a-q

starting compd 4	reaction time	yield, %		
		5	6	7
a	2 h	82		
b	2 h	85		
c	1.5 h	46	19	17
d	2 h	78		
e	1.5 h	79 ^a		
f	1.5 h	80 ^a		
g	2.5 h	71	12.5	12.5
h	5 h	79		
i	5 h	79		
j	5 h	33	35	6
k	2 h	88		
l	2 h	88		
m	2.5 h	86		
n	3 h	56	28	8
o	3 days	86 ^b		
p	3 days	76 ^b	13 ^b	9 ^b
q	3 days	16 ^b	41 ^b	15 ^b

^a Overall yield, starting from 3e and 3f, respectively. ^b Total yields for the formed cis (5-7) and trans isomers (8-10) starting from 3o-q.

step comprises a Knoevenagel condensation of the carbonyl group of 3 with malononitrile in toluene at room temperature to give 4a-g and 4h-n in high yields (Chart I). In the case of 2-(2-methyl- and 2-ethyl-1-piperidinyl)-benzaldehyde (3e and 3f) the malononitrile adduct could not be obtained pure because mixtures of the condensation product (4e or 4f) and the product of ring closure (5e or 5f) were isolated.

The condensation of the benzophenone derivatives (3o-q) did not take place under these conditions. However, by elevating the reaction temperature, the condensation of 3o-q to 4o-q could be accomplished. Since at the higher reaction temperatures ring closure proceeds (much) faster than condensation, hardly any condensation product was present in these reaction mixtures, and therefore no attempt was made to isolate compounds 4o-q.^{8,12}

Thermal Isomerization of 2-Vinyl-N,N-dialkylanilines 4a-q. The objective of our work was to study the regioselectivity of the cyclization of 2-vinyl-N,N-dialkylanilines (4a-q). Firstly, we will describe the effect on the regioselectivity by variation of R^2 (when $R^1 = H$) and the N,N-dialkylamino group is a pyrrolidinyl ($X = -$) or a piperidinyl ($X = CH_2$) moiety. Secondly, the effect of various R^2 groups will be described when $R^1 = CH_3$ ($X = -$, CH_2). Thirdly, the effect of R^2 on the regioselectivity when $R^1 = 4-C_6H_4CH_3$ ($X = -$) will be described.

Cyclization of 4a-g ($R^1 = H$). Compound 4a was reacted in refluxing 1-butanol to give 1,2,3,4a-tetrahydropyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5a) in a yield of 82% (Table I).² Heating of 4b in refluxing 1-butanol afforded selectively 5b with the substituent R^2

iently prepared from 2-fluorobenzaldehyde (1a), 2-fluoroacetophenone (1b), and 2-fluoro-4'-methylbenzophenone (1c). The latter compound was synthesized from 2-fluorobenzaldehyde via a Grignard reaction with (4-methylphenyl)magnesium bromide and subsequent oxidation of the resulting secondary alcohol with pyridinium chlorochromate (PCC) in dichloromethane.⁶

The first step in the synthesis of compounds 4 involves a nucleophilic substitution of the fluorine atom in these compounds by a pyrrolidine (2a-c) or a piperidine group (2d-g) in refluxing N,N-dimethylformamide (DMF) in the presence of K_2CO_3 .⁷ These reactions afford the substituted benzaldehydes 3a-g, acetophenones 3h-n, and benzophenones 3o-q, respectively (Chart I). The second

(8) Unfortunately, we could not isolate the compounds 4o-q, though several attempts have been made. A Knoevenagel condensation with $TiCl_4$ and base did not give the desired compounds;⁹ a Peterson olefination reaction¹⁰ and variations thereof and a simple Wittig reaction¹¹ to obtain the olefins without the cyano groups did not succeed.

(9) Lehnert, W. *Tetrahedron* 1973, 29, 635.

(10) Peterson, D. J. *J. Org. Chem.* 1968, 33, 780.

(11) Wittig, G.; Schölkopf, U. *Chem. Ber.* 1954, 87, 1318.

(12) The condensation reaction and subsequent cyclization, here in most cases described as a two step process, can also be performed in a one-pot procedure. This is demonstrated by the reaction of 3o-q with malononitrile in refluxing 1-butanol. Heating of the aldehyde (3a-g) and acetophenone derivatives 3h-n with malononitrile in either refluxing toluene or refluxing 1-butanol gives the cyclized products 5-7(a-n) in comparable yields.

(6) Corey, E. J.; Suggs, J. W. *Tetrahedron Lett.* 1975, 2647.

(7) Niewiadomski, K. B.; Suschitzky, H. *J. Chem. Soc., Perkin Trans. I* 1975, 1679.

(= CH₃) at the bridgehead carbon atom in 85% yield. This was concluded from the ¹H and ¹³C NMR spectral data [¹H NMR δ 1.26 (s, CH₃); ¹³C NMR δ 63.8 [s, NC(CH₃)]].

However, this regioselectivity was lost in the reaction of **4c** (R² = CH₂OCH₃). In this case we obtained a mixture of *one* diastereomer with the methoxymethyl substituent at C-3a (bridgehead carbon atom) (**5c**, 46%) and two diastereomers with a hydrogen atom at C-3a, that could be separated by column chromatography. Each of these two compounds exhibits very characteristic and different signals for the methylene protons of the CH₂OCH₃ group in the 200-MHz ¹H NMR spectrum (see the Experimental Section). On the basis of the ¹H NMR data, compared with those of **6j** and **7j** (vide infra), we concluded that these are **6c** (19%) and **7c** (17%).

Ring closure of the piperidinyl analogue **4d** (R¹ = R² = H) resulted in the formation of 2,3,4,4a-tetrahydro-1H-benzo[c]quinolizine-5,5(6H)-dicarbonitrile (**5d**) in a yield of 78%.² Heating of crude (vide supra) **4e** (R¹ = H, R² = CH₃) and of **4f** (R¹ = H, R² = CH₂CH₃) in refluxing 1-butanol gave selectively **5e** and **5f** in overall yields of 79% and 80% (based on **3e** and **3f**), respectively. In these reactions the same regioselectivity is observed as in the ring closure of the pyrrolidinyl analogue **4b**.

However, this regioselectivity was again lost when a methoxymethyl substituent is present. Reaction of **4g** (R¹ = H, R² = CH₂OCH₃) in refluxing 1-butanol resulted in the formation of *one* diastereomer with the methoxymethyl group at the bridgehead carbon atom (C-4a) (**5g**, 71%), and a mixture of *two* diastereomers with a hydrogen atom at C-4a (**6g** and **7g**, total yield 25%). Compound **5g** could be separated from these two compounds by column chromatography. Compounds **6g** and **7g** could not be separated in this way, but 200-MHz ¹H NMR spectroscopy revealed the 1:1 ratio of these two diastereomers. The two isomers exhibit different multiplets at δ 4.35–4.2 and 3.95–3.8 for the NCH(CH₂OCH₃) proton. However, upon fractional crystallization one isomer could be obtained pure.¹³ The methylene protons of the CH₂OCH₃ group of this isomer exhibit the same characteristic absorptions as those of **6n** (vide infra). Therefore, we could assign the structure **6g** to this compound.

Cyclization of 4h–n (R¹ = CH₃). Subsequently, we studied the reactions of the compounds **4h–j** (X = –) and **4k–n** (X = CH₂) in which the α-carbon atom of the vinyl moiety is a prochiral center. When the pyrrolidinyl derivative **4h** was heated in refluxing 1-butanol, exclusively **5h** was obtained in 79% yield. The piperidinyl analogue **4k** gave also selectively **5k** (88%).

The compounds **5i** (X = –, R² = CH₃), **5l** (X = CH₂; R² = CH₃), and **5m** (X = CH₂; R² = CH₂CH₃) were formed regioselectively by cyclization of **4i**, **4l**, and **4m** in yields of 79%, 88%, and 86%, respectively. Three compounds, **5j** (33%), **6j** (35%), and **7j** (6%), were isolated upon ring closure of **4j** (X = –, R² = CH₂OCH₃). Cyclization of **4n** (X = CH₂; R² = CH₂OCH₃) gave one isomer with the methoxymethyl substituent at the bridgehead carbon atom (**5n**, 56%) and a mixture of two diastereomers, **6n** and **7n**, with a hydrogen atom at the bridgehead carbon atom in a combined yield of 36%. These two diastereomers are formed in a ratio of 7:2, as was determined by ¹H NMR spectroscopy. The NCH(CH₂OCH₃) proton of the major

Chart II

Chart III

isomer (**6n**) exhibits a multiplet at δ 4.4–4.25, and of the minor isomer (**7n**) a multiplet at δ 3.95–3.8.

The structures of **5i** (X = –, R² = CH₃) and **5l** (X = CH₂, R² = CH₃) were determined by ¹H NOE difference spectroscopy. This indicated that in both compounds the methyl group at the bridgehead carbon atom and the hydrogen at the benzylic position are at the *same face* of the molecule. Irradiation of the bridgehead methyl group (CH₃-3a) of **5i** gave a NOE of 7% at H-5. In reverse, on irradiation of H-5 only a slight enhancement of 2% was noticed at CH₃-3a. This difference arises because the protons of CH₃-3a have additional and different protons contributing to their relaxation.¹⁴ The same difference was noticed by irradiating H-6 and CH₃-4a of compound **5l** (Chart II).

¹H NOE difference spectroscopy experiments with **5m** (X = CH₂; R² = C₂H₅) gave almost identical information about the relative positions of the ethyl group and the hydrogen atom at C-6. Irradiation of the methyl group of the CH₂CH₃ moiety gave a NOE of 4% at H-6 but upon irradiation of H-6 no NOE was measured at the ethyl group (Chart II). ¹H NOE difference spectroscopy of **5k** gave no reliable results about the position of the hydrogen atoms at C-4a and C-6. However, on the basis of the results obtained by ¹H NOE experiments with **5i**, **5l**, and **5m**, we assume that the relative configuration of **5k** is as depicted (Chart I). The relative configuration of the hydrogen atoms at C-3a and C-5 of compound **5h** could not be assigned in this way because the absorptions of the NCH₂ hydrogen atoms (H-1) and H-5 coincide. Irradiation of H-3a gave a NOE at H-5 and H-1 (≈3%). In reverse, irradiation of H-5 and H-1 gave an enhancement at H-3a (≈5%). Since we have never noticed any effect at H-1 by

(13) Unfortunately, we were not able to determine the relative configuration of this isomer by means of ¹H NOE difference spectroscopy: irradiation of the bridgehead hydrogen atom (H-4a) gave no measurable effect on H-1 and irradiation of H-1 gave no effect on H-4a. Attempts to measure the effect of irradiation of H-4a at the methylene hydrogen atoms of the CH₂OCH₃ group and reverse were doomed to fail, because the absorptions of these protons in the ¹H NMR spectrum coincide.

(14) Similar observations have been reported, e.g.: Bell, R. A.; Saunders, J. K. *Can. J. Chem.* 1970, 48, 4114. Bachers, G. E.; Schaefer, T. *Chem. Rev.* 1971, 71, 617. Noggle, J. H.; Schirmer, R. E. *The Nuclear Overhauser Effect*; Academic Press: New York, 1971; p 182 and references cited therein.

give singlets at δ 4.54 and 4.58, while the minor isomers, **9q** and **10q**, absorb at lower field, viz. δ 4.62 and 4.72, respectively.

The structures of **5o** ($R^2 = H$) and **5p** ($R^2 = CH_3$) were proven by 1H NOE difference spectroscopy. Compound **5o** ($R^2 = H$) could be obtained pure by fractional crystallization, and the cis relationship between the hydrogen atoms at the bridgehead carbon atom (C-3a) and at C-5 was determined by 1H NOE difference spectroscopy (Chart IV). Consequently, in the minor isomer (**8o**), H-3a and H-5 are related trans. The absorption of H-5 of the two isomers in the 1H NMR spectrum is very significant: in **5o** H-5 absorbs at higher field (δ 4.55) than in **8o** (δ 4.68). This is a general feature of the compounds with $R^1 = 4-C_6H_4CH_3$ (vide infra, assignment of **6q** and **7q**).

Compound **5p** ($R^2 = CH_3$), the major isomer obtained by cyclization of **4p**, could also be isolated pure by fractional crystallization. On basis of the absorption of the benzylic proton, we have assigned the structure **5p**. Compound **8p** was obtained as a mixture together with the two regioisomers with a hydrogen atom at C-3a. On basis of the absorptions of the benzylic protons of these two compounds we have assigned the structures **6p** and **7p** to these isomers.

The major isomer with the methoxymethyl group at the bridgehead carbon atom, obtained by cyclization of **4q**, could be obtained pure by fractional crystallization. On basis of 1H NMR absorptions, in particular the absorption of the benzylic proton (H-5), structure **5q** was assigned to this compound. The major isomers (**6q** and **7q**) of the two isomeric mixtures subsequently isolated could be obtained pure by fractional crystallization, and 1H NOE difference spectroscopy pointed out that the hydrogen atoms at C-3a and at C-5, in both compounds, are at the same face of the molecule (Chart IV). Also, in these compounds, **6q** and **7q**, H-5 absorbs at higher field than in the minor isomers **9q** and **10q**. The methylene protons of the CH_2OCH_3 group of **6q** reveals the same characteristic ABX pattern in the 1H NMR spectrum as compounds **6c** ($R^1 = H$) and **6j** ($R^1 = CH_3$). The absorptions of the corresponding protons in **7q** show a different specific pattern and can be compared with that of the compounds **7c** and **7j**. Therefore, the structures **6q** and **7q** were assigned to these two compounds.

Discussion

The thermal isomerization of the compounds **4** to the substituted pyrroloquinolines and benzoquinolizines **5–10** proceeds via a 1,5-hydrogen shift followed by subsequent cyclization.¹⁸

When $R^1 = R^2 = H$ the compounds **4a** ($X = -$) and **4d** ($X = CH_2$) can only give the ring closed products **5a** and **5d**, respectively. When R^2 is a methyl (**4b**, **4e**) or an ethyl group (**4f**), the cyclized product with the substituent R^2 at the bridgehead carbon atom is formed selectively (**5b**, **5e**, and **5f**, respectively). This is due to a more effective stabilization of a tetrasubstituted iminium double bond

Scheme I

in the dipolar intermediate (A, Scheme I), generated when the hydrogen atom at the substituted carbon atom undergoes a 1,5-hydrogen shift, in comparison with a tri-substituted double bond (B, Scheme I), generated when one of the α -methylene protons of the other carbon atom is transferred.

Therefore C–C bond formation will take place selectively at the carbon atom bearing the substituent R^2 .

When R^2 is a methoxymethyl group, a sterically more hindering group and one that stabilizes the iminium double bond less compared with a methyl (or ethyl) group because of the inductive electron-withdrawing effect of the oxygen atom, the regioselectivity is lost. Consequently, the 1,5-hydrogen shift will not take place *exclusively* from the carbon atom adjacent to nitrogen bearing the methoxymethyl substituent, and mixtures of regioisomers (**5**, **6**, and **7**) are formed.

Cyclization of **4h–n**, in which R^1 is a methyl group, gave additional information about the mechanism. The structure of **5i** ($X = -$; $R^1 = R^2 = CH_3$) and **5l** ($X = CH_2$; $R^1 = R^2 = CH_3$), for instance, that are formed *selectively* by reaction of **4i** and **4l** revealed that the hydrogen atom in the product molecule which underwent a 1,5-hydrogen shift is related *cis* to the substituent R^2 . This was confirmed recently by the formation of one of the *enantiomers* of **5j** and the subsequent X-ray analysis of a bromo derivative.¹⁵

The position of the vinyl moiety in **4** determines the stereochemistry of **5**, **6**, and **7**. X-ray analysis of **4a** showed that the β -carbon atom of the vinyl moiety of this molecule points away from the amine ring.¹⁹ Important for the hydrogen shift to take place is the proximity of the migrating hydrogen of the $NCHR^2$ moiety to the α -carbon atom of the vinyl moiety, when **5** is formed, or from the NCH_2 moiety when **6** and **7** are formed.

When the hydrogen atom attached to the carbon atom with the substituent R^2 is transferred (and compounds **5** are formed), it takes place from the conformer in which the β -carbon atom of the vinyl moiety is below the plane of the aromatic ring (Scheme II). It seems likely that the hydrogen migration takes place suprafacially. As the vinyl moiety is already rotated when this hydrogen is migrating, subsequent rotation of the former vinyl group will probably

(18) The crucial step in the mechanism is the transfer of hydrogen. We have always assumed that the thermal isomerizations of the compounds **4a–q** to the substituted pyrroloquinolines and benzoquinolizines proceeds via a concerted suprafacial 1,5-hydrogen shift followed by subsequent cyclization.²³ A concerted suprafacial hydrogen transfer explains the discrete chirality in the product molecules,¹⁸ and the primary isotope effect and solvent effect are supporting this. However, on basis of recent experiments we cannot exclude a direct intramolecular transfer of hydrogen with a partial negative charge,¹⁹ when the NCH_2 or $NCHR^1$ unit is in the proximity of the dicyanovinyl moiety. The hydrogen transfer may be an intramolecular variant of a Meerwein–Ponndorf–Verley reaction. In literature, there are precedences of such base-catalyzed intramolecular hydride transfers.^{20,21}

(19) Groenen, L. C.; Verboom, W.; Nijhuis, W. H. N.; Reinhoudt, D. N.; van Hummel, G. J.; Feil, D. *Tetrahedron* 1988, 44, 4637.

(20) Warnhof, E. W. *Can. J. Chem.* 1977, 55, 1635 and references cited therein.

(21) Menicagli, R.; Giacomelli, G. P.; Lardicci, L. *J. Organomet. Chem.* 1973, 50, C15.

Scheme II

continue in the same direction to enable C-C bond formation. The complete reaction is thus stereochemically well defined. Hence, compounds **5** are formed in which the substituent R^2 at the bridgehead is at the same face of the molecule as the hydrogen atom that has shifted.

In the compounds **6** and **7** the hydrogen atom at the bridgehead and the hydrogen which is transferred are at the same face of the molecule. From these results we conclude that in the dipolar intermediates the carbanion adds to the iminium double bond from the same face as where the hydrogen atom, which underwent a 1,5-shift, originally was located.

The configurations of **6c** and **7c** differ in the position of the hydrogen atom at C-3a related to the hydrogen atom located at the carbon atom bearing the methoxymethyl group (C-1). Because the shift of H_a and H_b , which determines the ratio of **6c** and **7c**, takes place in almost equal ratio, **6c** and **7c** are formed in a ratio of $\approx 1:1$.

In the formation of **6j** and **7j** one of the two hydrogen atoms, H_a or H_b , respectively, undergoes a 1,5-hydrogen shift. However, these shifts are not proceeding in a 1:1 ratio. CPK models reveal that in **4j**, when H_a or H_b migrates, the pyrrolidinyl ring is out of plane with the phenyl group because of steric hindrance caused by the substituent R^1 (in this case a methyl group). When H_b is transferred, and **7j** is formed, there is severe steric hindrance between the methoxymethyl group and the aromatic ring. In the product molecule **7j**, the methoxymethyl group and $H-9$ are in close proximity, as was determined by 1H NOE difference spectroscopy (vide supra). The steric hindrance between the methoxymethyl group and $H-9$ is much less in the case of H_a shifts, and **6j** is formed. Since the 1,5-hydrogen shift is the rate-determining step, the product ratio of **6j** and **7j** is determined

Scheme III

by the ratio of shifts of H_a and H_b , and so **6j** is formed favorably.

When R^1 is larger, as is the case in the formation of **6q** and **7q** (when $R^1 = 4-C_6H_4CH_3$), the increased steric hindrance causes the pyrrolidinyl ring with the methoxymethyl group to turn even more out of the plane with the aromatic ring, compared with the formation of **6j** and **7j**. As a result, the hindrance between the CH_2OCH_3 group and the aromatic ring, when H_b is transferred, and **7q** is formed, is much less than in the formation of **7j**. The ratio of **6q:7q** will therefore decrease compared with that of **6j:7j**. The total yield of **6q** and **7q** exceeds that of **6j** and **7j** because of the increased steric hindrance between the substituents R^1 and R^2 .

As was mentioned, the cyclization of **4a-n** takes place in a stereochemically well defined way, resulting in the formation of compounds **5-7** with the hydrogen or the substituent R^1 at the bridgehead at the same face of the molecule as the hydrogen that is transferred. However, the cyclization of **4o-q** when R^1 is a 4-methylphenyl group gives besides **5-7(o-q)** also the compounds **8-10(o-q)**, which have the opposite configuration at C-5. When R^1 is a methyl group or a hydrogen, no trace of these products, in which $H-3a$ and $H-5$ are at the opposite face of the molecule, could be detected. We can explain this result by assuming that there is an interchange of the position of the vinyl group (Scheme III), because of steric hindrance caused by the substituent R^1 at the α -position of the amine

ring. Cyclization then partly takes place from the conformation in which the 4-methylphenyl group points away from the amine ring (Scheme III). In this situation, hydrogen migration leads to the formation of an intermediate with a new asymmetric center that has the opposite configuration compared with the intermediates generated by formation of 5-7. Attack of the generated carbanion from the side the hydrogen was transferred gives the compounds 8-10. The bridgehead asymmetric center has the same configuration as in the products 5-7; the new asymmetric center at C-5, however, has the opposite configuration. So, in the case of cyclization of 40-q, the geometry of the transition state is less well defined.

In conclusion, we have demonstrated that by proper choice of the substituents R^1 and R^2 we can direct the regio- and stereoselectivity in the formation of pyrrolo[1,2-a]quinolines and benzo[c]quinolizines via the "tert-amino effect". When R^1 and R^2 are substituents like a hydrogen atom or a methyl group, the cyclization proceeds regioselectively. Substituents (R^2), like a methoxymethyl group, that are sterically more hindering and less stabilizing disturb the regioselectivity. The size of the substituent R^1 determines the relative configuration of the substituents at the bridgehead carbon atom and at the benzylic position in the product molecule because it directs the position of the vinyl moiety in the starting material.

Experimental Section

Melting points were determined with a Reichert melting point apparatus and are uncorrected. ^1H NMR spectra (CDCl_3) were recorded with a Bruker WP-80 or a Nicolet NT 200-WB spectrometer (Me_4Si as an internal standard). ^{13}C NMR spectra (CDCl_3) were recorded with a Nicolet NT 200-WB spectrometer (Me_4Si as an internal standard). Mass spectra were recorded with a Varian Mat 311A spectrometer and IR spectra with a Perkin-Elmer 257 spectrophotometer. Elemental analyses were carried out by A. Christenhusz of the Laboratory of Chemical Analysis of the University of Twente.

2-Fluorobenzaldehyde (1a) (Janssen), 2-fluoroacetophenone (1b) (Janssen), pyrrolidine (2a) (Merck), 2-methylpyrrolidine (2b) (Alfa), piperidine (2d) (Merck), 2-methylpiperidine (2e) (Merck), and 2-ethylpiperidine (2f) (Merck) were used without further purification.

($\pm$)-(2-Methoxymethyl)pyrrolidine (2c) was obtained from L-proline via a racemization to DL-proline²² and subsequent conversion to ($\pm$)-(2-methoxymethyl)pyrrolidine as described by Enders et al.²³ ($\pm$)-(2-Methoxymethyl)piperidine (2g) was obtained from ($\pm$)-(2-hydroxymethyl)piperidine via the same route as was used for the conversion of DL-proline to ($\pm$)-(2-methoxymethyl)pyrrolidine.²³

All reactions were carried out under a nitrogen atmosphere.

2-Fluoro-4'-methylbenzophenone (1c). To a solution of (4-methylphenyl)magnesium bromide in diethyl ether (0.65 M; 500 mL) was added a solution of 2-fluorobenzaldehyde (1a; 25.0 g; 0.2 mol) in diethyl ether (250 mL) at such a rate that the temperature in the flask stayed between -10 and -5°C . The mixture was allowed to warm to room temperature and was stirred for 2 h. The crude reaction mixture was added to a saturated NH_4Cl solution (400 mL). After separation of the organic phase, the water layer was extracted with diethyl ether (2×200 mL). The combined organic layers were dried (MgSO_4). The solvent was removed under reduced pressure, and the residue was purified by column chromatography (SiO_2 ; CH_2Cl_2) to give the alcohol as an oil in 75% yield: ^1H NMR δ 7.65-6.9 (m, 8 H, Ar H), 6.11 [d, 1 H, $J = 4.0$ Hz, $\text{CH}(\text{OH})$], 2.32 (s, 3 H, CH_3), 2.22 [d, 1 H, $J = 4.0$ Hz, $\text{CH}(\text{OH})$]; ^{13}C NMR δ 159.7 (d, $J_{\text{C-F}} = 245.7$ Hz, Ar CF), 138.1 and 137.3 (s, Ar C), 131.0 (d, $J_{\text{C-F}} = 12.2$ Hz, C-1), 129.1, 128.7, 127.4, 126.3, and 124.1 (d, Ar C), 115.0 (d, $J_{\text{C-F}} = 21.5$ Hz,

C-3), 69.7 [d, $\text{CH}(\text{OH})$], 21.0 (q, CH_3); IR (KBr) 3350 (OH) cm^{-1} ; mass spectrum, m/e 216.095 (M^+ , calcd for $\text{C}_{14}\text{H}_{13}\text{FO}$ 216.095).

A solution of the secondary alcohol (4.33 g; 20 mmol) in dry CH_2Cl_2 (15 mL) was added dropwise to a suspension of PCC (6.47 g; 30 mmol) in dry CH_2Cl_2 (25 mL). The mixture was stirred for 8 h and subsequently decanted. The solvent was removed under reduced pressure, and the residue was purified by column chromatography (SiO_2 ; CH_2Cl_2) to yield 1c as a solid in a yield of 92%: mp 74.5-75.5 $^\circ\text{C}$ (MeOH); ^1H NMR δ 7.85-7.0 (m, 8 H, Ar), 2.43 (s, 3 H, CH_3); ^{13}C NMR δ 193.0 (s, $\text{C}=\text{O}$), 159.9 (d, $J_{\text{C-F}} = 251.9$ Hz, Ar CF), 144.3 and 134.8 (s, Ar C), 132.7, 130.6, 129.9, and 128.1 (d, Ar C), 127.4 (d, $J_{\text{C-F}} = 18.0$ Hz, C-1), 124.2 (d, Ar C), 116.2 (dd, $J_{\text{C-F}} = 21.5$ Hz, C-3), 21.7 (q, CH_3); IR (KBr) 1650 ($\text{C}=\text{O}$) cm^{-1} ; mass spectrum, m/e 214.079 (M^+ , calcd 214.079). Anal. Calcd for $\text{C}_{14}\text{H}_{11}\text{FO}$ (M , 214.240): C, 78.49; H, 5.18. Found: C, 78.24; H, 5.11.

General Procedure for the Synthesis of the Benzaldehyde (3a-g), Acetophenone (3h-n), and Benzophenone Derivatives (3o-q). To a solution of 2-fluorobenzaldehyde (1a), 2-fluoroacetophenone (1b), or 2-fluoro-4'-methylbenzophenone (1c) (20 mmol) and a 2-substituted pyrrolidine or piperidine (23 mmol) in DMF (20 mL) was added potassium carbonate (3.18 g; 23 mmol), and the mixture was heated for several hours at 152°C . When the reaction was complete as followed from TLC, the reaction mixture was allowed to cool. The crude reaction mixture was taken up in water (100 mL) and extracted with ethyl acetate (3×75 mL). The combined organic layers were washed with a saturated NH_4Cl solution (3×75 mL) and subsequently dried (MgSO_4). The solvent was removed under reduced pressure, and the residue was purified either by distillation or by column chromatography (SiO_2 ; CH_2Cl_2). For experimental and spectral data, see Table II.

General Procedure for the Condensation of the Compounds 3a-n. Synthesis of the 2-Vinyl-N,N-dialkylanilines 4a-n. To a solution of the aldehydes 3a-g or the ketones 3h-n (10 mmol) in toluene (10 mL) was added malononitrile (0.67 g; 10 mmol) in one portion. After the mixture was stirred for several hours at room temperature, the solvent was removed under reduced pressure. The residue was purified either by crystallization, when solid compounds were obtained, or by column chromatography (SiO_2 ; CH_2Cl_2) when oils were obtained (except for 4e and 4f, see below). For experimental and spectral data, see Table III.

2-[[2-(2-Methyl-1-piperidinyl)phenyl]methylene]propanedinitrile (4e). After the mixture was stirred for 5 h, the solvent was removed under reduced pressure. The resulting oil could not be crystallized from organic solvents, and slow cyclization took place upon standing. When the oil was passed through a column of silica gel (CH_2Cl_2) also cyclization took place, affording a mixture of 4e and 5e, which could not be separated. 4e: ^1H NMR δ 8.39 (s, 1 H, $\text{CH}=\text{N}$), 8.25-8.05 (m, 1 H, Ar H), 7.7-7.45 (m, 1 H, Ar H), 7.3-7.1 (m, 2 H, Ar H), 3.3-2.6 (m, 3 H, NCH_2 , NCH), 2.05-1.15 (m, 6 H, CH_2), 0.87 (d, 3 H, $J = 6.4$ Hz, CH_3).

2-[[2-(2-Ethyl-1-piperidinyl)phenyl]methylene]propanedinitrile (4f). After the mixture was stirred for 28 h, the same procedure was followed as for 4e. A mixture of 4f and 5f was obtained, which could not be separated. 4f: ^1H NMR δ 8.29 (s, 1 H, $\text{CH}=\text{N}$), 8.25-8.0 (m, 1 H, Ar H), 7.7-7.4 (m, 1 H, Ar H), 7.3-6.9 (m, 2 H, Ar H), 3.2-2.6 (m, 3 H, NCH_2 , NCH), 2.1-1.7 (m, 6 H, CH_2), 1.45-1.25 (m, 2 H, CH_2CH_3), 0.70 (t, 3 H, $J = 7.6$ Hz, CH_3).

Cyclization of the Knoevenagel Products 4a-n. Synthesis of the Pyrrolo[1,2-a]quinoline [5-7(a-c and h-j)] and Benzo[c]quinolizine Derivatives [5-7(d-g and k-n)]. A solution of the condensation product 4 (10 mmol) in 1-butanol (10 mL) was heated (118°C) for several hours until all the starting material had disappeared according to TLC. The solvent was removed under reduced pressure and the residue purified by column chromatography (SiO_2 ; CH_2Cl_2). The reaction times and yields are summarized in Table I.

Reaction of the Benzophenone Derivatives 3o-q with Malononitrile. Synthesis of the Pyrrolo[1,2-a]quinolines 5-10(o-q). To a solution of the benzophenone derivative 3o,p, or q (10 mmol) in 1-butanol (10 mL) was added malononitrile (0.67 g; 10 mmol). The mixture was heated until all the starting material, including the in situ generated compounds 4, had disappeared according to TLC. The solvent was removed under re-

(22) Yamada, S.; Hongo, C.; Yoshioka, R.; Chibata, I. *J. Org. Chem.* 1983, 4, 843.

(23) Enders, D.; Eichenauer, H. *Chem. Ber.* 1979, 112, 2933.

Table II. Experimental and Spectral Data of the Aldehydes and Ketones 3^a

compd	reaction time, h	yield, %	bp, °C (mmHg)	n_D^{20}	mass spectrum, m/e (M^+ , calcd)	^1H NMR, δ	^{13}C NMR, δ
3b ^b	2.5	76	122–124 (1)	1.5978	189.115 (189.115)	10.11 (CHO)	190.1 (CHO)
3c ^c	2.5	81	126–129 (0.06)	1.5829	219.126 (219.126)	10.11 (CHO)	190.4 (CHO)
3e ^d	16	71	120–122 (1)	1.5563	203.131 (203.131)	10.52 (CHO)	192.4 (CHO)
3f ^e	25	52	154–156 (1)	1.5583	217.148 (217.147)	10.42 (CHO)	192.2 (CHO)
3g ^f	48	58	118–120 (0.07)	1.5530	233.141 (233.142)	10.47 (CHO)	192.2 (CHO)
3i ^g	3.5	82	93–95 (1.6)	1.5756	203.131 (203.131)	2.52 (COCH ₃)	202.7 (COCH ₃)
3j ^h	3	76	137–139 (0.12)	1.5614	233.141 (233.142)	2.54 (COCH ₃)	202.6 (COCH ₃)
3k	24	52	91–92 (0.3)	1.5557	203.131 (203.131)	2.64 (COCH ₃)	204.4 (COCH ₃)
3l ⁱ	72	53	98–100 (0.09)	1.5398	217.147 (217.147)	2.65 (COCH ₃)	204.1 (COCH ₃)
3m ^j	84	46	110–112 (0.1)	1.5390	231.160 (231.162)	2.65 (COCH ₃)	204.4 (COCH ₃)
3n ^k	84	45	132–134 (0.08)	1.5527	247.152 (247.157)	2.66 (COCH ₃)	203.6 (COCH ₃)
3o	10	81 (oil)			265.146 (265.147)	2.41 (ArCH ₃)	196.6 (C=O)
3p ⁱ	10	85 (oil)			279.159 (279.162)	2.40 (ArCH ₃)	197.2 (C=O)
3q ^m	12	81 (oil)			309.173 (309.173)	2.36 (ArCH ₃)	197.2 (C=O)

^a Compounds 3a,d,h are described in ref 2 and 5. All compounds 3 exhibit in the IR (KBr) spectrum an absorption between 1645 and 1680 cm^{-1} . ^{b–m} ^1H NMR δ : ^b 1.17 (d, 3 H, J = 5.9 Hz, CH₃); ^c 3.24 (s, 3 H, OCH₃); ^d 0.92 (d, 3 H, J = 6.3 Hz, CH₃); ^e 0.70 (t, 3 H, J = 7.2 Hz, CH₂CH₃); ^f 3.09 (s, 3 H, OCH₃); ^g 1.15 [d, 3 H, J = 5.6 Hz, NCH(CH₃)]; ^h 3.54 (ABX, 1 H, J = 9.1 and 7.1 Hz, CHHOCH₃), 3.26 (ABX, 1 H, J = 9.3 and 3.7 Hz, CHHOCH₃); ⁱ 0.84 [d, 3 H, J = 6.1 Hz, NCH(CH₃)]; ^j 0.69 (t, 3 H, J = 9.0 Hz, CH₂CH₃); ^k 2.66 (s, 3 H, CH₂OCH₃); ^l 1.15 [d, 3 H, J = 6.1 Hz, NCH(CH₃)]; ^m 3.55 (ABX, 1 H, J = 9.3 and 7.6 Hz, CHHOCH₃), 3.15 (ABX, 1 H, J = 9.3 and 3.7 Hz, CHHOCH₃).

Table III. Experimental and Spectral Data of the Knoevenagel Condensation Products 4a–n^a

compd	reaction time, h	yield, %	mp, °C (MeOH)	mass spectrum, m/e (M^+ , calcd)	^1H NMR, δ	^{13}C NMR, δ
4b ^b	5	86	58–60	237.125 (237.127)	7.99 (=CH)	159.0 (=CH)
4c	48	96	77.5–78.5	267.138 (267.137)	8.1–7.9 (=CH)	159.0 (=CH)
4g ^c	24	90	oil	281.152 (281.153)	8.43 (=CH)	158.1 (=CH)
4h	24 ⁱ	85	111–113	237.127 (237.127)	2.58 (CH ₃)	178.5 [=C(CH ₃)]
4i ^d	30 ⁱ	95	oil	251.143 (251.142)	2.58 (CH ₃)	179.0 [=C(CH ₃)]
4j ^e	40 ⁱ	82	oil	281.152 (281.153)	2.57 (CH ₃)	179.8 [=C(CH ₃)]
4k	72	96	96–100	251.141 (251.142)	2.67 (CH ₃)	178.7 [=C(CH ₃)]
4l ^f	48	90	oil	265.161 (265.158)	2.67 (CH ₃)	178.7 [=C(CH ₃)]
4m ^g	48	97	oil	279.174 (279.174)	2.67 (CH ₃)	179.9 [=C(CH ₃)]
4n ^h	72	96	oil	295.168 (295.173)	2.67 (CH ₃)	179.8 [=C(CH ₃)]

^a Compounds 4a and 4d are described in ref 2. All the compounds 4a–n exhibit in the IR (KBr) spectrum an absorption between 2220 and 2240 (CN) cm^{-1} . Satisfactory elemental analyses were obtained for 4b,c,h,k (C, ± 0.30 ; H, ± 0.10 ; N, ± 0.30). ^{b–h} ^1H NMR δ : ^b 1.13 (d, 3 H, J = 6.1 Hz, CH₃); ^c 3.13 (s, 3 H, OCH₃); ^d 1.16 [d, 3 H, J = 5.9 Hz, NCH(CH₃)]; ^e 3.26 (s, 3 H, OCH₃); ^f 0.88 [d, 3 H, J = 6.3 Hz, NCH(CH₃)]; ^g 0.73 (t, 3 H, J = 7.2 Hz, CH₂CH₃); ^h 3.14 (s, 3 H, OCH₃). ⁱ Reaction performed at 40 °C.

duced pressure, and the residue was purified by column chromatography (SiO₂; CH₂Cl₂). The reaction times and yields are summarized in Table I.

1,2,3,3a-Tetrahydro-3a-methylpyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5b): oil; ^1H NMR (80 MHz) δ 7.3–6.95 (m, 2 H, Ar H), 6.85–6.5 (m, 2 H, Ar H), 3.42 (s, 2 H, ArCH₂), 3.75–3.2 (m, 2 H, NCH₂), 2.5–2.0 (m, 4 H, CH₂), 1.26 (s, 3 H, CH₃); ^{13}C NMR δ 141.1 (s, C-9a), 129.0, 117.2, and 113.2 (d, Ar C), 115.4, 114.3, and 112.9 (s, C-5a and CN), 63.8 [s, NC(CH₃)], 47.6 (t, NCH₂), 38.7 [s, C(CN)₂], 37.0 (t, ArCH₂), 34.4 and 21.6 (t, CH₂), 20.6 (q, CH₃); IR (KBr) 2245 (CN) cm^{-1} ; mass spectrum, m/e

237.126 (M^+ , calcd for C₁₅H₁₅N₃ 237.127).

1,2,3,3a-Tetrahydro-3a-(methoxymethyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5c): mp 135–136 °C (MeOH); ^1H NMR (80 MHz) δ 7.3–7.0 and 6.9–6.6 (m, 4 H, Ar H), 3.75–3.5 (m, 2 H, NCH₂), 3.47 (s, 2 H, ArCH₂), 3.37 (s, 2 H, CH₂OCH₃), 3.30 (s, 3 H, OCH₃), 2.65–2.0 (m, 4 H, CH₂); ^{13}C NMR δ 141.4 (s, C-9a), 129.1, 129.0, 117.6, and 113.8 (d, Ar C), 114.9, 114.6, and 113.7 (s, C-5a and CN), 73.7 (t, CH₂OCH₃), 66.1 [s, NC-(CH₂OCH₃)], 59.6 (q, OCH₃), 48.8 (t, NCH₂), 36.9 [s, C(CN)₂], 34.7, 33.5, and 22.3 (t, ArCH₂ and CH₂); IR (KBr) 2240 (CN) cm^{-1} ; mass spectrum, m/e 267.138 (M^+ , calcd 267.137). Anal. Calcd

for $C_{16}H_{17}N_3O$ (M_r 267.331): C, 71.89; H, 6.41; N, 15.72. Found: C, 72.10; H, 6.73; N, 15.61.

cis-(±)-1,2,3,3a-Tetrahydro-1-(methoxymethyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (6c): mp 156–158 °C (MeOH); 1H NMR (200 MHz) δ 7.25–7.1 and 7.05–7.0 (m, 2 H, Ar H), 6.8–6.7 (m, 2 H, Ar H), 4.2–4.05 [m, 1 H, NCH(CH₂OCH₃)], 3.91 [dd, 1 H, J = 3.7 and 8.2 Hz, NCH (bridgehead)], 3.53 (ABX, 1 H, J = 9.5 and 5.2 Hz, CHHOCH₃) and 3.49 (ABX, 1 H, J = 9.5 and 4.0 Hz, CHHOCH₃), 3.46 (s, 2 H, ArCH₂), 3.35 (s, 3 H, OCH₃), 2.7–2.5 (m, 1 H, CHH), 2.4–2.2 (m, 2 H, CH₂), 2.05–1.95 (m, 1 H, CHH); ^{13}C NMR δ 141.2 (s, C-9a), 129.3, 129.1, 117.7, and 112.4 (d, Ar C), 115.4 and 114.6 (s, CN), 113.4 (s, C-5a), 73.5 (t, CH₂OCH₃), 62.6 [d, NCH (bridgehead)], 60.3 [d, NCH(CH₂OCH₃)], 59.3 (q, OCH₃), 38.2 (t, ArCH₂), 33.6 [s, C(CN)₂], 28.5 and 25.9 (t, CH₂); IR (KBr) 2250 (CN) cm^{-1} ; mass spectrum, m/e 267.138 (M^+ , calcd 267.137). Anal. Calcd for $C_{16}H_{17}N_3O$ (M_r 267.331): C, 71.89; H, 6.41; N, 15.72. Found: C, 71.97; H, 6.60; N, 15.75.

trans-(±)-1,2,3,3a-Tetrahydro-1-(methoxymethyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (7c): mp 99–101 °C (MeOH); 1H NMR (200 MHz) δ 7.25–7.15 and 7.1–7.0 (m, 2 H, Ar H), 6.8–6.65 (m, 2 H, Ar H), 4.1–3.95 [m, 1 H, NCH(CH₂OCH₃)], 3.73 [dd, 1 H, J = 5.2 and 9.1 Hz, NCH (bridgehead)], 3.57 (dd, 1 H, J = 9.4 and 2.5 Hz, CHHOCH₃), 3.55 and 3.44 (AB q, 2 H, J = 16.4 Hz, ArCH₂), 3.36 (s, 3 H, OCH₃), 3.14 (dd, 1 H, J = 9.4 and 9.4 Hz, CHHOCH₃), 2.45–2.35 (m, 1 H, CHH), 2.3–2.15 (m, 3 H, CHH and CH₂); ^{13}C NMR δ 140.8 (s, C-9a), 129.3, 128.9, 117.8, and 112.2 (d, Ar C), 114.7 and 113.1 (s, CN), 114.4 (s, C-5a), 71.7 (t, CH₂OCH₃), 63.0 [d, NCH (bridgehead)], 59.1 (q, OCH₃), 56.9 [d, NCH(CH₂OCH₃)], 38.0 (t, ArCH₂), 34.8 [s, C(CN)₂], 27.6 and 27.0 (t, CH₂); IR (KBr) 2250 (CN) cm^{-1} ; mass spectrum, m/e 267.138 (M^+ , calcd 267.137). Anal. Calcd for $C_{16}H_{17}N_3O$ (M_r 267.331): C, 71.89; H, 6.41; N, 15.72. Found: C, 72.04; H, 6.60; N, 15.75.

2,3,4,4a-Tetrahydro-4a-methyl-1H-benzo[c]quinolizine-5,5(6H)-dicarbonitrile (5e): mp 131–132 °C (MeOH); 1H NMR (80 MHz) δ 7.35–6.65 (m, 4 H, Ar H), 3.9–3.55 (m, 1 H, NCH_{eq}H_{ax}), 3.52 (s, 2 H, ArCH₂), 3.0–2.5 (m, 1 H, NCH_{eq}H_{ax}), 2.3–1.5 (m, 6 H, CH₂), 1.23 (s, 3 H, CH₃); ^{13}C NMR 143.7 (s, C-10a), 129.2, 128.8, 119.0, and 113.4 (d, Ar C), 115.1 (s, C-6a), 115.0 (s, CN), 57.2 [s, NC(CH₃)], 43.1 [s, C(CN)₂], 42.6 (t, NCH₂), 34.3, 24.9, and 19.4 (t, CH₂), 35.2 (t, ArCH₂), 13.8 (q, CH₃); IR (KBr) 2255 (CN) cm^{-1} ; mass spectrum, m/e 251.142 (M^+ , calcd 251.142). Anal. Calcd for $C_{16}H_{17}N_3$ (M_r 251.332): C, 76.46; H, 6.82; N, 16.72. Found: C, 76.39; H, 6.81; N, 16.32.

2,3,4,4a-Tetrahydro-4a-ethyl-1H-benzo[c]quinolizine-5,5(6H)-dicarbonitrile (5f): mp 132–133.5 °C (MeOH); 1H NMR (80 MHz) δ 7.3–6.95 (m, 2 H, Ar H), 6.9–6.7 (m, 2 H, Ar), 3.8–3.4 (m, 1 H, NCH_{eq}H_{ax}), 3.51 (s, 2 H, ArCH₂), 3.05–2.7 (m, 1 H, NCH_{eq}H_{ax}), 2.55–1.25 (m, 8 H, CH₂, CH₂CH₃), 1.03 (t, 3 H, J = 7.5 Hz, CH₂CH₃); ^{13}C NMR δ 143.9 (s, C-10a), 129.0, 128.8, 118.8, and 113.3 (d, Ar C), 115.7, 115.5, and 115.4 (s, C-6a and CN), 59.7 [s, NC(CH₂CH₃)], 42.6 [s, C(CN)₂], 43.5 (t, NCH₂), 31.9, 24.6, and 19.6 (t, CH₂), 34.4 (t, ArCH₂), 23.1 (t, CH₂CH₃), 9.3 (q, CH₂CH₃); IR (KBr) 2255 (CN) cm^{-1} ; mass spectrum, m/e 265.158 (M^+ , calcd 265.158). Anal. Calcd for $C_{17}H_{19}N_3$ (M_r 265.359): C, 76.95; H, 7.22; N, 15.83. Found: C, 76.74; H, 7.37; N, 15.80.

2,3,4,4a-Tetrahydro-4a-(methoxymethyl)-1H-benzo[c]quinolizine-5,5(6H)-dicarbonitrile (5g): mp 109–110 °C (MeOH); 1H NMR (200 MHz) δ 7.25–7.15 and 7.1–7.0 (m, 2 H, Ar H), 6.9–6.75 (m, 2 H, Ar H), 3.8–3.65 (m, 1 H, NCH_{eq}H_{ax}), 3.62 and 3.37 (AB q, 2 H, J = 10.3 Hz, CH₂OCH₃), 3.50 (s, 2 H, ArCH₂), 3.30 (s, 3 H, OCH₃), 2.9–2.6 (m, 3 H, NCH_{eq}H_{ax}, CH₂), 2.0–1.5 (m, 4 H, CH₂); ^{13}C NMR δ 143.7 (s, C-10a), 129.2, 128.7, and 118.8 (d, Ar C), 115.8 (s, C-6a), 115.6 and 114.7 (s, CN), 112.8 (d, Ar C), 68.0 (t, CH₂OCH₃), 59.4 (q, OCH₃), 59.0 [s, NC(CH₂OCH₃)], 43.1 (t, NCH₂), 41.1 [s, C(CN)₂], 34.2 (t, ArCH₂), 30.3, 24.5, and 19.0 (t, CH₂); IR (KBr) 2255 (CN) cm^{-1} ; mass spectrum, m/e 281.150 (M^+ , calcd 281.152). Anal. Calcd for $C_{17}H_{19}N_3O$ (M_r 281.359): C, 72.57; H, 6.81; N, 14.92. Found: C, 72.29; H, 6.76; N, 14.90.

trans-(±)-2,3,4,4a-Tetrahydro-1-(methoxymethyl)-1H-benzo[c]quinolizine-5,5(6H)-dicarbonitrile (6g): mp 168–170 °C (MeOH); 1H NMR (200 MHz) δ 7.3–7.15 (m, 1 H, Ar H), 7.05–6.9 (m, 2 H, Ar H), 6.85–6.7 (m, 1 H, Ar H), 4.35–4.2 [m, 1 H, NCH(CH₂OCH₃)], 3.55 (ABX, 1 H, J = 9.3 and 9.3 Hz,

CHHOCH₃) and 3.37 (ABX, 1 H, J = 9.3 and 4.1 Hz, CHHOCH₃), 3.44 and 3.40 (AB q, 2 H, J = 9.9 Hz, ArCH₂), 3.6–3.45 [m, 1 H, NCH (bridgehead)], 3.30 (s, 3 H, OCH₃), 2.4–2.25 (m, 1 H, CHH), 2.1–1.6 (m, 5 H, CHH and CH₂); ^{13}C NMR δ 142.9 (s, C-10a), 129.7, 129.4, and 118.8 (d, Ar C), 115.3 (s, C-6a), 114.9 and 113.3 (s, CN), 113.6 (d, Ar C), 69.2 (t, CH₂OCH₃), 59.1 (q, OCH₃), 56.7 [d, NCH (bridgehead)], 52.0 [d, NCH(CH₂OCH₃)], 37.6 [s, C(CN)₂], 37.5 (t, ArCH₂), 29.7, 25.3, and 18.2 (t, CH₂); IR (KBr) 2230 (CN) cm^{-1} ; mass spectrum, m/e 281.149 (M^+ , calcd 281.152). Anal. Calcd for $C_{17}H_{19}N_3O$ (M_r 281.359): C, 72.57; H, 6.81; N, 14.92. Found: C, 72.29; H, 6.76; N, 14.71.

trans-(±)-1,2,3,3a-Tetrahydro-5-methylpyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5h): mp 130–131 °C (MeOH); 1H NMR δ 7.35–7.1 (m, 2 H, Ar H), 6.85–6.7 (m, 1 H, Ar H), 6.6–6.5 (m, 1 H, Ar H), 3.88 [dd, 1 H, J = 8.2 and 5.9 Hz, NCH (bridgehead)], 3.6–3.3 [m, 3 H, NCH₂, ArCH(CH₃)], 2.6–2.45 and 2.35–1.95 (m, 4 H, CH₂), 1.75 [d, 3 H, J = 6.8 Hz, ArCH(CH₃)]; ^{13}C NMR δ 142.0 (s, C-9a), 129.0, 126.8, 117.5, and 111.9 (d, Ar C), 118.7 (s, C-5a), 115.2 (s, CN), 63.0 (d, NCH), 48.1 (t, NCH₂), 42.3 [s, C(CN)₂], 41.1 [d, ArCH(CH₃)], 30.3 and 22.7 (t, CH₂), 16.1 (q, CH₃); IR (KBr) 2245 (CN) cm^{-1} ; mass spectrum, m/e 237.128 (M^+ , calcd for $C_{15}H_{15}N_3$ 237.127). No satisfactory elemental analysis could be obtained.

trans-(±)-1,2,3,3a-Tetrahydro-3a,5-dimethylpyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5i): mp 112–113 °C (MeOH); 1H NMR (200 MHz) δ 7.35–7.15 (m, 2 H, Ar H), 6.85–6.7 and 6.65–6.55 (m, 2 H, Ar H), 3.75–3.6 (m, 1 H, NCH_{eq}H_{ax}), 3.55–3.35 [m, 2 H, NCH_{eq}H_{ax}, ArCH(CH₃)], 2.55–2.1 (m, 4 H, CH₂), 1.76 [d, 3 H, J = 6.8 Hz, ArCH(CH₃)], 1.28 [s, 3 H, NC(CH₃)]; ^{13}C NMR δ 140.8 (s, C-9a), 128.9, 127.7, 117.3, and 113.1 (d, Ar C), 118.4 (s, C-5a), 115.5 (s, CN), 64.0 [s, NC(CH₃)], 47.9 (t, NCH₂), 37.6 (t, CH₂), 36.6 [d, ArCH(CH₃)], 21.5 (t, CH₂), 21.0 [q, NC(CH₃)], 16.8 [q, ArCH(CH₃)]; IR (KBr) 2240 (CN) cm^{-1} ; mass spectrum, m/e 251.142 (M^+ , calcd 251.142). Anal. Calcd for $C_{16}H_{17}N_3$ (M_r 251.332): C, 76.46; H, 6.82; N, 16.72. Found: C, 76.13; H, 7.00; N, 16.81.

trans-(±)-1,2,3,3a-Tetrahydro-3a-(methoxymethyl)-5-methylpyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5j): mp 139–141 °C (MeOH); 1H NMR (200 MHz) δ 7.3–7.15 (m, 2 H, Ar H), 6.85–6.75 and 6.7–6.6 (m, 2 H, Ar H), 3.8–3.65 (m, 1 H, NCH_{eq}H_{ax}), 3.6–3.3 [m, 4 H, NCH_{eq}H_{ax}, CH₂OCH₃, ArCH(CH₃)], 3.29 (s, 3 H, OCH₃), 2.7–2.1 (m, 4 H, CH₂), 1.76 (d, 3 H, J = 6.7 Hz, CH₃); ^{13}C NMR δ 141.7 (s, C-9a), 128.8, 127.1, 117.7, and 113.7 (d, Ar C), 119.2 (s, C-5a), 114.9 and 113.4 (s, CN), 73.7 (t, CH₂OCH₃), 66.2 [s, NC(CH₂OCH₃)], 59.6 (q, OCH₃), 49.0 (t, NCH₂), 45.5 [s, C(CN)₂], 36.4 (d, ArCH(CH₃)), 34.1 and 22.1 (t, CH₂), 16.8 (q, CH₃); IR (KBr) 2260 (CN) cm^{-1} ; mass spectrum, m/e 281.152 (M^+ , calcd 281.153). Anal. Calcd for $C_{17}H_{19}N_3O$ (M_r 281.358): C, 72.57; H, 6.81; N, 14.93. Found: C, 72.92; H, 6.94; N, 15.19.

(1 α ,3 α ,5 β)-1,2,3,3a-Tetrahydro-1-(methoxymethyl)-5-methylpyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (6j): mp 123–125 °C (MeOH); 1H NMR (200 MHz) δ 7.25–7.1 and 6.85–6.65 (m, 4 H, Ar H), 4.2–4.1 (m, 1 H, NCH(CH₂OCH₃)), 4.01 [dd, 1 H, J = 8.1 and 4.0 Hz, NCH (bridgehead)], 3.53 (ABX, 1 H, J = 12.5 and 5.4 Hz, CHHOCH₃), 3.48 (ABX, 1 H, J = 12.5 and 3.7 Hz, CHHOCH₃), 3.44 [q, 1 H, J = 6.8 Hz, ArCH(CH₃)], 3.34 (s, 3 H, OCH₃), 2.65–2.5 (m, 1 H, CHH), 2.4–2.2 (m, 2 H, CH₂), 2.1–1.9 (m, 1 H, CHH), 1.75 [d, 3 H, J = 6.8 Hz, ArCH(CH₃)]; ^{13}C NMR δ 141.2 (s, C-9a), 128.6, 126.9, 117.7, and 112.1 (d, Ar C), 118.6 (s, C-5a), 115.5 and 112.3 (s, CN), 73.4 (t, CH₂OCH₃), 62.7 [d, NCH (bridgehead)], 60.5 [d, NCH(CH₂OCH₃)], 59.3 (q, OCH₃), 42.1 [s, C(CN)₂], 40.8 [d, ArCH(CH₃)], 28.8 and 26.0 (t, CH₂), 15.9 (q, CH₃); IR (KBr) 2240 (CN) cm^{-1} ; mass spectrum, m/e 281.152 (M^+ , calcd 281.153). Anal. Calcd for $C_{17}H_{19}N_3O$ (M_r 281.358): C, 72.57; H, 6.81; N, 14.93. Found: C, 72.21; H, 6.92; N, 15.00.

(1 α ,3 α ,5 α)-1,2,3,3a-Tetrahydro-1-(methoxymethyl)-5-methylpyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (7j): mp 139–141 °C (MeOH); 1H NMR (200 MHz) δ 7.25–7.15 and 6.8–6.1 (m, 4 H, Ar H), 4.2–4.05 [m, 1 H, NCH(CH₂OCH₃)], 3.83 [dd, 1 H, J = 9.2 and 4.5 Hz, NCH (bridgehead)], 3.57 (dd, 1 H, J = 9.3 and 2.7 Hz, CHHOCH₃), 3.50 [q, 1 H, J = 6.9 Hz, ArCH(CH₃)], 3.36 (s, 3 H, OCH₃), 3.15 (dd, 1 H, J = 9.3 and 9.3 Hz, CHHOCH₃), 2.5–2.35 (m, 1 H, CHH), 2.3–2.0 (m, 3 H, CHH and CH₂), 1.74 [d, 3 H, J = 6.9 Hz, ArCH(CH₃)]; ^{13}C NMR δ 140.8

(s, C-9a), 129.3, 128.9, 117.9, and 112.2 (d, Ar C), 119.9 (s, C-5a), 114.8 and 113.1 (s, CN), 71.8 (t, CH_2OCH_3), 63.1 [d, NCH (bridgehead)], 59.1 (q, OCH_3), 56.9 [d, NCH(CH_2OCH_3)], 43.8 [s, $\text{C}(\text{CN})_2$], 42.6 [d, ArCH(CH_3)], 28.0 and 27.2 (t, CH_2), 16.9 (q, CH_3); IR (KBr) 2240 (CN) cm^{-1} ; mass spectrum, m/e 281.151 (M^+ , calcd 281.153). Anal. Calcd for $\text{C}_{17}\text{H}_{19}\text{N}_3\text{O}$ (M , 281.358): C, 72.57; H, 6.81; N, 14.93. Found: C, 72.37; H, 6.77; N, 14.63.

trans-(±)-2,3,4,4a-Tetrahydro-6-methyl-1H-benzo[c]-quinolizine-5,5(6H)-dicarbonitrile (5k): mp 145.5–146.5 °C (MeOH); ^1H NMR (200 MHz) δ 7.25–7.1 and 6.95–6.8 (m, 4 H, Ar H), 3.99 (br d, 1 H, $J = 11.3$ Hz, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$), 3.52 [q, 1 H, $J = 6.9$ Hz, ArCH(CH_3)], 3.33 [dd, 1 H, $J = 11.3$ and 3.2 Hz, NCH (bridgehead)], 2.68 [ddd, 1 H, $J = 11.3$, 11.3, and 3.4 Hz, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$], 2.45–2.3 (m, 1 H, CHH), 2.1–1.4 (m, 5 H, CHH and CH_2), 1.75 [d, 3 H, $J = 6.9$ Hz, ArCH(CH_3)]; ^{13}C NMR δ 144.6 (s, C-10a), 128.8 and 126.9 (d, Ar C), 121.3 (s, C-6a), 119.5 and 113.9 (d, Ar C), 115.0 and 112.3 (s, CN), 60.1 (d, NCH), 48.1 (t, NCH_2), 46.2 [s, $\text{C}(\text{CN})_2$], 39.9 [d, ArCH(CH_3)], 30.0, 24.8, and 22.7 (t, CH_2), 16.6 (q, CH_3); IR (KBr) 2260 (CN) cm^{-1} ; mass spectrum, m/e 251.143 (M^+ , calcd 251.142). Anal. Calcd for $\text{C}_{18}\text{H}_{17}\text{N}_3$ (M , 251.332): C, 76.40; H, 6.84; N, 16.86. Found: C, 76.46; H, 6.82; N, 16.72.

trans-(±)-2,3,4,4a-Tetrahydro-4a,6-dimethyl-1H-benzo[c]-quinolizine-5,5(6H)-dicarbonitrile (5l): mp 146–148 °C (MeOH); ^1H NMR (200 MHz) δ 7.3–6.7 (m, 4 H, Ar H), 3.69 (br d, 1 H, $J = 11.9$ Hz, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$), 3.46 [q, 1 H, $J = 6.6$ Hz, ArCH(CH_3)], 2.65 [ddd, 1 H, $J = 11.9$, 11.9, and 3.5 Hz, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$], 2.2–1.5 (m, 6 H, CH_2), 1.76 [d, 3 H, $J = 6.6$ Hz, ArCH(CH_3)], 1.21 [s, 3 H, NC(CH_3)]; ^{13}C NMR δ 143.5 (s, C-10a), 128.8 and 127.2 (d, Ar C), 121.0 (s, C-6a), 119.1 (d, Ar C), 115.2 and 113.5 (s, CN), 113.3 (d, Ar C), 57.8 [s, NC(CH_3)], 51.8 [s, $\text{C}(\text{CN})_2$], 42.8 (t, NCH_2), 36.0 [d, ArCH(CH_3)], 35.7, 24.9, and 19.2 (t, CH_2), 17.3 and 14.1 (q, CH_3); IR (KBr) 2240 (CN) cm^{-1} ; mass spectrum, m/e 265.156 (M^+ , calcd 265.158). Anal. Calcd for $\text{C}_{17}\text{H}_{19}\text{N}_3$ (M , 265.359): C, 76.95; H, 7.22; N, 15.83. Found: C, 76.66; H, 7.34; N, 15.95.

trans-(±)-2,3,4,4a-Tetrahydro-4a-ethyl-6-methyl-1H-benzo[c]-quinolizine-5,5(6H)-dicarbonitrile (5m): mp 113.5–114.5 °C (MeOH); ^1H NMR (200 MHz) δ 7.3–7.1 and 6.9–6.7 (m, 4 H, Ar H), 3.64 (br d, 1 H, $J = 11.8$ Hz, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$), 3.46 [q, 1 H, $J = 7.0$ Hz, ArCH(CH_3)], 2.82 [ddd, 1 H, $J = 11.8$, 11.8, and 3.6 Hz, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$], 2.6–2.4 (m, 1 H, CHH), 2.15–1.5 (m, 7 H, CHH, CH_2 , CH_2CH_3), 1.77 [d, 3 H, $J = 7.0$ Hz, ArCH(CH_3)], 1.00 (t, 3 H, $J = 7.5$ Hz, CH_2CH_3); ^{13}C NMR δ 143.7 (s, C-10a), 128.7 and 128.6 (d, Ar C), 121.2 (s, C-6a), 118.8 (d, Ar C), 115.5 and 114.2 (s, CN), 113.1 (d, Ar C), 59.8 [s, NC(C_2H_5)], 51.1 [s, $\text{C}(\text{CN})_2$], 43.5 (t, NCH_2), 36.0 [d, ArCH(CH_3)], 32.2, 24.5, 23.2, and 19.5 (t, CH_2), 17.4 [q, ArCH(CH_3)], 9.2 (q, CH_2CH_3); IR (KBr) 2245 (CN) cm^{-1} ; mass spectrum, m/e 279.170 (M^+ , calcd 279.173). Anal. Calcd for $\text{C}_{18}\text{H}_{21}\text{N}_3$ (M , 279.386): C, 77.38; H, 15.04; N, 7.58. Found: C, 77.44; H, 15.12; N, 7.59.

trans-(±)-2,3,4,4a-Tetrahydro-4a-(methoxymethyl)-6-methyl-1H-benzo[c]-quinolizine-5,5(6H)-dicarbonitrile (5n): mp 153–155 °C (MeOH); ^1H NMR (200 MHz) δ 7.3–7.15 and 6.9–6.75 (m, 4 H, Ar H), 3.75–3.5 (m, 1 H, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$), 3.62 and 3.37 (AB q, 2 H, $J = 10.3$ Hz, CH_2OCH_3), 3.47 [q, 1 H, $J = 6.7$ Hz, ArCH(CH_3)], 3.29 (s, 3 H, OCH_3), 2.85–2.6 (m, 3 H, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$, CH_2), 2.1–1.5 (m, 4 H, CH_2), 1.79 [d, 3 H, $J = 6.7$ Hz, ArCH(CH_3)]; ^{13}C NMR δ 143.5 (s, C-10a), 128.6 and 127.2 (d, Ar C), 121.5 (s, C-6a), 118.9 (d, Ar C), 114.6 and 114.1 (s, CN), 112.7 (d, Ar C), 68.5 (t, CH_2OCH_3), 59.4 (q, OCH_3), 59.3 [s, NC(CH_2OCH_3)], 49.6 [s, $\text{C}(\text{CN})_2$], 43.4 (t, NCH_2), 35.7 [d, ArCH(CH_3)], 30.8, 24.4, and 19.0 (t, CH_2), 17.1 (q, CH_3); IR (KBr) 2260 (CN) cm^{-1} ; mass spectrum, m/e 295.171 (M^+ , calcd 295.169). Anal. Calcd for $\text{C}_{18}\text{H}_{21}\text{N}_3\text{O}$ (M , 295.385): C, 73.19; H, 7.17; N, 14.22. Found: C, 72.99; H, 7.08; N, 13.98.

(1 α ,3 α ,5 β)-(±)-2,3,4,4a-Tetrahydro-1-(methoxymethyl)-6-methyl-1H-benzo[c]-quinolizine-5,5(6H)-dicarbonitrile (6n): mp 135–136 °C (MeOH); ^1H NMR (200 MHz) δ 7.3–7.1 and 7.0–6.75 (m, 4 H, Ar H); 4.4–4.25 [m, 1 H, NCH(CH_2OCH_3)], 3.7–3.6 [m, 1 H, NCH (bridgehead)], 3.53 (ABX, 1 H, $J = 9.0$ and 9.0 Hz, CHHOCH_3), 3.34 (ABX, 1 H, $J = 9.0$ and 4.0 Hz, CHHOCH_3), 3.43 [q, 1 H, $J = 6.8$ Hz, ArCH(CH_3)], 3.28 (s, 3 H, OCH_3), 2.45–2.3 and 2.15–2.0 (m, 2 H, CH_2), 1.9–1.7 (m, 4 H, CH_2), 1.73 [d, 3 H, $J = 6.8$ Hz, 3 H, ArCH(CH_3)]; ^{13}C NMR δ 143.0 (s, C-10a), 129.1 and 126.9 (d, Ar C), 120.9 (s, C-6a), 118.8 and 113.5

(d, Ar C), 115.0 and 112.0 (s, CN), 68.9 (t, CH_2OCH_3), 59.0 (q, OCH_3), 57.2 (d, NCH), 52.0 [d, NCH(CH_2OCH_3)], 46.2 [s, $\text{C}(\text{CN})_2$], 39.8 [d, ArCH(CH_3)], 30.1, 25.3, and 18.2 (t, CH_2), 16.1 (q, CH_3); IR (KBr) 2250 (CN) cm^{-1} ; mass spectrum, m/e 295.166 (M^+ , calcd 295.168). Anal. Calcd for $\text{C}_{18}\text{H}_{21}\text{N}_3\text{O}$ (M , 295.385): C, 73.19; H, 7.17; N, 14.22. Found: C, 73.03; H, 7.30; N, 14.29.

trans-(±)- and cis-(±)-1,2,3,3a-tetrahydro-5-(4-methylphenyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5o and 8o): ratio 82:18. Major isomer (5o): mp 207–208 °C (MeOH); ^1H NMR (200 MHz) δ 7.4–7.15 (m, 5 H, Ar H), 6.8–6.6 (m, 3 H, Ar H), 4.55 [s, 1 H, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 4.03 [dd, 1 H, $J = 7.8$ and 6.3 Hz, NCH (bridgehead)], 3.65–3.4 (m, 2 H, NCH_2), 2.39 (s, 3 H, CH_3), 2.60–2.0 (m, 4 H, CH_2); ^{13}C NMR δ 142.7 (s, C-9a), 138.9 and 133.1 (s, Ar C), 129.5 and 129.2 (d, Ar C), 118.3 (s, C-5a), 117.2 (d, Ar C), 114.5 and 112.6 (s, CN), 112.1 (d, Ar C), 63.5 [d, CH($\text{C}_6\text{H}_4\text{CH}_3$)], 53.1 (d, NCH), 47.9 (t, NCH_2), 43.3 [s, $\text{C}(\text{CN})_2$], 30.1 and 22.6 (t, CH_2), 21.2 (q, CH_3); IR (KBr) 2260 (CN) cm^{-1} ; mass spectrum, m/e 313.153 (M^+ , calcd 313.157). Anal. Calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3$ (M , 313.403): C, 80.49; H, 6.11; N, 13.41. Found: C, 80.39; H, 6.11; N, 13.42.

trans-(±)- and cis-(±)-1,2,3,3a-tetrahydro-3a-methyl-5-(4-methylphenyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5p and 8p): ratio 98:2. Major isomer (5p): mp 184–186 °C (MeOH); ^1H NMR (200 MHz) δ 7.5–7.1 (m, 5 H, Ar H), 6.95–6.5 (m, 3 H, Ar H), 4.52 [s, 1 H, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 3.8–3.4 (m, 2 H, NCH_2), 2.6–2.1 (m, 4 H, CH_2), 2.38 (s, 3 H, ArCH $_3$), 1.39 [s, 3 H, CH_3 (bridgehead)]; ^{13}C NMR δ 141.7 (s, C-9a), 138.8 and 133.7 (s, Ar C), 129.9, 129.4, and 129.0 (d, Ar C), 117.5 (s, C-5a), 117.0 (d, Ar C), 114.9 and 113.7 [s, $\text{C}(\text{CN})_2$], 113.2 (d, Ar C), 64.9 [s, NC(CH_3)], 48.9 [d, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 47.7 (t, NCH_2), 37.6 and 21.5 (t, CH_2), 21.2 and 20.8 (q, CH_3); IR (KBr) 2255 (CN) cm^{-1} ; mass spectrum, m/e 327.168 (M^+ , calcd 327.173). Anal. Calcd for $\text{C}_{22}\text{H}_{21}\text{N}_3$ (M , 327.413): C, 80.70; H, 6.46; N, 12.83. Found: C, 80.34; H, 6.39; N, 12.71.

trans-(±)- and cis-(±)-1,2,3,3a-tetrahydro-3a-(methoxymethyl)-5-(4-methylphenyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (5q and 8q): ratio 95:5. Major isomer (5q): mp 222–223 °C (MeOH); ^1H NMR (200 MHz) δ 7.4–7.2 (m, 5 H, Ar H), 6.85–6.6 (m, 3 H, Ar H), 4.56 [s, 1 H, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 3.85–3.7 (m, 1 H, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$), 3.61 and 3.46 (AB q, 2 H, $J = 10.1$ Hz, CH_2OCH_3), 3.6–3.4 (m, 1 H, $\text{NCH}_{\text{eq}}\text{H}_{\text{ax}}$), 3.36 (s, 3 H, OCH_3), 2.65–2.15 (m, 4 H, CH_2), 2.39 (s, 3 H, CH_3); ^{13}C NMR δ 142.0 (s, C-9a), 138.9 and 133.7 (s, Ar C), 129.9, 129.4, and 129.0 (d, Ar C), 118.3 (s, C-5a), 117.3 (d, Ar C), 114.3 and 114.0 (s, CN), 113.8 (d, Ar C), 73.7 (t, CH_2OCH_3), 67.1 [s, NC(CH_2OCH_3)], 59.7 (q, CH_2OCH_3), 48.8 (t, NCH_2), 48.7 [d, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 46.3 [s, $\text{C}(\text{CN})_2$], 34.2 and 22.3 (t, CH_2), 21.2 (q, $\text{C}_6\text{H}_4\text{CH}_3$); IR (KBr) 2260 (CN) cm^{-1} ; mass spectrum, m/e 357.186 (M^+ , calcd 357.184). Anal. Calcd for $\text{C}_{23}\text{H}_{23}\text{N}_3\text{O}$ (M , 357.456): C, 77.28; H, 6.49; N, 11.75. Found: C, 77.01; H, 6.53; N, 11.71.

(1 α ,3 α ,5 β)-(±)- and (1 α ,3 α ,5 α)-(±)-1,2,3,3a-tetrahydro-1-(methoxymethyl)-5-(4-methylphenyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (6q and 9q): ratio 96:4. Major isomer (6q): mp 192–193 °C (MeOH); ^1H NMR (200 MHz) δ 7.4–7.15 (m, 5 H, Ar H), 6.8–6.6 (m, 3 H, Ar H), 4.54 [s, 1 H, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 4.3–4.15 [m, 1 H, NCH(CH_2OCH_3)], 4.19 [dd, 1 H, $J = 3.6$ and 8.3 Hz, NCH (bridgehead)], 3.58 (ABX, 1 H, $J = 9.5$ and 5.5 Hz, CHHOCH_3), 3.52 (ABX, 1 H, $J = 9.5$ and 3.8 Hz, CHHOCH_3), 3.36 (s, 3 H, OCH_3), 2.40 (s, 3 H, CH_3), 2.7–2.2 (m, 3 H, CH_2 and CHH), 2.15–1.95 (m, 1 H, CHH); ^{13}C NMR δ 141.9 (s, C-9a), 139.0 and 133.2 (s, Ar C), 129.8, 129.6, and 129.0 (d, Ar C), 118.1 (s, C-5a), 117.3 (d, Ar C), 114.8 and 113.0 (s, CN), 112.4 (d, Ar C), 73.3 (t, CH_2OCH_3), 63.94 [d, NCH (bridgehead)], 60.4 [d, NCH(CH_2OCH_3)], 59.3 (q, CH_2OCH_3), 53.1 [d, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 43.1 [s, $\text{C}(\text{CN})_2$], 28.7 and 26.1 (t, CH_2), 21.2 (q, $\text{C}_6\text{H}_4\text{CH}_3$); IR (KBr) 2255 (CN) cm^{-1} ; mass spectrum, m/e 357.178 (M^+ , calcd 357.184). Anal. Calcd for $\text{C}_{23}\text{H}_{23}\text{N}_3\text{O}$ (M , 357.456): C, 77.28; H, 6.49; N, 11.75. Found: C, 77.08; H, 6.46; N, 11.69.

(1 α ,3 α ,5 α)-(±)- and (1 α ,3 α ,5 β)-(±)-1,2,3,3a-tetrahydro-1-(methoxymethyl)-5-(4-methylphenyl)pyrrolo[1,2-a]quinoline-4,4(5H)-dicarbonitrile (7q and 10q): ratio 97:3. Major isomer (7q): mp 170–173 °C (MeOH); ^1H NMR (200 MHz) δ 7.4–7.1 (m, 5 H, Ar H), 6.8–6.6 (m, 3 H, Ar H), 4.58 [s, 1 H, ArCH($\text{C}_6\text{H}_4\text{CH}_3$)], 4.1–3.9 [m, 2 H, NCH(CH_2OCH_3) and NCH (bridgehead)], 3.66 (dd, 1 H, $J = 9.3$ and 2.6 Hz, CHHOCH_3), 3.23 (dd, 1 H, $J = 9.3$ and 9.3 Hz, CHHOCH_3), 3.39 (s, 3 H, OCH_3),

2.5–2.0 (m, 4 H, CH₂), 2.31 (s, 3 H, CH₃); ¹³C NMR δ 141.8 (s, C-9a), 139.0 and 133.4 (s, Ar C), 130.3, 129.5, and 129.0 (d, Ar C), 119.3 (s, C-5a), 117.7 (d, Ar C), 114.3 and 112.5 (s, CN), 112.0 (d, Ar C), 72.1 (t, CH₂OCH₃), 64.0 [d, NCH (bridgehead)], 59.1 (q, OCH₃), 57.1 and 53.2 [d, NCH(CH₂OCH₃) and ArCH(C₆H₄CH₃)], 44.6 [s, C(CN)₂], 28.0 and 26.9 (t, CH₂), 21.2 (q, CH₃); IR (KBr) 2260 (CN) cm⁻¹; mass spectrum, *m/e* 357.189 (M⁺, calcd 357.184). Anal. Calcd for C₂₃H₂₃N₃O (M_r 357.456): C, 77.28; H, 6.49; N, 11.75. Found: C, 76.91; H, 6.58; N, 11.71.

X-ray Structure Determination of 6n. The crystal structure of **6n** was determined by X-ray diffraction. Crystal data C₁₈H₂₁N₃O: triclinic, space group *P* $\bar{1}$; *a* = 15.031 (1) Å, *b* = 8.190 (2) Å, *c* = 6.938 (1) Å, α = 79.86 (1)°, β = 78.55 (1)°, γ = 94.82 (1)°; *V* = 817.7 (3) Å³; *Z* = 2, *d*_{calcd} = 1.20 g cm⁻³; μ = 0.7 cm⁻¹. Reflections were measured in the ω/2θ scan mode, using graphite monochromated Mo Kα radiation [scan width (ω) 1.40 + 0.6 tan θ]. The structure was solved by direct methods and refined with full-matrix least-squares methods. A total of 1583 reflections with *F*_o² > 3σ(*F*_o²) was used in the refinement. The number of parameters refined was 284 [scale factor, extinction parameter, positional parameters of all atoms, and thermal parameters (isotropic for H atoms, anisotropic for others)]. The final *R* factors were *R* = 3.6%, *R*_w = 4.6%. All calculations were done with SDP.²⁴

Acknowledgment. This investigation was supported by the Netherlands Foundation of Chemical Research

(24) Structure Determination Package; Frenz, B. A. and Associates Inc., College Station, TX, and Enraf-Nonius, Delft, 1983.

(SON) with financial aid from the Netherlands Organization for Scientific Research (NWO).

Registry No. **1a**, 446-52-6; **1b**, 445-27-2; **1c**, 68295-43-2; **2a**, 123-75-1; **2b**, 765-38-8; **2c**, 76946-27-5; **2d**, 110-89-4; **2e**, 109-05-7; **2f**, 1484-80-6; **2g**, 117607-54-2; **3a**, 58028-74-3; **3b**, 117677-92-6; **3c**, 117677-93-7; **3d**, 34595-26-1; **3e**, 117607-55-3; **3f**, 117607-56-4; **3g**, 117607-57-5; **3h**, 70243-84-4; **3i**, 117677-94-8; **3j**, 117677-95-9; **3k**, 39911-06-3; **3l**, 117607-58-6; **3m**, 117607-59-7; **3n**, 117607-60-0; **3o**, 117607-61-1; **3p**, 117607-62-2; **3q**, 117607-63-3; **4a**, 87698-95-1; **4b**, 107743-55-5; **4c**, 107743-57-7; **4d**, 87698-96-2; **4e**, 117607-64-4; **4f**, 117607-65-5; **4g**, 117607-66-6; **4h**, 117607-67-7; **4i**, 117607-68-8; **4j**, 107797-45-5; **4k**, 117607-69-9; **4l**, 117607-70-2; **4m**, 117607-71-3; **4n**, 117607-72-4; **5a**, 117607-73-5; **5b**, 107743-56-6; **5c**, 107743-58-8; **5d**, 117607-74-6; **5e**, 117607-75-7; **5f**, 117607-76-8; **5g**, 117607-77-9; **5h**, 117607-78-0; **5i**, 117677-96-0; **5j**, 107797-46-6; **5k**, 117607-79-1; **5l**, 117607-80-4; **5m**, 117607-81-5; **5n**, 117607-82-6; **5o**, 117607-83-7; **5p**, 117607-84-8; **5q**, 117607-85-9; **6c**, 107743-59-9; **6g**, 117607-86-0; **6j**, 107797-47-7; **6n**, 117607-87-1; **6p**, 117607-88-2; **6q**, 117607-89-3; **7c**, 107743-64-6; **7g**, 117607-90-6; **7j**, 107797-48-8; **7n**, 117677-97-1; **7p**, 117677-98-2; **7q**, 117677-99-3; **8o**, 117607-91-7; **8p**, 117607-92-8; **8q**, 117607-93-9; **9q**, 117678-00-9; **10q**, 117678-01-0; 2,4'-H₃CC₆H₄CH(OH)C₆H₄F, 117607-94-0; (4-methylphenyl)magnesium bromide, 4294-57-9; malononitrile, 109-77-3.

Supplementary Material Available: Tables of positional and thermal parameters, bond distances and bond angles for **6n** (6 pages). Ordering information is given on any current masthead page.

Stereochemical Aspects of the "tert-Amino Effect". 2. Enantio- and Diastereoselectivity in the Synthesis of Quinolines, Pyrrolo[1,2-*a*]quinolines, and [1,4]Oxazino[4,3-*a*]quinolines

Walter H. N. Nijhuis,[†] Willem Verboom,[†] A. Abu El-Fadl,^{†,§} Gerrit J. van Hummel,[†] and David N. Reinhoudt^{*,†}

Laboratories of Organic Chemistry and Chemical Physics, University of Twente, 7500 AE Enschede, The Netherlands

Received June 20, 1988

Thermal isomerization of the optically pure 2-vinyl-*N,N*-dialkylanilines, with a methyl or an ethyl substituent (R²) at the α-position of the *N,N*-dialkyl moiety, **4a** (R¹ = H), **4b** (R¹ = CH₃), **5a** (R¹ = H), and **6a** (R¹ = H), affords *enantioselectively* the optically pure pyrrolo[1,2-*a*]quinolines **7a** and **7b** and the [1,4]oxazino[4,3-*a*]quinoline **11**, with the methyl or ethyl substituent (R²) at the bridgehead carbon atom, and the quinoline **13**, respectively. The optical purity of these quinoline derivatives was determined by ¹H NMR spectroscopy in the presence of chiral shift reagents. Heating of the optically pure analogues in which R² is a methoxymethyl group (**4c** and **4d**) in refluxing 1-butanol yields, besides the compounds **7c,d** with the methoxymethyl group at the bridgehead carbon atom, also the regioisomers **8c,d** and **9c,d** that are enantiomerically pure. Mixtures of the diastereomers **12a,b** and **14a,b** were obtained by cyclization of compound **5b**, with a 3-ethylmorpholinyl group, and of the acyclic amine **6b**, respectively, in refluxing 1-butanol. The compounds **12a** and **14a,b** were proven enantiomerically pure. The configuration of the compounds **8**, **9**, **12**, and **14** was determined by X-ray analysis [(±)-**12a**] and ¹H NMR and ¹H NOE difference spectroscopy. These results provide conclusive evidence for the mechanism of these cyclization reactions, which are further examples of the "tert-amino effect". The effect of substituents on the enantio- and diastereoselectivity of the cyclization is discussed.

Introduction

In a previous paper on the C–C bond formation via the "tert-amino effect", we have described the influence of steric and electronic effects of substituents on the regioselectivity of the cyclization of 2-vinyl-*N,N*-dialkyl-

anilines, yielding pyrrolo[1,2-*a*]quinolines and benzo[*c*]quinolizines.¹

As a further extension we have investigated the possible synthesis of optically pure quinoline derivatives by thermal conversion of optically pure 2-vinyl-*N,N*-dialkylanilines. Moreover, cyclization of chiral 2-vinyl-*N,N*-dialkylanilines could provide conclusive evidence for the proposed

[†] Laboratory of Organic Chemistry.

[‡] Laboratory of Chemical Physics.

[§] Present address: Physics Department, Faculty of Science, Assiut University, Assiut, Egypt.

(1) Nijhuis, W. H. N.; Verboom, W.; Abu El-Fadl, A.; Harkema, S.; Reinhoudt, D. N. *J. Org. Chem.*, preceding paper in this issue.