Differential Metal Ion Stabilization of Reactants and Transition States in the Transacylation of Crown Ether Aryl Acetates

Roberta Cacciapaglia,[†] Arie R. van Doorn,[‡] Luigi Mandolini,^{*,†} David N. Reinhoudt,^{*,†} and Willem Verboom[‡]

Contribution from Centro CNR Meccanismi di Reazione and Dipartimento di Chimica, Università la Sapienza, 00185 Roma, Italy, and Laboratory of Organic Chemistry, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands. Received June 24, 1991

Abstract: Transfer to methoxide ion of the acetyl group of a series of crown ether aryl acetates ranging from 2-AcO-15C4 to 2-AcO-27C8 is accelerated by alkali and alkaline earth metal ions. The magnitude of the rate enhancements is dependent on the substrate-cation combination, the divalent metals being in general much stronger catalysts than the monovalent metals. A maximum catalytic effect of 760 is observed in the Ba²⁺-catalyzed reaction of 2-AcO-18C5. The dependence of the observed second-order rate constants (k_{obsd}) on metal ion concentration was quantitatively accounted for in terms of 1:1 associations of the metal ions with reactants and transition states. A definite contribution from an additional reaction path involving two metal ions was detected in the cases of the Na⁺- and K⁺-catalyzed reactions of 2-AcO-24C7 and of the K⁺- and Rb⁺-catalyzed reactions of 2-AcO-27C8. A comparative analysis of structure and metal ion effects on the strength of binding of metal ions with reactants and transition states indicates that transition-state stabilization by metal ions occurs through cooperation of coordination to the oxygen donors of the bridge and electrostatic binding to the negative charge being transferred from methoxide ion to the carbonyl oxygen.

Introduction

We have reported¹ that the polyether bridges of 2-AcO-15C4 and 2-AcO-21C6 greatly increase the extent of catalysis brought about by alkali and alkaline-earth cations on rates of acetyl transfer from phenyl acetate to methoxide ion in methanol (eq 1). The

observed effects are highly dependent on cation nature and bridge length. On account of their inherent simplicity, these reactions lent themselves to a thorough kinetic investigation from which the strength of interaction of metal ions with transition states and reactants could be carefully measured. A fundamental understanding of the way in which additional binding energy rendered available by some extra group is utilized in the catalyst-substrate (CS) and in the transition state (CT[‡]) complexes² is crucial for the understanding of the mechanism of catalysis. Therefore, we have extended our investigations over a wide range of ring sizes, with the aim of obtaining a complete picture of differential metal ion stabilization of reactants and transition states through complexation. Binding of metal ions by crown ether hosts is known to be highly dependent on the type of metal ion and the macrocyclic ring size.³ Accordingly, variations in the length of the polyether bridge spanning the 1,3-xylyl unit are expected to produce structured reactivity patterns.

In this paper, we report an extensive kinetic investigation of the effect of monovalent (Na, K, Rb, and Cs) and divalent (Sr, Ba) metal ions on the acetyl-transfer reaction between methoxide ion and a series of crown ether acetates ranging from a ring size of 15 to a ring size of 27, in MeOH at 25.0 °C. 2,6-Bis[(2-ME)M]PA was also included as an acyclic model compound.

Results

Acyl-transfer reactions were carried out by reacting very dilute (ca. 0.1 mM) substrate solutions with excess Me₄NOMe (1–10 mM), either in the absence or presence of metal bromides. These were added over a wide concentration range. The above conditions are strictly comparable to those of the previous investigation,¹ with the only significant difference being that in the present work the total salt concentration is adjusted to 0.15 M with Et₄NBr in order to keep the ionic strength essentially constant and to simplify the treatment of the reactions of the divalent metals.⁴

Spectral changes at 300 nm due to the aryl oxide products showed close adherence to first-order kinetics in all cases, with stable and reproducible infinity readings corresponding to virtually quantitative reactions. The results are expressed as second-order rate constants calculated from the observed pseudo-first-order rate constants and the stoichiometric concentration of methoxide ion. A complete list of rate data is given in the supplementary material. Plots of k_{obsd}/k_0 against concentration of added salt are shown in Figure 1 for the monovalent and in Figure 2 for the divalent ions. Here k_{obsd} is the second-order rate constant measured in the presence of added salt, and k_0 refers to reactions run under conditions where tetraalkylammonium ions are the only cations present in solution. Since the latter rate constants closely represent free methoxide reactivity,^{1,4} they provide a convenient reference for a quantitative evaluation of the effect of metal ions on transacylation rates.

It is apparent from Figures 1 and 2 that in all cases acetyl transfer to methoxide proceeds much faster in the presence of metal ions. The reactivity pattern is highly structured, with effects strongly dependent on the substrate-cation combination. In general, the magnitude of the observed rate enhancements increases on increasing metal ion concentration and eventually reaches a plateau value in many cases, but a reactivity drop is present in the high-concentration region of some of the rate profiles

(1) Cacciapaglia, R.; Lucente, S.; Mandolini, L.; van Doorn, A. R.; Reinhoudt, D. N.; Verboom, W. *Tetrahedron* 1989, 45, 5293–5304.

(2) Fersht, A. *Enzyme Structure and Mechanism*; W. H. Freeman and Co.: New York, 1985; Chapter 12.

(3) (a) Izatt, R. M.; Bradshaw, J. S.; Nielsen, S. A.; Lamb, J. D.; Christensen, J. J. *Chem. Rev.* 1985, 85, 271–339. (b) Inoue, Y.; Liu, Y.; Hakushi, T. *Cation Binding by Macrocycles*; Inoue, Y., Gokel, G. W., Eds.; Marcel Dekker: New York, 1990; p 1.

(4) Ercolani, G.; Mandolini, L. *J. Am. Chem. Soc.* 1990, 112, 423–427.

[†] Università la Sapienza.

[‡] University of Twente.

Figure 1. Effect of alkali metal bromides on the rate of the basic methanolysis of 2-AcO-[3(*n* + 1)]C*n* at 25 °C. The curves are calculated using either eq 3 or eq 10 and the parameters listed in Table I.

related to the divalent metals (Figure 2). Maximum observed accelerations range from 2 to 20 with the alkali ions and from 2 to 3 orders of magnitude with the alkaline-earth ions.

Unlike the larger alkali metal ions, lithium proved to be ineffective in all of the reactions investigated. The low solubility of calcium and magnesium methoxide prevented investigation of the effects of these ions.

In contrast with the behavior displayed by the ring compounds, the reaction of the open chain model 2,6-bis[(2-ME)M]PA is only slightly affected by alkali metal ions. Rate enhancements of less than 30 percent were observed in all cases at the highest attainable concentrations of added salts. With the divalent metals we observed significant rate enhancements (Figure 2), though much less pronounced than with the cyclic substrates.

Treatment of Rate Data

Alkali Metal Ions. We found that in all cases k_{obsd} varies in proportion to the concentration of added salt in the low-concentration region, but the apparent order in metal salt usually decreases on increasing concentration and eventually becomes zero (saturation kinetics) in many cases. This is clearly indicative of significant associations of the metal ions with reactant(s). Since to a close approximation the alkali metal methoxides, like the alkali metal bromides, are strong electrolytes in MeOH,⁵ the only

(5) Fernandez-Prini, R. *Physical Chemistry of Organic Solvent Systems*; Covington, A. K., Dickinson, T., Eds.; Plenum: London, 1973; p 568-571.

Figure 2. Effect of strontium (a) and barium (b) bromides on the rate of the basic methanolysis at 25 °C of 2,6-bis[(2-ME)M]PA (Δ), 2-AcO-15C4 ($\bullet$), 2-AcO-18C5 ($\blacktriangledown$), 2-AcO-21C6 ($\circ$), 2-AcO-24C7 ($\blacktriangle$), and 2-AcO-27C8 ($\square$). The curves are calculated using either eq 11 or eq 12 and the parameters listed in Table II.

reactant-state interaction to be taken into account is that with the crown ether substrate (eq 2). Treatment of rate data was

$$\frac{k_{\text{obsd}}}{k_0} = \frac{1 + K_T[M^+]}{1 + K_S[M^+]} \quad (3)$$

carried out as before^{1,4} by means of eq 3, which is a binding isotherm for the case of 1:1 stoichiometry, having the familiar form of a rectangular hyperbola.⁶ Equation 3, easily derived from transition-state theory,⁷ applies when besides the uncatalyzed reaction path proceeding through the rate-limiting transition state T^* , a metal ion assisted path whose transition state has the composition T^*M^+ contributes to the overall rate. The quantity K_T^*

$$\lim_{[M^+] \rightarrow \infty} \frac{k_{\text{obsd}}}{k_0} = \frac{k_{\text{cat}}}{k_0} = \frac{K_T^*}{K_S} \quad (5)$$

has the meaning of the equilibrium constant⁸ for binding of the metal ion to the transition state (eq 4), and k_{cat} is the second-order rate constant⁹ for reaction of methoxide ion with the substrate fully saturated with the metal ion. Equation 3 can be written in the equivalent form of eq 6.

$$k_{\text{obsd}} = \frac{k_0 + k_{\text{cat}}K_S[M^+]}{1 + K_S[M^+]} \quad (6)$$

The data related to the reactions of 2-AcO-15C4, 2-AcO-18C5, and 2-AcO-21C6 strictly fit to eq 3 (6), as shown by the curves drawn in Figure 1a–c, which were calculated by means of the parameters summarized in Table I.¹⁰ The situation with the larger macrocycles 2-AcO-24C7 and 2-AcO-27C8 is more complex, as

Table I. Alkali Metal Ion Assisted Acetyl Transfer in MeOH at 25 °C

	K_S (M^{-1})	K_T^* (M^{-1})	$K_T^*M^+$ (M^{-1})	k_{cat} ($M^{-1} s^{-1}$)	k_{cat}/k_0
2-AcO-15C4 ($k_0 = 9.63 \times 10^{-2} M^{-1} s^{-1}$)					
Na	$\leq 3^a$	3.9×10		> 1.2	> 12
K	11	3.2×10^2		2.8	29
Rb	13	3.0×10^2		2.2	23
Cs	10	1.7×10^2		1.6	17
2-AcO-18C5 ($k_0 = 6.61 \times 10^{-2} M^{-1} s^{-1}$)					
Na	5.1×10	4.5×10^2		0.58	8.8
K	1.8×10^3	3.6×10^4		1.30	20
Rb	2.2×10^3	1.5×10^4		0.45	6.8
Cs	1.7×10^3	4.9×10^3		0.19	2.9
2-AcO-21C6 ($k_0 = 6.66 \times 10^{-2} M^{-1} s^{-1}$)					
Na	5.2×10	9.8×10		0.12	1.8
K	7.2×10^2	4.5×10^3		0.42	6.3
Rb	1.3×10^3	1.2×10^4		0.61	9.2
Cs	1.8×10^3	2.2×10^4		0.81	12
2-AcO-24C7 ($k_0 = 6.72 \times 10^{-2} M^{-1} s^{-1}$)					
Na	9.3^b	4.1×10	8.3	0.30	4.5
K	3.2×10^2	5.5×10^2	6.2	0.12	1.8
Rb	4.8×10^2	1.2×10^3		0.17	2.5
Cs	7.1×10^2	2.9×10^3		0.27	4.0
2-AcO-27C8 ($k_0 = 6.28 \times 10^{-2} M^{-1} s^{-1}$)					
Na	9.8^b	4.8×10		0.31	4.9
K	1.2×10^2	3.2×10^2	10	0.17	2.7
Rb	1.4×10^2	3.8×10^2	3	0.17	2.7
Cs	2.2×10^2	5.7×10^2		0.16	2.5

^a See footnote 10. ^b From NMR titration. ^c From spectrophotometric titration.

shown by inspection of the pertinent rate profiles (Figure 1d,e). An undeniable tendency toward saturation is apparent in the Cs reactions of both substrates, as well as in the Rb reaction of 2-AcO-24C7. However, in the Rb reaction of 2-AcO-27C8 and in the K reactions of both substrates, saturation values are not reached in the high-concentration regions but, rather, regular rate increases with increasing metal ion concentration are observed. The quasi-linear dependence of k_{obsd} upon $[Na^+]$, which is apparent in the profile of the Na reaction of 2-AcO-24C7, would suggest at first sight a first-order dependence on metal ion, but we found that this is definitely not the case (vide infra). The relatively featureless profile of the Na reaction of 2-AcO-27C8 bears a close resemblance to the profiles of the K and Rb reactions of the same substrate, but again we realized that the resemblance is more apparent than real and that the rate equation for the Na reaction is different from that of the K and Rb reactions. Let us now summarize our analyses starting from the simplest cases.

The Cs reactions of 2-AcO-24C7 and 2-AcO-27C8 and the Rb reaction of 2-AcO-24C7 were analyzed in the usual way on the

(6) Connors, K. A. *Binding Constants*; Wiley Interscience: New York, 1987; Chapter 6.

(7) Hammett, L. P. *Physical Organic Chemistry*; McGraw-Hill: New York, 1970; pp 136–140, 212–213.

(8) K_T^* , which is defined in terms of concentrations, is related to the limiting value K_T^{*0} at zero salt concentration by means of the relationship $K_T^* = \gamma^2 K_T^{*0}$, where γ is the activity coefficient of singly charged ions at the ionic strength of the kinetic experiments. It should be noted that $K_S = K_S^0$, as based on the assumptions that activity coefficients of species of like charge are equal ($\gamma_{SM^+} = \gamma_{M^+}$) and that the activity coefficient of neutral species is unity ($\gamma_S = 1$).

(9) $k_{\text{cat}} = \gamma^2 k_{\text{cat}}^0$, where k_{cat}^0 refers to zero ionic strength (see footnote 8).

(10) The profile for the Na reaction of 2-AcO-15C4 (Figure 1a) is linear, with no tendency to saturation. This indicates that the extent to which the substrate combines with Na ion is too small to affect the kinetics, i.e., $K_S[M^+] \ll 1$. The data were fit to the simple equation $(k_{\text{obsd}}/k_0) = 1 + K_T^*[M^+]$. There was spectroscopic evidence obtained for a very weak association. Addition of $NaClO_4$ to a methanol solution of 2-AcO-15C4 caused a hypochromic effect which was significant at salt concentrations in the range 0.2–0.5 M, but negligibly small at lower concentrations. Although these data are not such as to allow a precise estimate of the equilibrium constant, they provide a rough indication that K_S is probably on the order of $2\text{--}3 M^{-1}$.

basis of the hypothesis of metal ion interactions of 1:1 stoichiometry. As shown in Figure 1d,e, the data are well-fit by eq 3. On the other hand, the data for the K reactions of 2-AcO-24C7 and 2-AcO-27C8 and for the Rb reaction of 2-AcO-27C8 gave a very poor fit to eq 3. Furthermore, we note that the order of catalytic efficiency at low metal ion concentration (<0.01 M) is $\text{Cs} > \text{Rb} > \text{K} > \text{Na}$ both with 2-AcO-24C7 and 2-AcO-27C8, but when the concentration is raised to 0.05–0.1 M, the order becomes $\text{Na} \approx \text{Cs} > \text{Rb} \approx \text{K}$ with 2-AcO-24C7 and $\text{K} > \text{Rb} > \text{Na} \approx \text{Cs}$ with 2-AcO-27C8. On the basis of the analogy of what has been observed in related cases,^{11,12} we can only attribute this behavior to the superposition of an additional reaction path that requires the involvement of two metal ions.

The kinetic equation that accounts for the additional operation of a two-metal-ion mechanism (eq 7) is easily obtained by expansion of eq 3. The quantities $K_{T^*M^+}$ and K_{SM^+} are defined in

$$\frac{k_{\text{obsd}}}{k_0} = \frac{1 + K_{T^*}[M^+](1 + K_{T^*M^+}[M^+])}{1 + K_S[M^+](1 + K_{SM^+}[M^+])} \quad (7)$$

eqs 8 and 9, respectively.¹³ Whenever there is insignificant formation of the initial-state 1:2 complex $S(M^*)_2$, i.e., when $K_{SM^+}[M^+] \ll 1$, eq 7 reduces to the simpler form of eq 10.

$$\frac{k_{\text{obsd}}}{k_0} = \frac{1 + K_{T^*}[M^+](1 + K_{T^*M^+}[M^+])}{(1 + K_S[M^+])} \quad (10)$$

It is apparent that eqs 7 and 10 are very adjustable in that they contain too many unknown parameters to be determined by curve fitting of the data. In order to avoid computational artifacts, we measured by independent techniques the equilibrium constants for 1:1 stoichiometry and tried to find some evidence for associations of 1:2 stoichiometry. This was done in most cases by means of ^1H NMR spectroscopy and occasionally by means of UV spectrophotometry. Chemical shifts or optical absorption changes upon complexation with the metal ions were in all cases very small. Nevertheless, meaningful titration curves were obtained from which reasonably accurate K_S values were calculated (Table I). In a control experiment, ^1H NMR analysis gave a K_S for binding of K^+ to 2-AcO-18C5 of $1.86 \times 10^3 \text{ M}^{-1}$, which compared extremely well with the value from the kinetic analysis.¹⁴ No evidence whatsoever was found for associations with 1:2 stoichiometry. Therefore, treatment of rate data was carried out by means of eq 10. The independently determined K_S values were used to develop our analysis as exemplified by the K^+ -catalyzed reaction of 2-AcO-27C8. On introduction of the known K_S values, eq 10 reduces to an equation which contains only two adjustable parameters. The solid curve drawn in Figure 3 is a plot of eq 10 calculated by means of the parameters listed in Table I. The dotted curve is a plot of the two-parameter eq 3. It is evident that the contribution resulting from incursion of the two- K^+ mechanism increases on increasing concentration, whereas the contribution becomes negligibly small when $[\text{K}^+]$ approaches zero.

The other cases where there was some evidence of the "disturbance" due to an additional metal ion were treated in the same way. The pertinent parameters are listed in Table I. It is worth noting that the finding that K_S for binding of Na^+ to 2-AcO-24C7 is not negligibly small definitely rules out the possibility that the reaction is simply first-order in $[\text{Na}^+]$ in the whole concentration range, as is found to be the case for the Na reaction of 2-AcO-15C4. The oddity of the seemingly linear rate profile (Figure 1d) is simply due to the fact that K_S and $K_{T^*M^+}$, which appear in the denominator and numerator, respectively, of eq 10,

Figure 3. The basic methanolysis of 2-AcO-27C8 as carried out in the presence of KBr: calculated with the inclusion of the two-metal-ion pathway (—); calculated with the exclusion of the two-metal-ion pathway (---).

are very similar to each other, so that a substantial cancellation occurs.

A last comment is devoted to the Na reaction of 2-AcO-27C8, whose rate data could be fit with good precision to eq 3. This indicates that the two-metal-ion pathway is unimportant in this case.

Alkaline Earth Metal Ions. Treatment of rate data as obtained in the presence of the divalent metal ions is less straightforward, because association with both methoxide ion and bromide ion is significant.¹⁴ If only interactions of 1:1 stoichiometry are involved, the functional relation between k_{obsd} and the total concentration of added salt, c_{salt} , is the following:

$$\frac{k_{\text{obsd}}}{k_0} = \frac{1 + K_{T^*}c_{\text{salt}}}{(1 + K_S c_{\text{salt}})(1 + K_{\text{MeO}} c_{\text{salt}})} \quad (11)$$

where binding to methoxide ion is accounted for by inclusion of a proper binomial in the denominator. It is likely that the metal ions are distributed among different species, namely, M^{2+} , $(\text{MBr})^+$, and MBr_2 , but the essentially constant concentration of bromide ions in the kinetic runs ensures invariance of the relative proportions of the various metal species in solution. Therefore, the K_{T^*} , K_S , and K_{MeO} quantities which appear in eq 11 have the meaning of conditional constants which strictly hold for solutions having the same ionic composition as those of the kinetic experiments.

When the substrate S does not bind metal ions to a significant extent, the binomial containing K_S is indistinguishable from unity, and eq 11 reduces to the simple form of eq 12.

$$\frac{k_{\text{obsd}}}{k_0} = \frac{1 + K_{T^*}c_{\text{salt}}}{1 + K_{\text{MeO}}c_{\text{salt}}} \quad (12)$$

We found that the dependence of k_{obsd} on metal ion concentration for the reactions of 2-AcO-15C4, 2-AcO-18C5, and the open chain model 2,6-bis[(2-ME)M]PA is well described by eq 12 (Figure 2), which means that binding of the divalent metal species by these substrates is negligible. The pertinent parameters are listed in Table II. It is worth noting that the K_{MeO} values are not only in fair agreement with each other, but also compare reasonably well with earlier determinations^{1,4} under similar, but not identical, conditions.

With the larger macrocycles, the complete form of eq 11 was used. In order to avoid application of a curve-fitting procedure to a three-parameter equation, the average K_{MeO} values as obtained from the reactions of 2,6-bis[(2-ME)M]PA, 2-AcO-15C4, and 2-AcO-18C5 were used as known values to leave only two unknown parameters to be evaluated by a curve fit of the data. Figure 2 shows the close adherence of the data to eq 11. The observed reactivity inversions are due to the fact that the squared term in c_{salt} in the denominator becomes significant at high salt concentrations. The effects are much more marked in the Ba reactions than in the Sr reactions, because the K_S values are much

(11) Mandolini, L.; Masci, B. *J. Am. Chem. Soc.* **1984**, *106*, 168–174.

(12) Antonini Vitali, C.; Masci, B. *Tetrahedron* **1989**, *45*, 2213–2222.

(13) $K_{T^*M^+} = K_{T^*}^0 + \gamma^2 K_{SM^+}^0$ (see footnote 8).

(14) The nearly perfect agreement may be incidental. A realistic estimate of the experimental uncertainty is $\pm 20\%$, or $\pm 10\%$ at best.

Figure 4. Binding of alkali metal ions to reactants (a) and transition states (b) in the basic methanolysis of 2-AcO-15C4 (●), 2-AcO-18C5 (▼), 2-AcO-21C6 (○), 2-AcO-24C7 (▲), and 2-AcO-27C8 (□).

is much greater than K_S , which is tantamount to saying that the metal ions are rate-enhancing in all of the studied reactions. The general behavior is fully consistent with the basic idea that transition-state stabilization by metal ions occurs through cooperation of two major modes of interaction, namely, electrostatic binding²⁰ to the negative charge being transferred from methoxide ion to the carbonyl oxygen and coordination to the oxygen donors of the bridge. In the absence of the bridge, as in the parent phenyl acetate,^{1,4} or when the bridge is replaced by two bidentate side arms to give the open chain reference compound 2,6-bis[(2-ME)M]PA, electrostatic binding to the monovalent metal ions is hardly appreciable. However, it is significant with the divalent metal ions. In fact, it is stronger than binding to the localized negative charge of methoxide ion, which has been interpreted^{1,4} as due to a chelate interaction involving both the methoxide oxygen and the carbonyl oxygen. It appears, therefore, that the transition state for transacylation may be reasonably pictured as a lariat-type macrocyclic ligand with a short side arm bearing a negatively charged donor site.

Alkali Metal Ions. A graphical comparison of K_S and K_T (Figure 4) shows that there is a substantial degree of resemblance between the two sets of values, including the fact that binding of metal ions, irrespective of the nature of the metal ion itself, is strongest at ring sizes of 18 and 21, in both the reactant and the transition states. A noticeable difference is the substantial selectivity toward K⁺ of the transition state of the reaction of 2-AcO-18C5 that the substrate itself lacks.

As for the important question of differential cation binding to transition state and reactants, we note that there is a general tendency for K_T to increase on increasing K_S , but a plot of log K_T against log K_S (not reported here) shows considerable scatter. This is not surprising, as linear free energy relationships, which have contributed so much to our present understanding of the relations between structure and reactivity, are more the exception than the rule in crown ether chemistry. As a consequence, the relation between catalysis and binding is far from being a simple one.

Since k_{cat} refers to substrates fully saturated by metal ion, only a fraction of the binding energy provided by the polyether bridges is utilized in catalysis,² namely, the differential binding energy with the transition state and with the reactant. Table I shows that, apart from such irregularities as the exceedingly low values of the Cs reaction of 2-AcO-18C5 and the Na reaction of 2-AcO-21C6, there is a general tendency for k_{cat} to decrease as the size of the ring becomes larger. This means that utilization of binding energy in catalysis by alkali metal ions generally decreases on increasing ring size, irrespective of the magnitude of the binding

energy itself. It is remarkable that the reaction of 2-AcO-15C4, which is a very weak cation binder in itself, exhibits the highest (k_{cat}/k_o) values. These considerations suggest that T^*M^+ can be described as an ion pair which is tight (strong electrostatic binding) in the lower terms of the series where the metal ion is obviously forced to sit very close to the reaction zone, but becomes increasingly more loose (weak electrostatic binding) when the ring size increases. In the limit of very large ring structures, complexation takes place far away from the reaction zone with little or no extra stabilization from the negative charge. Such a situation is met practically with the large ring substrates 2-AcO-24C7 and 2-AcO-27C8, for which the k_{cat}/k_o values are very low.

We found that an additional metal ion can be accommodated in the macrocyclic cavity of the transition state, giving rise to well-defined two-metal-ion pathways, which are kinetically detected in the Na and K reactions of 2-AcO-24C7 and in the K and Rb reactions of 2-AcO-27C8. The involvement of two metal ions in sufficiently large transition states having crown ether structure is well precedented. Following an earlier observation,¹¹ it has been recently demonstrated¹² that two K⁺ ions act as a template in the formation of benzo-27C9 and benzo-30C10. Furthermore, strong indication was obtained of the operation of the template effect of two Na⁺ ions in the formation of 24C8. Fully consistent with the above evidence are the findings that in the present system the metal ions that can be doubly accommodated in the cavity of 2-AcO-24C7 are Na⁺ and K⁺ and that with the larger 2-AcO-27C8 K⁺ still works, but the smaller Na⁺ is replaced by the larger Rb⁺. With Cs⁺, which is the largest cation, the effect is not appreciable. Thus, there appears to be a definite correlation between cation size and fit to the macrocyclic cavity. No doubt, there must be a steric compression of one metal ion against the other in the doubly occupied transition state, with the result that one of the metal ions is forced to be closer to the site where the negative charge is concentrated. We believe that this reinforcement of electrostatic binding relative to the loose ion pairing in T^*M^+ acts as a major driving force for the formation of transition-state complexes $T^*(M^+)_2$ of definite stability. Consistently, the neutral 24- and 27-membered macrocyclic reactants fail to give 1:2 complexes of definite stability.

Alkaline Earth Metal Ions. Table II shows that the K_T values increase regularly on increasing ring size up to a flat maximum centered at a ring size of 24 with both Sr²⁺ and Ba²⁺, with somewhat larger values being found for the latter ion in all cases. Again, the picture which emerges strongly indicates that metal ion stabilization of the transition state arises from cooperation between electrostatic binding and coordination to the polyether bridges, but the importance of the former is much greater with the divalent metal ions than with the monovalent ones. Interestingly, in the reactions of 2-AcO-15C4 and 2-AcO-18C5, K_S is negligibly small both with Sr²⁺ and with Ba²⁺. Here the only initial-state stabilization by metal ions which has to be paid for

(20) Fyles, T. M. *Cation Binding by Macrocycles*; Inoue, Y., Gokel, G. W., Eds.; Marcel Dekker: New York, 1990; p 203.

(in the denominator of eq 11) is that of methoxide ion. This means that, unlike the alkali metal ion reactions of the same substrates, the additional binding energy rendered available by the polyether bridges relative to the open chain model shows up in the transition states only and is utilized fully in catalysis. With the higher homologues, binding of metal ions to substrate becomes significant. This is particularly so in the case of Ba^{2+} . The result is that only a fraction of the available binding energy is utilized in catalysis. This is clearly shown, for example, by a comparison of the Ba reactions of 2-AcO-18C5 and 2-AcO-24C7. Although K_T^+ is larger with the latter, the former is more strongly catalyzed, as shown by the $(k_{\text{obsd}}/k_o)_{\text{max}}$ values listed in the last column of Table II. With the Sr^{2+} ion, the K_S values are so small that our rate measurements practically refer to conditions where the substrates are far from being saturated with metal ion. As a consequence, there is a nearly complete utilization of binding energy in catalysis, as clearly shown by the close parallelism between K_T^+ and $(k_{\text{obsd}}/k_o)_{\text{max}}$. This delicate interplay between transition state and substrate stabilization by metal ions explains well why Sr^{2+} is a more effective catalyst than Ba^{2+} in the reactions of the larger substrates, but less effective in the reactions of the smaller ones.

We finally note that the much greater catalytic power of the divalent as compared with the monovalent metal ions is easily understood in light of the above discussion and the K_S and K_T^+ data listed in Tables I and II. With the alkaline earth metal ions, there is not only a much larger electrostatic stabilization of the anionic transition state but also a better utilization of coordinative interactions with the neutral donors of the polyether bridges.

Experimental Section

^1H NMR spectra were recorded in CDCl_3 with a Bruker WP-80 spectrometer, and ^{13}C NMR spectra were recorded with a Nicolet MT 200 spectrometer. Mass spectra were obtained with a Varian MAT 311 A spectrometer. UV spectral measurements were carried out on an HP 8452 diode array spectrometer.

Materials. Dry methanol and a stock solution of tetramethylammonium methoxide in methanol were prepared and handled as previously reported.¹ Alkali and alkaline-earth bromides were from a previous investigation.¹ The bromide content of the methanolic solutions was checked by argentometric titration. Tetraethylammonium bromide (Aldrich, gold label) was crystallized from absolute ethanol-acetone (1:3.8, v/v) and dried under vacuum at 110 °C for 2 h. BuLi in hexane (Merck) was titrated with 2-butanol in the presence of 1,10-phenanthroline.²¹ Diethyl ether was stored over 4-Å molecular sieves. THF was distilled from sodium benzophenone ketyl before use. $\text{B}(\text{OMe})_3$ was also freshly distilled. 2-Bromo-1,3-bis(bromomethyl)benzene was prepared according to a literature procedure.²² Other chemicals were used as obtained.

2-AcO-[3(*n* + 1)]Cn (*n* = 4–8). These compounds were prepared (70–80% yield) by acetylation of the parent phenols²³ according to a published procedure.²⁴ Purification was carried out by flash chromatography on silica gel (230–400 mesh) using ethanol-chloroform (2:98, v/v) as eluent. Purity was found to be >99% by GLC or HPLC analysis. ^1H NMR, IR, and mass spectra were as expected.

1,3-Bis[(2-methoxyethoxy)methyl]-2-bromobenzene. To a refluxing stirred suspension of NaH (0.72 g, 0.024 mol, 80% in oil) in THF (125 mL) was added a solution of 2-methoxyethanol (1.67 g, 0.022 mol) in THF (25 mL) under nitrogen. After 1 h, a solution of 2-bromo-1,3-bis(bromomethyl)benzene (3.43 g, 0.010 mol) in THF (25 mL) was added and reacted to completion (TLC, SiO_2 , CHCl_3). Subsequently, water (100 mL) was added, and most of the THF was evaporated in vacuo. The aqueous layer was extracted with CHCl_3 (3 × 50 mL), whereupon the combined CHCl_3 layers were dried over MgSO_4 and concentrated in vacuo. The pure product was obtained as an oil after vacuum distillation: yield, 92%; bp 155–157 °C/0.06 mmHg; ^1H NMR δ 7.4–7.2 (m, 3 H, ArH), 4.65 (s, 4 H, ArCH_2), 3.8–3.6 (m, 8 H, OCH_2),

3.41 (s, 6 H, OCH_3); ^{13}C NMR δ 137.4 (s, C-2, 6), 127.7 (d, C-3, 5), 127.0 (d, C-4), 122.4 (s, C-1), 72.6, 71.8, 70.0 (t, OCH_2), 58.9 (q, OCH_3); mass spectrum, m/z 332.061 (M^+ , calcd for $\text{C}_{14}\text{H}_{21}\text{BrO}_4$ 332.062).

2,6-Bis[(2-methoxyethoxy)methyl]phenol. To a solution of 1,3-bis[(2-methoxyethoxy)methyl]-2-bromobenzene (3.02 g, 0.009 mol) in THF (15 mL) was added BuLi (6.7 mL, 1.5 M in hexane, 0.010 mol) dropwise at –78 °C under nitrogen. The mixture was stirred for 2 h, whereupon an excess of $\text{B}(\text{OMe})_3$ (3.2 mL, 0.027 mol) was added. After an additional 2 h of stirring at –78 °C, the reaction mixture was slowly warmed at room temperature. Subsequently, 4 M HCl (30 mL) was added, and the mixture was stirred for 15 h. The THF was distilled off, and the water phase was extracted with CHCl_3 (3 × 75 mL). The combined CHCl_3 layers were concentrated (ca. 50 mL) and extracted with 3 M NaOH (3 × 50 mL) and water (50 mL). The combined aqueous layers were washed with CHCl_3 (50 mL), acidified to pH 2, extracted with CHCl_3 (3 × 150 mL), and dried over MgSO_4 . After evaporation of the solvent, the residual colorless oil was dissolved in diethyl ether (100 mL), and 10% hydrogen peroxide (3.0 mL) and 10% NaHCO_3 (1.0 mL) were added. The reaction mixture was stirred for 6 h. Subsequently, 2 M HCl (30 mL) was added, and the mixture was extracted with CHCl_3 (3 × 125 mL). The combined CHCl_3 layers were washed with water (50 mL), dried over MgSO_4 , and concentrated in vacuo to obtain the pure product as a colorless oil: yield, 37%; ^1H NMR δ 7.8–7.7 (br s, 1 H, OH), 7.2–6.7 (m, 3 H, ArH), 4.69 (s, 4 H, ArCH_2), 3.8–3.5 (m, 8 H, OCH_2), 3.40 (s, 6 H, OCH_3); ^{13}C NMR δ 154.2 (s, C-1), 128.5 (d, C-3, 5), 123.8 (s, C-2, 6), 119.5 (d, C-4), 71.7, 70.2, 69.5 (t, OCH_2), 59.0 (q, OCH_3); mass spectrum, m/z 270.147 (M^+ , calcd for $\text{C}_{14}\text{H}_{22}\text{O}_5$ 270.147).

2,6-Bis[(2-methoxyethoxy)methyl]phenyl Acetate (2,6-Bis[(2-ME)-M]PA). The parent phenol was acetylated as above in 79% yield. Purification was carried out by flash chromatography on silica gel (230–400 mesh) using hexane-ethyl acetate (1:2, v/v) as eluent: ^1H NMR δ 7.4–7.2 (m, 3 H, ArH), 4.48 (s, 4 H, ArCH_2), 3.51 (s, 8 H, OCH_2), 3.36 (s, 6 H, OCH_3), 2.30 (s, 3 H, COCH_3); IR (CH_2Cl_2) 1763 cm^{-1} . Anal. Calcd for $\text{C}_{16}\text{H}_{24}\text{O}_6$: C, 61.52; H, 7.74. Found: C, 61.62; H, 7.81.

Equilibrium and Rate Measurements. The spectrophotometric determination of K_S was carried out as previously described.²⁵ The ^1H NMR determination was carried out in methanol- d_4 at 25.0 °C using tetramethylsilane and methylene chloride as internal standards. Calculated amounts of titrating solution of the alkali metal bromide were dispensed by microsyringe to a 1 mM substrate solution. After each addition, the chemical shift value of the methyl group was recorded after thermal equilibration. The data were fit by means of a nonlinear least-squares procedure to eq 13, where Δ is the chemical shift difference measured at the given metal ion concentration, and the subscript ∞ refers to substrate fully complexed by M^+ . The quantities Δ_∞ and K_S were treated as adjustable parameters.

$$\Delta = \frac{\Delta_\infty K_S [\text{M}^+]}{1 + K_S [\text{M}^+]} \quad (13)$$

Rate measurements were carried out as previously reported.^{1,4} In the runs carried out in the presence of divalent metal ions, the concentration of methoxide ion never exceeded 1 mM. This allowed us to neglect the fraction of M^{2+} sequestered by methoxide ion. Whenever required both in the equilibrium and rate measurements, the fraction of metal ion sequestered by the substrate was calculated by means of an iteration procedure, and the total metal ion concentration was corrected accordingly.

Computational Details. Nonlinear least-squares calculations were carried out using the program Sigma Plot (Jandel Scientific). Curve fitting was carried out on the logarithmic scale in order to give a more sound weight to the data.¹ Standard deviations from the regression line were in the range of 0.002–0.05 log unit (0.5–11%).

Acknowledgments. Work was carried out in the frame of EC joint projects, contracts No. ST2J-0215 and No. SC1*-0359. Thanks are due to EC for financial support, as well as to MURST for the part of work carried out in Rome.

Supplementary Material Available: Listings of second-order rate constants at the various concentrations of alkali and alkaline earth metal bromides for the reactions of compounds 2-AcO-[3(*n* + 1)]Cn (*n* = 4–8) and 2,6-bis[(2-ME)M]PA (4 pages). Ordering information is given on any current masthead page.

(21) Watson, S. C.; Eastman, J. F. *J. Organomet. Chem.* **1967**, *9*, 165–171.

(22) Newcomb, M.; Moore, S. S.; Cram, D. J. *J. Am. Chem. Soc.* **1977**, *99*, 6405–6410.

(23) Skowronska-Ptasinska, M.; Aarts, V. M. L. J.; Egberink, R. J. M.; van Eerden, J.; Harkema, S.; Reinhoudt, D. N. *J. Org. Chem.* **1988**, *53*, 5484–5491.

(24) Taylor, E. C.; McLay, G. W.; McKillop, A. *J. Am. Chem. Soc.* **1968**, *90*, 2422–2423.

(25) Ercolani, G.; Mandolini, L.; Masci, B. *J. Am. Chem. Soc.* **1981**, *103*, 7484–7489.