EARLY ECONOMIC EVALUATION

of technologies for emerging
interventions to personalize
breast cancer treatment

Anna Miquel Cases

**EARLY ECONOMIC EVALUATION OF TECHNOLOGIES FOR
EMERGING INTERVENTIONS TO PERSONALIZE BREAST
CANCER TREATMENT**

Anna Miquel Cases

Address of correspondence

Anna Miquel Cases
Molenwerf 4, F5
1014AG Amsterdam
The Netherlands

Copyright © Anna Miquel Cases, Amsterdam, 2016

All rights reserved. No part of this thesis may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage or retrieval system, without permission in writing from the author, or, when appropriate, from the publishers of the publications.

ISBN: 978-90-365-4055-1

Cover design: Anna Miquel Cases

Lay-out: Gildeprint

Printed by: Gildeprint

The research presented in this thesis was performed within the framework of the Center for Translational Molecular Medicine; project breast CARE.

The printing of this thesis was financially supported by:

- The Netherlands Cancer Institute.
- AMGEN B.V.
- Boehringer Ingelheim B.V.

**EARLY ECONOMIC EVALUATION OF TECHNOLOGIES FOR
EMERGING INTERVENTIONS TO PERSONALIZE BREAST
CANCER TREATMENT**

DISSERTATION

to obtain
the degree of doctor at the University of Twente,
on the authority of the rector magnificus
prof. dr. H. Brinksma,
on account of the decision of the graduation committee,
to be publicly defended
on Friday 1st April 2016 at 12.45h

by
Anna Miquel Cases
born on 15 December 1987
in Igualada, Spain

Supervisor

Prof. dr. W.H. van Harten (University of Twente)

Co-supervisor

Dr. L.M.G. Steuten (Fred Hutchinson Cancer Research Center)

Assessment committee:

Prof.dr. Th.A.J. Toonen (Chairman and secretary; University of Twente)

Prof.dr. R. Torenvlied (University of Twente)

Prof. dr. A.P.W.P. van Montfort (University of Twente)

Prof. dr. S. Siesling (University of Twente)

Dr. G.S. Sonke (Netherlands Cancer Institute)

Prof. dr. E. Buskens (University Medical Center Groningen)

Prof. dr. ir. J.J.M. van der Hoeven (Radboud University Medical Centre)

Paranymphs:

Jacobien Kieffer

Lisanne Hummel

Per al padrí, l' avi i el Josep

(to my granddads)

Table of contents

Part I Introduction

Chapter 1	General introduction	11
-----------	----------------------	----

Part II Predictive biomarkers: personalize systemic treatment

Chapter 2	(Very) early health technology assessment and translation of predictive biomarkers in breast cancer <i>Submitted for publication</i>	25
Chapter 3	Early stage cost-effectiveness analysis of a BRCA1-like test to detect triple negative breast cancers responsive to high dose alkylating chemotherapy <i>The Breast 2015, Aug;24(4):397-405.</i>	59
Chapter 4	Decisions on further research for predictive biomarkers of high dose alkylating chemotherapy in triple negative breast cancer: A value of information analysis <i>Value in Health 2016, in press</i>	79

Part III Imaging techniques: monitoring systemic treatment

Chapter 5	Imaging performance in guiding response to neoadjuvant therapy according to breast cancer subtypes: A systematic literature review <i>Submitted for publication</i>	107
Chapter 6	Exploratory cost-effectiveness analysis of response-guided neoadjuvant chemotherapy for hormone positive breast cancer patients <i>Accepted with minor revisions</i>	135
Chapter 7	Cost-effectiveness and resource use of implementing MRI-guided NACT in ER-positive/HER2-negative breast cancers <i>Revised submission</i>	163

Part IV Imaging techniques: screening for distant metastasis

Chapter 8	¹⁸ F-FDG PET/CT for distant metastasis screening in stage II/III breast cancer patients: A cost-effectiveness analysis from a British, US and Dutch perspective <i>Submitted for publication</i>	195
-----------	--	-----

Part V General discussion and Annex

Chapter 9	General discussion	235
Annex	Summary	255
	Samenvatting	259
	Acknowledgements	263
	List of publications	265
	Curriculum vitae	267

PART I

INTRODUCTION

CHAPTER 1

General introduction

Health technology assessment and economic evaluations

Health Technology Assessment (HTA) has been called “the bridge between evidence and policy-making” [1]. It is a discipline that aims to inform health-care decision-makers, on the properties, effects, and/or other impacts of health care technologies, as cited by the International Society of Technology Assessment in Health Care, 2002. The type of evidence typically considered in HTA includes safety, efficacy, cost and cost-effectiveness of a technology. However, with the increase of limitations in national budgets, partly motivated by the financial crisis of 2008, the increase in life expectancy due to presence of more effective health care interventions, and the ever-increasing costs of health care, cost-effectiveness considerations are becoming more central. In other words, there is greater awareness and urgency in considering whether money is wisely spent. As a consequence of this, in a growing number of countries cost-effectiveness (CE) is being used as a criterion for pricing and reimbursement decision-making [2–4] as well as a method to prioritize public and private resources into specific health problems and related interventions.

Economic Evaluations (EE) are the tool used to measure CE. They provide knowledge on the financial resources required to implement effective medical technologies and how money invested relates to outcomes achieved [5]. They are often performed in late stages of a technology’s development to demonstrate value for money [2,3] and thus facilitate its incorporation into the healthcare marketplace. The most recognized type of Economic Evaluation is Cost-Effectiveness Analysis (CEA). CEA compares the costs and the health effects of an intervention to assess the extent to which it can be regarded as providing value for money. The most common measures of health improvement are Life Years (LY) and Quality Adjusted Life Years (QALY) [5]. CEAs execution is often via health economic models, which provide of a framework to synthesize available clinical and economic evidence on the technology [6].

Early Economic Evaluations

A less common application of EE takes place in the early development of medical technologies. This application emerged in view of the high research and development costs of new technologies [7], especially in the late stages of development when patients have been included in trials [8]. The disadvantage of evaluations in later stages of development is that developers at this point have made a substantial capital investment in the technology, both in terms of developing the product itself and the evidence supporting its clinical role in care. Hence an unfavorable EE at this point creates severe problems for the manufacturer, particularly if the negative assessment is based on uncertainties regarding key aspects of performance (i.e., sensitivity) or the impact of the diagnostic on clinical outcomes versus alternatives. In fact, any factor that ‘drives’ an unfavorable assessment beyond price implies that the developer will have to make additional investments

in research, causing delays in access and further costs. Early EE could have identified this in a timely fashion, allowing technology developers to improve upon this and make sure a reasonable level of CE can be reached. Thus the aim of early EE is to inform strategic decisions in the early development stages, before embarking into phase II and III clinical trials.

Early EE can be used for many purposes [4]. The first application is to prioritize pipeline candidates for further research. A second application is to inform go/no-go decisions if results reveal that further development of the technology is not interesting from a health economic viewpoint. A third application is the guidance of product development by identifying economically favorable product characteristics. Lastly, early EE can be used to identify data gaps and optimal study designs to cover those data gaps. The differences between performing EE early versus late in the product development process are presented in table 1.

Health economic modeling is the central method to early EE. However, as early EE is a relatively new field, there is no unified framework on how to use health economic modeling alongside product development. Health economic modeling can be complemented by other type of HTA methods. Currently the use of these additional methods depends on the decision that needs to be informed [9]. While Bayesian techniques and Value of Information analysis (VOI) seem useful for updating information during research and development (R&D) and continuously informing decision-making [4,10], the headroom method can be valuable for informing the maximum reimbursable price of a technology [11]. Furthermore, scenario analysis can be used for trend extrapolation and for envisioning alternative paths into the future. Additionally, resource modeling analysis allows to quantitatively capture the resource implications of the future implementation a new technology in clinical practice [12].

Table 1: Key differences between early and mainstream EE, adapted from IJzerman et al [13].

Characteristics	Early economic evaluations	Mainstream economic evaluations
Objectives	Strategic R&D decision making	Reimbursement
	Preliminary market assessment	Pricing decisions
	Product development	
	Design clinical trials	
	Price determination	
Target informants	Manufacturer's	Policymakers
	Policymakers	Payers
Evidence	Elicitation from experts	Clinical trials
	Prior similar technologies	
	Animal studies	
	Small clinical studies	

Aim of this thesis

Even though the idea of starting EE early in the product life cycle has gained popularity in the past few years, its use in real-life situations is not fully exploited yet (VOI analysis [14,15], headroom analysis [11,16–18], scenario analysis [19], resource modeling analysis [20,21]). Therefore, this thesis contributes to this literature, particularly to that on early CEAs, VOI analysis and resource modeling analysis. We applied these methods to technologies for emerging breast cancer interventions with the aim to inform strategic decision-making in these technologies. This research was part of the Medical Technology Assessment work package of the Breast CARE project, funded by the public-privately Center for Translational Molecular Medicine consortium [22].

Breast cancer diagnosis and treatment

In Europe and worldwide, the incidence of breast cancer is between 25% and 29% of the total female population [23]. The last decades' decline in breast cancer mortality [24–26] is mainly caused by 1) the addition of drug treatment to the local treatment modalities of surgery and radiation therapy, and 2) earlier diagnosis as a result of breast cancer screening by mammography [27–31]. More recently, mortality rates have stabilized [26] and breast cancer remains the leading cause of cancer death in women [23]. Thereby, new approaches to its treatment are still needed.

Personalized medicine is an emerging approach to patient care, whose aim is to find the right treatment for the right patient at the right time [32]. It is an evolving field in medicine with many resources dedicated to searching for diagnostic, prognostic, and predictive technologies that can be used to guide clinical decision-making. It is expected that the translation of such technologies into routine clinical practice will improve current breast cancer survival rates.

Technologies for emerging breast cancer interventions

The Breast CARE project was our source for identifying technologies for emerging breast cancer interventions. The project was designed to discover and validate new technologies to personalize breast cancer treatment. A core idea was rapid translational research, so that scientific results could be applied as quickly as possible in actual patient care [22]. To stimulate this, the Neoadjuvant Chemotherapy (NACT) setting (where chemotherapy is given prior to surgery) was chosen as a research model. This had the advantage of providing an 'in vivo' model where new technology's effectiveness could be rapidly assessed. The project involved two types of technologies: predictive biomarkers and imaging techniques.

Predictive biomarkers: personalize systemic treatment

Predictive biomarkers are biological entities in a patient's body that associate with an outcome after a specific treatment is given and thus serve as a guide to personalize patients' treatment [33]. Although there is plenty of research on predictive biomarkers few of those are currently implemented in the daily practice, with ER/PR and HER2 being the main examples in breast cancer. Within the breast CARE project, three promising predictive biomarkers emerged: the BRCA1-like, the XIST, and the 53BP1. All three were determined to be predictive of high-dose alkylating chemotherapy [34,35] and are currently being validated in the framework of prospective or retrospective studies. These three biomarkers were involved as case studies in our early EE assessments.

Imaging techniques: monitoring systemic treatment

The combination of MRI and PET/CT as a tool to monitor treatment response during NACT was investigated in the Breast CARE project. Unfortunately, due to time constraints, we could not involve this project in this thesis. Yet as the idea of "response-guided NACT" seemed promising, we found alternative projects on this approach that could proportionate data within this thesis time-frame. One project explored the effectiveness of "response-guided NACT" by using MRI [36] and the other by using ultrasound [37]. These projects came from the Netherlands Cancer Institute (NKI), and the German Breast Group (GBG) in Germany respectively. These case studies were also involved in our early EE assessments.

Imaging techniques: screening for distant metastasis

The last intervention we assessed was the use of PET/CT for distant metastasis screening in stage II/III breast cancers. Although this intervention fall outside of the breast CARE scope, this research was motivated by the fact that PET/CT is a costly modality and emerging evidence suggests that it is expected to be more accurate than current standard imaging [38–42]. Therefore the interest in knowing its added value.

The technologies and emerging interventions that we assessed using early Economic Evaluation are presented in Figure 1.

VOI methods allow quantifying the uncertainty around cost-effectiveness estimates derived from early CEAs and valuing whether investing in additional research is worthwhile. The underlying principle of this framework is to compare the costs and benefits generated by gathering additional information with the costs of investing in further research [7,30]. The incorporation of VOI methods into early health economic models was done for two purposes. The first was to decide on whether investment in further research endeavors is worthwhile, and in case affirmative, the second was to identify the type of data and study designs that are most worthwhile to perform this additional research.

Resource modeling analysis is a method that typically falls outside the health economic evaluation scope but within the HTA framework. Resource modeling allows the quantitative capture of the resource implications of implementing a new technology in clinical practice [12]. As the ultimate goal of decision makers is implementation of cost-effective health-care interventions into routine clinical practice, this method can be of great help to health services planners who are challenged by implementation issues normally not addressed in CEAs.

Thesis outline

This thesis consists of three parts, distinguished by the type of technologies assessed: predictive biomarkers (chapter 2 – chapter 4), imaging techniques to monitor NACT response (chapter 5 – chapter 7) and imaging techniques to screen for distant metastasis (chapter 8). Specific research questions that are addressed in these chapters and that contribute to the overall aim of this thesis are presented here.

Predictive biomarkers: personalize systemic treatment

In **chapter 2** we discuss the current development stage of predictive biomarkers for NACT in breast cancer and suggest on ways to improve the translational process from a clinical, biological and HTA perspective. This chapter is motivated by the decision of Breast CARE to use the NACT setting as a model for biomarker discovery.

In **chapter 3** we estimate the expected cost-effectiveness of a biomarker strategy to personalize high dose alkylating chemotherapy in a subgroup of breast cancers (triple negative breast cancer). Furthermore, we determine the minimum prevalence of the biomarker and the minimum predictive value of its diagnostic test for the implementation of this biomarker strategy to be cost effective in clinical practice. This chapter illustrates the usefulness of threshold sensitivity analysis as a complementary method to early health economic modeling.

In **chapter 4** we present a model that estimates the expected cost-effectiveness of the various biomarker combinations that can be used to personalize high dose alkylating chemotherapy. We determine 1) the decision uncertainty in a possible adoption decision based on current information, 2) whether it is worth investing in further research to reduce decision uncertainty, and if so, 3) how to perform this research most efficiently. This paper is an illustration of the full VOI methodology based on an early health economic model.

Imaging techniques: monitoring systemic treatment

In **chapter 5** we present an overview of the literature on the performance of various imaging techniques in monitoring NACT response by taking into account the different breast cancer subtypes. This chapter is motivated by the emergence of literature highlighting the differences in imaging performance depending on subtype.

In **chapter 6** we present a model that compares the expected cost-effectiveness of a response-guided NACT using ultrasound in a subgroup of breast cancers (hormone-receptor positive patients). This paper illustrates the usefulness of early health economic modeling as a tool to estimate the expected cost-effectiveness of the technology as it were to be applied in clinical practice.

In **chapter 7** we present another model on the response-guided NACT approach, this time with MRI applied to another subgroup of breast cancers (ER-positive/HER2-negative patients). We estimated its expected cost-effectiveness and the resources required for its implementation compared to conventional NACT. This chapter illustrates the use of resource modeling analysis in addition to CEA considering a current and a full implementation scenario of response-guided NACT.

Imaging techniques: screening for distant metastasis

In **chapter 8** we calculate the expected cost-effectiveness of ^{18}F -FDG-PET/CT for distant metastasis screening in stage II-III patients from a perspective of the United Kingdom, the Netherlands, and the United States. This chapter illustrates the cost-effectiveness consequences of analyzing the same early health economic model from different country perspectives.

In **chapter 9** we conclude this thesis with a summary of answers to research questions, present a discussion on the possible methodological and treatment policy consequences and directions for future research.

References

- [1] R N Battista MJH. The evolving paradigm of health technology assessment: reflections for the millennium. *CMAJ* 1999;160(10):1464–7.
- [2] Johannesson M. Economic evaluation of drugs and its potential uses in policy making. *PharmacoEconomics* 1995;8:190–8.
- [3] Drummond MF. Economic evaluation of pharmaceuticals: science or marketing? *PharmacoEconomics* 1992;1:8–13.
- [4] Annemans L, Genesté B, Jolain B. Early modelling for assessing health and economic outcomes of drug therapy. *Value Health J Int Soc Pharmacoeconomics Outcomes Res* 2000;3:427–34. doi:10.1046/j.1524-4733.2000.36007.x.
- [5] Drummond et al. *Methods for the economic evaluation of health care programmes*. Oxford University Press; 2005.
- [6] Briggs et al. *Decision modelling for Health Economic Evaluation*. Oxford: Oxford university press; 2006.
- [7] Dorsey ER, de Roulet J, Thompson JP, Reminick JJ, Thai A, White-Stellato Z, et al. Funding of US biomedical research, 2003–2008. *JAMA* 2010;303:137–43. doi:10.1001/jama.2009.1987.
- [8] Moses H, Dorsey ER, Matheson DHM, Thier SO. Financial Anatomy of Biomedical Research. *JAMA* 2005;294:1333. doi:10.1001/jama.294.11.1333.
- [9] Markiewicz K, van Til JA, IJzerman MJ. MEDICAL DEVICES EARLY ASSESSMENT METHODS: SYSTEMATIC LITERATURE REVIEW. *Int J Technol Assess Health Care* 2014;30:137–46. doi:10.1017/S0266462314000026.
- [10] Miller P. Role of pharmacoeconomic analysis in R&D decision making: when, where, how? *PharmacoEconomics* 2005;23:1–12.
- [11] Cosh E, Girling A, Lilford R, McAteer H, Young T. Investing in new medical technologies: A decision framework. *J Commer Biotechnol* 2007;13:263–71. doi:10.1057/palgrave.jcb.3050062.
- [12] Thokala P, Dixon S, Jahn B. Resource Modelling: The Missing Piece of the HTA Jigsaw? *PharmacoEconomics* 2014. doi:10.1007/s40273-014-0228-9.
- [13] IJzerman MJ, Steuten LMG. Early assessment of medical technologies to inform product development and market access: a review of methods and applications. *Appl Health Econ Health Policy* 2011;9:331–47. doi:10.2165/11593380-000000000-00000.
- [14] Retèl VP, Grutters JPC, van Harten WH, Joore MA. Value of research and value of development in early assessments of new medical technologies. *Value Health J Int Soc Pharmacoeconomics Outcomes Res* 2013;16:720–8. doi:10.1016/j.jval.2013.04.013.
- [15] Steuten L, van de Wetering G, Groothuis-Oudshoorn K, Retèl V. A systematic and critical review of the evolving methods and applications of value of information in academia and practice. *PharmacoEconomics* 2013;31:25–48. doi:10.1007/s40273-012-0008-3.
- [16] McAteer H, Cosh E, Freeman G, Pandit A, Wood P, Lilford R. Cost-effectiveness analysis at the development phase of a potential health technology: examples based on tissue engineering of bladder and urethra. *J Tissue Eng Regen Med* 2007;1:343–9. doi:10.1002/term.36.
- [17] Cao Q, Postmus D, Hillege HL, Buskens E. Probability elicitation to inform early health economic evaluations of new medical technologies: a case study in heart failure disease management. *Value Health J Int Soc Pharmacoeconomics Outcomes Res* 2013;16:529–35. doi:10.1016/j.jval.2013.02.008.
- [18] Vaidya A, Joore MA, Cate-Hoek AJ ten, Cate H ten, Severens JL. Cost-effectiveness of risk assessment and tailored treatment for peripheral arterial disease patients. *Biomark Med* 2014;8:989–99. doi:10.2217/bmm.14.45.
- [19] Retèl VP, Joore MA, Linn SC, Rutgers EJT, van Harten WH. Scenario drafting to anticipate future developments in technology assessment. *BMC Res Notes* 2012;5:442. doi:10.1186/1756-0500-5-442.
- [20] Sharp L, Tilson L, Whyte S, Ceilleachair AO, Walsh C, Usher C, et al. Using resource modelling to inform decision making and service planning: the case of colorectal cancer screening in Ireland. *BMC Health Serv Res* 2013;13:105. doi:10.1186/1472-6963-13-105.
- [21] Stein ML, Rudge JW, Coker R, van der Weijden C, Krumkamp R, Hanvoravongchai P, et al. Development of a resource modelling tool to support decision makers in pandemic influenza preparedness: The AsiaFluCap Simulator. *BMC Public Health* 2012;12:870. doi:10.1186/1471-2458-12-870.

- [22] Cetner for Translational Molecular Medicine (CTMM) n.d. <http://www.ctmm.nl/>.
- [23] Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JWW, Comber H, et al. Cancer incidence and mortality patterns in Europe: Estimates for 40 countries in 2012. *Eur J Cancer* 2013;49:1374–403. doi:10.1016/j.ejca.2012.12.027.
- [24] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015: Cancer Statistics, 2015. *CA Cancer J Clin* 2015;65:5–29. doi:10.3322/caac.21254.
- [25] Cossetti RJD, Tyldesley SK, Speers CH, Zheng Y, Gelmon KA. Comparison of breast cancer recurrence and outcome patterns between patients treated from 1986 to 1992 and from 2004 to 2008. *J Clin Oncol Off J Am Soc Clin Oncol* 2015;33:65–73. doi:10.1200/JCO.2014.57.2461.
- [26] Allemani C, Weir HK, Carreira H, Harewood R, Spika D, Wang X-S, et al. Global surveillance of cancer survival 1995-2009: analysis of individual data for 25,676,887 patients from 279 population-based registries in 67 countries (CONCORD-2). *Lancet Lond Engl* 2015;385:977–1010. doi:10.1016/S0140-6736(14)62038-9.
- [27] Integraal Kankercentrum Nederland (IKNL). Breast Cancer Guideline, NABON 2012. n.d.
- [28] Smith RA, Manassaram-Baptiste D, Brooks D, Doroshenk M, Fedewa S, Saslow D, et al. Cancer screening in the United States, 2015: a review of current American cancer society guidelines and current issues in cancer screening. *CA Cancer J Clin* 2015;65:30–54. doi:10.3322/caac.21261.
- [29] England NCSP-PH. NHS Breast Screening Programme. 2014.
- [30] Force USPST. Screening for breast cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2009;151:716–26.
- [31] van Luijt PA, Fracheboud J, Heijnsdijk EAM, Heeten GJ den, de Koning HJ, National Evaluation Team for Breast Cancer Screening in Netherlands Study Group (NETB). Nation-wide data on screening performance during the transition to digital mammography: observations in 6 million screens. *Eur J Cancer Oxf Engl* 1990 2013;49:3517–25. doi:10.1016/j.ejca.2013.06.020.
- [32] Jackson SE, Chester JD. Personalised cancer medicine. *Int J Cancer J Int Cancer* 2015;137:262–6. doi:10.1002/ijc.28940.
- [33] Hayes DF. Biomarker validation and testing. *Mol Oncol* 2015;9:960–6. doi:10.1016/j.molonc.2014.10.004.
- [34] Schouten PC, Marme F, Aulmann S, Sinn H-P, Van Essen DF, Ylstra B, et al. Breast cancers with a BRCA1-like DNA copy number profile recur less often than expected after high-dose alkylating chemotherapy. *Clin Cancer Res Off J Am Assoc Cancer Res* 2014. doi:10.1158/1078-0432.CCR-14-1894.
- [35] Vollebbergh MA, Lips EH, Nederlof PM, Wessels LFA, Schmidt MK, van Beers EH, et al. An aCGH classifier derived from BRCA1-mutated breast cancer and benefit of high-dose platinum-based chemotherapy in HER2-negative breast cancer patients. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2011;22:1561–70. doi:10.1093/annonc/mdq624.
- [36] Rigtter LS, Loo CE, Linn SC, Sonke GS, van Werkhoven E, Lips EH, et al. Neoadjuvant chemotherapy adaptation and serial MRI response monitoring in ER-positive HER2-negative breast cancer. *Br J Cancer* 2013;109:2965–72. doi:10.1038/bjc.2013.661.
- [37] von Minckwitz G, Blohmer JU, Costa SD, Denkert C, Eidtmann H, Eiermann W, et al. Response-Guided Neoadjuvant Chemotherapy for Breast Cancer. *J Clin Oncol* 2013;31:3623–30. doi:10.1200/JCO.2012.45.0940.
- [38] Fuster D, Duch J, Paredes P, Velasco M, Munoz M, Santamaria G, et al. Preoperative Staging of Large Primary Breast Cancer With [18F]Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography Compared With Conventional Imaging Procedures. *J Clin Oncol* 2008;26:4746–51. doi:10.1200/JCO.2008.17.1496.
- [39] Riegger C, Herrmann J, Nagarajah J, Hecktor J, Kuemmel S, Otterbach F, et al. Whole-body FDG PET/CT is more accurate than conventional imaging for staging primary breast cancer patients. *Eur J Nucl Med Mol Imaging* 2012;39:852–63. doi:10.1007/s00259-012-2077-0.
- [40] Koolen BB, Vrancken Peeters M-JTFD, Aukema TS, Vogel WV, Oldenburg HSA, van der Hage JA, et al. 18F-FDG PET/CT as a staging procedure in primary stage II and III breast cancer: comparison with conventional imaging techniques. *Breast Cancer Res Treat* 2012;131:117–26. doi:10.1007/s10549-011-1767-9.
- [41] Groheux D, Giacchetti S, Delord M, Hindié E, Vercellino L, Cuvier C, et al. 18F-FDG PET/CT in staging patients with locally advanced or inflammatory breast cancer: comparison to conventional staging. *J Nucl Med Off Publ Soc Nucl Med* 2013;54:5-11. doi:10.2967/jnumed.112.106864.

- [42] Groheux D, Giacchetti S, Espié M, Vercellino L, Hamy A-S, Delord M, et al. The yield of 18F-FDG PET/CT in patients with clinical stage IIA, IIB, or IIIA breast cancer: a prospective study. *J Nucl Med Off Publ Soc Nucl Med* 2011;52:1526-34. doi:10.2967/jnumed.111.093864.
- [43] Hill S, Freemantle N. A role for two-stage pharmacoeconomic appraisal? Is there a role for interim approval of a drug for reimbursement based on modelling studies with subsequent full approval using phase III data? *PharmacoEconomics* 2003;21:761–7.
- [44] Sculpher M, Drummond M, Buxton M. The iterative use of economic evaluation as part of the process of health technology assessment. *J Health Serv Res Policy* 1997;2:26–30.
- [45] Claxton KP, Sculpher MJ. Using value of information analysis to prioritise health research: some lessons from recent UK experience. *PharmacoEconomics* 2006;24:1055–68.

PART II

PREDICTIVE BIOMARKERS: PERSONALIZE SYSTEMIC TREATMENT

CHAPTER 2

(Very) early health technology assessment and translation of predictive biomarkers in breast cancer

Anna Miquel-Cases*

Philip C Schouten*

Lotte MG Steuten

Valesca P Retèl

Sabine C Linn

Wim H van Harten

* First shared authorship

Submitted for publication

Abstract

Predictive biomarkers can guide treatment decisions in breast cancer. Many studies are undertaken to discover and translate these biomarkers, yet few are actually used for clinical decision-making. For implementation, predictive biomarkers need to demonstrate analytical validity, clinical validity and clinical utility. While attaining analytical and clinical validity is relatively straightforward by following methodological recommendations, achievement of clinical utility is more challenging. It requires demonstrating three associations: the biomarker with the outcome (prognostic association), the effect of treatment independent of the biomarker, and the differential treatment effect between the prognostic and the predictive biomarker (predictive association). Next to medical and biological issues, economical, ethical, regulatory, organizational and patient/doctor-related aspects are also influencing clinical translation. Traditionally, these aspects do not receive much attention until the formal approval or reimbursement of a biomarker test is at stake (via health technology assessment; HTA type of studies), at which point the clinical utility and sometimes price of the test can hardly be influenced anymore. However, if HTA analyses were performed earlier, during biomarker research and development, it could prevent the further development of those biomarkers unlikely to ever provide sufficient added value to society and rather facilitate translation of the promising ones. The use of early HTA is increasing and particularly relevant for the predictive biomarker field, as expensive medicines are increasingly under pressure and the urge for biomarkers to guide their appropriate use is huge. Closer interaction between clinical researchers and HTA experts throughout the translational research process will ensure that available data and methodologies are being used most efficiently to facilitate biomarker translation.

Introduction

Biomarkers are measurements of biological processes or disease that represent their state or activity. Since biomarkers signify a level of biological understanding, they can be exploited to improve research and clinical decision-making. For cancer treatment outcome, two types of biomarkers exist. Prognostic biomarkers associate with outcome and can help identify whether a patient should be treated. Predictive biomarkers, associate with outcome after a specific treatment and can guide the choice of treatment for an individual patient [1].

The neo-adjuvant (NACT) setting provides an *in vivo* research setting to identify predictive biomarkers, as in this setting the expression of biomarkers can be characterized prior to systemic treatment and the response to the therapy can subsequently be measured in the surgical specimen. Significant amounts of effort and money have been put in identifying predictive biomarkers to systemic NACT [2]. However, despite many studies being undertaken, few of these biomarkers are actually used for clinical decision making [3]. Several reasons may prevent more effective translation. Statistically studies are often poorly designed, clinically they lack a relevant use, and biologically they underestimate the complexity of drug mechanism of action and signaling pathways that confer sensitivity and resistance. Furthermore, economical, ethical, regulatory, organizational and patient/doctor-related aspects can affect translation as well.

Health Technology Assessment (HTA) is a multidisciplinary process that scientifically evaluates the medical, health economic, social and ethical aspects related to the adoption, implementation and use of a new technology or intervention. It aims to inform decisions on safe and effective health policies by seeking best value for money [4]. Traditionally, HTA does not receive much attention until the formal approval or reimbursement of a biomarker test is at stake. Early HTA refers to assessing these aspects alongside the basic, translational and clinical research process [5,6]. Early HTA can thus improve biomarker translation by preventing the further development of those biomarkers unlikely to ever provide sufficient added value to society, while facilitating translation of the promising ones [7]. Furthermore, it can be used to prevent late unfavorable assessments at the time the technology is being evaluated for cost-effectiveness and after big investments are done [8]. Common early HTA methods include literature reviews, evidence synthesis, decision analysis and health economic modeling as well as formal qualitative methods to elicit expert opinions and perform multi-criteria assessments for example in focus group discussions [5,9].

In this manuscript we discuss the clinical challenges in the translation of predictive biomarkers for NACT in breast cancer and provide concrete guidance on how the use of early HTA methods can support this process.

Types of treatment biomarkers

For treatment outcomes two types of biomarkers exist. Prognostic biomarkers inform on who to treat and predictive biomarkers inform on how to treat. The investigations of predictive biomarkers have to take into account three associations: the biomarker with the outcome (prognostic association), the effect of treatment independent of the biomarker, and the differential treatment effect between the prognostic and the predictive biomarker group (predictive association) [10–17]. Understanding these relations is important to choose the proper clinical action: to treat or not to treat in situations of good or very poor prognosis (prognostic biomarker), or to apply a treatment that is effective only in a subgroup of patients (predictive biomarker). For a hypothetical biomarker, survival curves that demonstrate prognostic value, treatment effects and predictive value are shown in figure 1. The overall landscape of the use of biomarkers for a particular population of patients can be illustrated by the therapeutic response surface [18] as shown in figure 2. This figure describes the relationship between treatment (drug and/or doses), sorted by prognostic characteristics, and clinical benefit of adding the treatment of a biologically homogeneous group of cancers. Through that figure one can identify patients for whom treatment should be spared, due to their exceptional prognosis or due to their increased risk of suffering from toxicities, and patients for whom additional treatment is likely to be beneficial, due to their poor prognosis in combination with on target treatment.

Figure 1: Prognostic, treatment and predictive effect. In this figure, hypothetical Kaplan-Meier curves resulting from biomarker negative and positive cases are shown. Patients have been treated with a specific treatment (A) or nothing. Two treatment effects can be observed (1 and 2), the prognostic effect is the difference between the non-treated biomarker-positive and negative patients. A differential treatment effect gives the predictive value.

Figure 2: Therapeutic response surface plotting clinical prognostic characteristics on the x-axis, treatment regimen and dose on the y-axis and clinical benefit on the z-axis. Several important regions are signaled: prognostic marker area, predictive biomarker area, the overlap between prognostically poor and predictive biomarker area in which a predictive biomarker adds benefit, the areas in which treatments are not working, and the area in which treatments may work but do not give benefit due to for example high toxicity. The easiest area being that of ineffective treatment i.e., the treatment does not add any benefit, despite the fact that some patients may seem to do well due to the good prognosis of their tumor. Some early stage tumors may have such good outcome that treatment is not advised, prognostic markers or characteristics should be used to identify these and spare patients the treatment. If one would use a predictive biomarker in this group, it could select patients and the therapy could seem efficacious given the good outcome. The extra benefit however would be smaller or non-existent due to the good prognosis from the outset. Predictive biomarkers can be identified as those markers that find groups of patients that benefit especially from a specific treatment (or dose). Suppose that the figure describes a homogenous group that can be identified by one biomarker. There would be one treatment option that adds benefit to all patients except those with good prognosis. This is illustrated by the ridge halfway the treatment axis in the figure. Additionally, some treatments may only add benefit to patients with intermediate prognostic characteristics and not those with poor characteristics. This may describe treatment burden-toxicity considerations. For example, in the case of two patients; one being young and without comorbidities, and one being older with many comorbidities, a treatment associated with high toxicity may only benefit the first, as shown in the figure by benefit decreasing in the area representing characteristics associated with poor prognosis.

Relevant HTA aspects			Example			HTA methods		
-Biomarker' effectiveness -LOE of available evidence -(expected) costs of testing -(expected) costs of research	-1st stage CEA (calculate the potential) -Tests' analytical validity -Costs of testing -Implementation and regulations demands -Patients' comfort -Ethical concerns	-1st stage CEA (calculate the potential) - Invested money - preliminary evidence -biomarker existence -logical - research plan - study design - Expected health gain	-Prospective vs retrospective -Study design -Regimen or single drug -Costs -Endpoint	-2nd stage CEA (calculate potential) -Final CEA -Organizational demands -Optimal implementation	-Does the test require personal training? / New working pathways in hospitals? / New material/ machinery? -What's the most efficient / cost-effective way to implement the test?			
-Biomarker A? PPV= 90%, testing= €3000, LOE medium, research= 2M -Biomarker B? 80%, €300, LOE high, 500K -Biomarker C? 70%, €200, LOE high, 300K	-Test 1 (biomarker A)? PPV= 90%, testing= €3000, new 30K machine, 1 week TOT, no patient discomfort (blood) - Test 2 (biomarker A)? 80%, €300, old infrastructure, 2 weeks TOT	-At which performance is the test CE? - See table 3	-Study 1? Retrospective, RCT, drug A vs drug B, 50K -Study 2? Prospective, RCT, drug A vs B, 2M -Study 3? Retrospective, case-control, drug A vs drug B, 5K	- Is the CE estimate uncertain? If so: - Which model parameters cause this uncertainty? - Is it worthwhile investing to gather more data? -Or is it best to wait for others' ongoing research to finish?	- Is the CE estimate uncertain? If so: - Which model parameters cause this uncertainty? - Is it worthwhile investing to gather more data? -Or is it best to wait for others' ongoing research to finish?	-Combination of the prior methods		
Quantitative: -CA -MCDA -AHP Qualitative: -Interviews, -discussions, -surveys -focus groups (Delphi method)	Quantitative: -CA, -MCDA -AHP Qualitative: -Interviews, -discussions, -surveys -focus groups (Delphi method)	- CEA model -SA Quantitative: -CA, -MCDA -AHP Qualitative: -Interviews, -discussions, -surveys -focus groups (Delphi method)	- CEA model -ROI Quantitative: -CA, -MCDA -AHP Qualitative: -Interviews, -discussions, -surveys -focus groups (Delphi method)	-CEA model -VOI -ROA Quantitative: -CA, -MCDA -AHP Qualitative: -Interviews, -discussions, -surveys -focus groups (Delphi method)	-CEA model -VOI -ROA Quantitative: -CA, -MCDA -AHP Qualitative: -Interviews, -discussions, -surveys -focus groups (Delphi method)	2		

Figure 3: Moment and type of decisions that (very) early and mainstream HTA can inform along the predictive biomarker research continuum.

*POP= proof of principle study, refers to the first in-human study. From an HTA perspective it is important to discern this because it provides the first Abbreviations: CE= cost-effectiveness analysis (CEA); CA= Conjoint analysis; MCDA=Multi criteria decision analysis; AHP= hierarchical analytical process; VOI= value of information analysis; ROA= real options analysis; RCT= randomized clinical trial; TOT= turnaround time; ROI= return on investment; LOE= level of evidence; PPV= positive predictive value; SA= sensitivity analysis; Bk-Tx-Ox= Biomarker-treatment-outcome; HTA= health technology assessment

Translating predictive biomarkers

To translate a biomarker from bench to bedside evidence is required that the test is reliable, that it separates a population in clinically relevant subgroups, and that applying the test results in improvement of clinical outcomes compared to not applying the test, respectively [19–23]. To address these criteria, predictive biomarker investigations typically involve multiple, often overlapping stages [24–31] (see Figure 3). After discovery, investigations range from laboratory experiments, to data mining exercises or clinical studies that aim to understand biological and/or clinical outcomes. Subsequently, the test may be improved. This can be done sequentially or in parallel with demonstrating its use in clinical studies [1,12,32]. The amount of evidence needed to demonstrate clinical utility will be weighed on a per-biomarker basis. The process may consist of differing combinations of studies [1]. Multiple rounds of testing may be performed until sufficient quality of the test and validation has been reached for regulatory approval. This differs between countries. For instance in the US, approval is granted by the FDA while in Europe this is the responsibility of national certified bodies. Commercialized biomarker tests are high risk medical devices [33,34]. In Europe this means demonstration of safety and performance suffices to get the CE- mark [35]. In the US demonstration of safety and effectiveness is required (premarket approval [34]). Yet if biomarkers tests are developed as in-house tests, performed in specific health care institutions, the situation differs. While in the US lab certification according to the Clinical Laboratory Improvement Amendments (CLIA)[36] is needed, in the EU there is no applicable regulation yet, although the medical device directive is currently being revised [37]. Reimbursement is the procedure that will facilitate wide spread use of the biomarker test; it is country specific and nowadays generally based on a cost-based criteria. However, value-based criteria are expected to become the norm as is the case for pharmaceuticals.

Studies on predictive biomarkers do not reach a high level of evidence (Case study: predictive biomarkers for NACT in breast cancer)

We performed a systematic search to identify tumor biomarkers that predict NACT response in breast cancer (n= 134, specific methods are described in the annex). Based on the type and quality of the identified studies, we concluded that biomarkers of NACT for breast cancer are in early stage evaluation. The characteristics of the identified studies are summarized in Figure 4. We found that drugs involved were generally standard NACT (regimens), that few genes have been investigated more than once (either in different studies or with different tests) and that all studies had a control for biomarker negative patients. On the other hand, only 8% (11/134) of the studies used control groups without the treatment of interest, and even those that had options for controlling did not. Based on the reported analysis interpretation, many studies found that the marker under investigation could be predictive. In those without control groups the amount of

'positive' studies was about 69% (85/123) versus 60% (6/10) in those with control groups. These conclusions can be misleading in the absence of control groups.

Challenges in translating predictive biomarkers

Our review showed that biomarkers of NACT for breast cancer are in early stage evaluation. The underlying success in the translation of a predictive biomarker is the final demonstration of clinical utility. This requires an a priori right choice of biomarker, treatment and outcome to investigate a particular application, as well as a continuous pursuance to correctly establish the link between these three entities in validation studies.

With regards to the biomarker, in principle, any biomarker/mechanism or biological entity can be investigated. Similarly any single drug or drug regimen can be investigated in relation to the biomarker. It is likely that resistance and sensitivity mechanisms are drug specific, hence for the dissection of such mechanisms, ideally, only one treatment variable should be tested in the study design. The design could be drug A versus nothing, drug A versus AB, or combo AB versus ABC, etc. Instead, if drug A is compared to drug B, or combo ABC with combo CDE, it won't be possible to dissect single drug resistance or drug sensitivity mechanisms anymore. However, treatment in the NACT setting is in principle curative, therefore, it is ethically impossible to withhold proven or apply only unproven treatment, thus many studies have mixed effects. That is why trying to identify biomarkers in these studies could be heavily confounded. Knowing this, it is important to include control groups for the biomarker (negative and positive) and for the treatment (treatment of interest and a comparator) and derive the treatment effect, prognostic effect and predictive effect of the biomarker [10–17]. If the theoretically best control is not available, resorting to a control group with the current clinical best practice is essential as it sets the minimal expected performance.

Regarding the clinical outcome, it remains important to carefully choose the endpoint that fits with the intended application and aim. The NACT setting provides rapid assessment of biomarker effectiveness by means of pathologic complete response (pCR), a surrogate endpoint of long-term survival [38,39]. Although pCR has gained acceptance in research and in the clinics, its association with long-term survival is not straightforward [40]. While pCR is a measure of local treatment effect, which measures tumor shrinkage, long-term survival is a measure of systemic treatment effect, which measures the presence or absence of events as consequence of the presence or absence of micro-metastasis. The outcome measure should give insight into the sensitivity of the cancer cell population (e.g. (a clone of the) primary lesion, metastatic lesion, a stem-cell population, etc.) that determines the overall prognosis.

Differences between the measured population and this population will lead to unexpected results, i.e., bad outcome where expected a good one, or vice versa. The interpretations that may derive from the use of pCR to predict survival are summarized in table 1. In some cases the early response measured by pCR translates well into improved patient survival, this is the case of patients in the case mix in the grey row. However in most of the cases it does not, as shown in the white rows. The majority of breast cancer subtypes in the case mix where pCR does not translate into improved breast cancer specific survival i.e., luminal B/HER2-positive or luminal A tumors probably fall in these last categories. Hard endpoints like relapse free survival (RFS), distant metastasis free interval (DMFI) or overall survival (OS) are measures of systemic treatment effect. Their downside is the confounding due to additional adjuvant and/or metastatic treatment and due to competing risks, next to the lengthy time required for its measurement.

The combination of a specific biomarker, treatment and outcome sets the stage for the envisioned application and investigations need. This combination needs to show analytical validity, clinical validity and clinical utility. While many problems that can arise during the analytical validity and clinical validity phases i.e., using correct study designs or analytical robustness, can be tackled by strictly following known methodological recommendations or guidelines [1,10,21,41], demonstrating clinical utility is rather difficult. This is the consequence of the majority of clinical datasets not providing high levels of evidence (LOE), for example due to missing control groups. Furthermore, for some applications, no suitable clinical dataset may be available. For example, biomarker-drug combinations that were identified in modeling systems may not have a clinical dataset in the neoadjuvant setting. Additionally, many neoadjuvant biomarker studies do not use a control treatment since it is thought that pCR is a direct proof of *specific* treatment efficacy. When data-mining is performed in such cohorts it is easy to identify confounded associations as interesting. These are examples that show that identifying and establishing the predictive value of a biomarker may be jeopardized by design limitations [17].

Concluding, for any biomarker-treatment-outcome analysis intended for implementation, the application is a specific case for which high LOE needs to be gathered, as from this application a particular clinical decision will follow i.e., withholding or giving a specific treatment. Any non-high-level, circumstantial evidence or evidence that fits another application should thus be considered too early. Randomized trials provide the most optimal setting in which this interaction can be investigated properly.

Table 1: Interpretations that derive from the use of pCR to predict survival

Measured outcomes		Interpretation	Underlying research question
Surrogate outcome	Long term outcome	Can treatment downsize tumour for breast conserving surgery?	Can treatment eliminate micrometastases that would otherwise grow into macrometastases using pCR as a read-out?
pCR – at diagnosis no micrometastases present that could turn into macrometastases	Favourable	Yes	Incorrect interpretation that treatment can eliminate micrometastases that could turn into macrometastases
No pCR – at diagnosis no micrometastases present that could turn into macrometastases	Favourable	Depends on amount of downsizing achieved, tumour size at diagnosis, and breast size	Confounder in ‘poor’ prognosis distant-recurrence free interval curve – since no pCR was achieved
pCR – at diagnosis micrometastases present that could turn into macrometastases	Favourable	Yes	Correct interpretation that treatment can eliminate micrometastases that could turn into macrometastases
pCR – at diagnosis micrometastases present that could turn into macrometastases	Distant recurrence	Yes	Incorrect interpretation that treatment can eliminate micrometastases that could turn into macrometastases; primary tumor is eliminated, but not micrometastatic tumor cells
No pCR – at diagnosis micrometastases present that could turn into macrometastases	Favourable	Depends on amount of downsizing achieved, tumour size at diagnosis, and breast size	Incorrect interpretation that treatment cannot eliminate micrometastases that could turn into macrometastases; primary tumor is not completely eliminated, but micrometastatic tumor cells are
No pCR – at diagnosis micrometastases present that could turn into macrometastases	Distant recurrence	Depends on amount of downsizing achieved, tumour size at diagnosis, and breast size	Correct interpretation that treatment cannot eliminate micrometastases that could turn into macrometastases, since primary tumor cells cannot be eliminated completely either

The role of early Health Technology Assessment

While medicine and biology are the basis for predictive biomarker research, economical, ethical, regulatory, organizational and patient/doctor-related aspects influence biomarkers' translation and adoption as well. These aspects are often assessed nearing decisions on coverage or reimbursement. However, if HTA analyses were performed earlier ((very) early HTA), during biomarker research and development, it could prevent the further development of those biomarkers unlikely to ever provide sufficient added value to society and rather facilitate translation of the promising ones. Furthermore, it could help appraising other relevant aspects timely, as the trade-offs with alternate approaches or the performance requirements for a specific technology to reach cost-effectiveness. [7].

In figure 3, we present the moment and the type of decisions that (very) early and mainstream HTA can inform along the predictive biomarker research continuum. The difference between very early and early HTA mainly lies on the availability of evidence from the assessed technology (very limited at the time of using very early HTA), and the methodology used (more use of modeling methods and assumptions in very early HTA). Furthermore, in figure 3 we provide a sample of common HTA methods used to inform these decisions. This does not provide all existing HTA methods (most of them can be found in references [5,9,42]), but highlights those that seem specifically useful for predictive biomarker research. Descriptions of the technical methods are provided in supplementary table 2.

(Very) early HTA is not yet used to assess predictive biomarkers

(Case study: predictive biomarkers for NACT in breast cancer)

We performed a systematic search to identify the current use of early HTA methods during the research and translation process of predictive biomarkers for NACT treatment in breast cancer (n= 31, specific methods are described in the supplementary material). These studies were classified on being on very early, early or mainstream HTA according to Figure 3, and on whether they described clinical, economic, ethical, organizational and patient/doctor related aspects. The identified studies were classified either as early or mainstream HTA, but none as very early HTA. Almost all early HTA articles reported on the comparative effectiveness of testing techniques [43–47]. Only one article presented an early stage cost-effectiveness analysis [48]. Another article presented an organizational and/or implementation aspect; the increase in uptake of a biomarker test as a consequence of new potential clinical applications [49]. Opinion leaders attitudes were used to gather potential issues arising from 'treatment-focused' genetic testing in one article [50]. The findings of this exploratory review on early HTA were similar to those of a 2014 review on early HTA for medical devices [9], where no studies for predictive biomarkers for breast cancer were found.

Improving the translation of predictive biomarkers from an HTA perspective

Our systematic review found that (very) early HTA is not applied along the research process of predictive biomarkers for NACT treatment in breast cancer. Different HTA aspects are relevant to address different type of decisions during the research process and can facilitate translation (figure 3 contains all references to methods).

Biomarker identification (a, figure 3)

At this stage, the presence of limited budgets and/or time can force researchers into decisions on which biomarker to involve in further investigations i.e., biomarker A (90% positive predictive value (PPV), medium LOE, €3000 expected testing costs and 2M expected validation costs), biomarker B (80%, high LOE, €300 and 500K) or biomarker C (70%, high LOE, €2000 and 300K)? As illustrated, aspects likely to play a role on this decision are the biomarker's PPV, the LOE of this evidence, the expected costs of testing and the expected costs for its validation. The conjoint analysis (CA), the multi criteria decision analysis (MCDA) and the analytical hierarchical process (AHP) are methods that can be used to prioritize these biomarkers, in a step-wise approach by using the aforementioned relevant aspects to compare and judge them. These judgments are made by a selected group of doctors, patients, developers, payers and policymakers. They are all decision-makers along the development process and can provide useful knowledge to the decision. In some situations, the evidence to characterize the aspects of the biomarkers will not yet be there i.e., the PPV of the test is not clear. In such cases, prior to starting the CA, MDCA or AHP process, estimates for these aspects can be derived by means of expert elicitation methods (via CA, MCDA, AHP or other elicitation methods) or by extrapolation from similar biomarker-drug cases (see methods of supplementary table 2 with references to case studies). In other situations, a quantitative-driven decision may not seem applicable yet. In this case, biomarker selection can be made via (semi) qualitative methods such as interviews, discussions, survey or focus groups (Delphi method). These methods allow a more flexible decision-making process and they are already common practice.

Biomarker translation (b, figure 3)

After biomarker selection has been made and the first proof of principle (POP) study has been conducted (refers to the first in-human study), the researcher questions whether more research towards biomarker validation should be continued. Assuming the endpoint of research is maximizing health outcomes with the resources available to society, this question can be answered by using the value of information analysis (VOI) method. VOI execution requires a prior construction of a CE model (with the POP data) and a first stage CEA. VOI analysis will translate

the magnitude of uncertainty around this first cost-effectiveness estimate into a monetary value that could lead to full certainty on the biomarkers' CE. This value (the expected value of perfect information (EVPI)) is subsequently compared to the expected costs of conducting further research, and if these are lower, it suggests that conducting further research is worthwhile. Further calculations of the VOI analysis can help determining for which data type is most beneficial to conduct research i.e., PPV of the test or quality of life of the administered treatment (the expected value of partial perfect information (EVPI)), and with which type and magnitude of study designs should this be conducted (Expected value of sampling information (EVSII)). A next relevant question is the timing to start these studies. The real option analysis (ROA) method helps deciding when it is most worthwhile to undertake this research. Whether it is best to invest on further research immediately or whether it is best to wait for current ongoing studies to be finished before investing. Maybe these studies already provide some evidence that can increase the CE uncertainty without needing investment. This option takes into account the costs of withholding the use of the biomarker and thus the possibility of giving suboptimal treatment to patients in the meantime. ROA is especially useful at these stages of development, when large investments are still expected.

Upon the decision of starting further biomarker validations, a biomarker test needs to be chosen. Available tests to measure one biomarker may have very different characteristics i.e. test 1 (PPV 90%, €3000 expected testing costs, new 30K machine, 1 week turnaround time (TOT), patient comfort (blood)) or test 2 (80%, €300, old infrastructure, 2 weeks TOT)? As illustrated, aspects likely to play a role on this decision are the tests' analytical validity, the expected costs of testing, its implementation and regulatory demands, the patients' comfort, and ethical concerns. This choice can be made by using the same methods described in the biomarker identification stage. Yet in the case evidence to define the biomarkers' aspects is lacking, other methods than the previously described are useful. For instance, usability testing to determine patients' comfort during the usage of a specific test, or the multipath mapping tool to forecast the implementation demands of the test (see supplementary table 1).

Biomarker tests performance has traditionally been guided by effectiveness. By accounting for the costs associated to false cases, a more realistic minimum performance that can warrant the tests' clinical application can be determined. This can be achieved by using the already built CE model together with the one-way sensitivity analysis (SA) method. This means varying model parameter values that represent performance in the model to determine the minimum performance values where cost-effectiveness remains and to see which parameters drive the cost-effectiveness. The SA method can be used any time during biomarker development to explore how new test features affect CE. It is essential that this goes along with updates on clinical and economic evidence in the CE model.

Another consideration that may be relevant at this point is to anticipate the expected yield of future investigations and its associated investments. Its evaluation can be done by using the concept of returns on investment (ROI). By drawing a likely research plan for the specific biomarker and considering the amount of money invested and the expected health outcomes gained in return. Hypothetical scenarios on possible 'research plans' for predictive biomarker development and its economic and health consequences are explained in table 2. The scenarios show that opting for the speedy solutions with wrong study designs when there is low level of preliminary evidence can lead to futile expenditures. On the other hand, investing in basic research endeavors or prospective validation studies, that seem more costly at the onset, is likely to lead to improved health outcomes.

While ROI type of analysis can provide an overview of the consequences of a specific research plan, the use of CA, MCDA or AHP methods can help optimally designing each validation study. The basis is to consider the high costs of setting up new studies with the optimal features these can offer versus the of use already available data which is less costly but comes with limitations (retrospective, presence/absence control group, availability of hard endpoints or drug administered alone) i.e., choice between study 1 (retrospective, RCT, drug A vs drug B, 50K), study 2 (prospective, RCT, drug A vs B, 2M) or study 3 (retrospective, case-control, drug A vs drug B, 5K)? This choice will be driven by the timing of the study (prospective vs retrospective), the understanding of the underlying biological mechanism, the study design, the presence of a drug regimen or single drug, the costs of the study and the endpoint. In this case, the execution of CA, MCDA or AHP methods should include other specialized experts, such as statisticians, molecular biologists and/or epidemiologists. The final choice can be further investigated by using clinical trial simulations (CTS) that can explore the effects of specific design assumptions to the expected outcomes.

Biomarker validation (c, figure 3)

Prior to each validation study, one will reflect upon the need for a further study, the nature of the study and the timing of such study. By updating the CE model with the newly generated evidence and using the CEA, VOI and ROA methods, as explained in the biomarker translation phase, these questions can be answered taking the broader health economic perspective. Furthermore, decisions on study design characteristics can be assessed at any time as explained in the biomarker translation phase.

Table 2: Hypothetical scenarios on possible ‘research plans’ for predictive biomarker development and its economic and health consequences. These scenarios are composed of four characteristics: 1) whether a consistent path of investigations for the aim is followed, 2) whether the studies are designed properly; 3) whether the preliminary evidence is strong and reliable; 4) whether the biomarker under investigation actually exists. Based on those, we hypothesized discovery paths a biomarker may follow and whether approval and reimbursement of the biomarker test can be obtained.

Reliable preliminary evidence	Biomarker exists or test is reliable	Logical steps for the plan / all evidence is contributing	Proper study designs		Basic research/ retrospective trials	POP / First in Human	Prospective Trials	Evidence sufficient for approval and use	Sufficiently cost-effective for reimbursement	Total investment compared to best case scenario							Economic outcome	Health outcomes
			yes	yes						yes	yes	reference	well invested	improved				
yes	yes	yes	yes	yes	yes		yes	yes	yes	equal to reference	high loss (invested money)	high loss (not used)						
no	yes	yes	yes	maybe	yes	no	no	no	n/a	lower	low loss (based on wrong evidence)	high loss (not improved)						
no	no	yes	no		yes	yes	no	no	n/a	higher	high loss (invested money)	low loss						
no	yes	no	no		no	no	no	n/a		lower	low loss (based on wrong evidence)	depends						
no	yes	no	no	yes		yes	yes	yes	yes	higher	low loss (unnecessary studies)	improved (but wasted time and money)						
yes	no	no	no	yes	yes	yes	no	n/a	n/a	equal to reference	high loss (invested money)	low loss						

health: high loss = biomarker not used or biomarker does not improve outcomes
costs: high loss = many studies performed vs. early stop of studies

Finally, once biomarker clinical utility is almost demonstrated, questions on future adoption and implementation demands become relevant. For instance, does the test require personnel training, the generation of new working pathways or the purchase of machinery? It is likely that during prior stages of the biomarker development process these questions have already been addressed (via previously mentioned methods like interviews, discussions or MCDA type of methods). Additional issues to address at this stage are the availability of resources for immediate implementation of the biomarker. A quantitative method specially formulated to anticipate and quantify demands is resource-modeling analysis. Also important is to determine the optimal implementation scenario for the test. This can be determined by using the SA method together with the final updated version CE model. For instance, it can determine the optimal turn-around time for the test by varying the parameter values that represent material and personnel requirements. Last, the final cost-effectiveness of the test can be determined. Recently, Coverage with Evidence Development (CED) programs were initiated throughout Europe and the US. These programs contain a (randomized) controlled trial including a broad Health Technology Assessment, where the new technology/drug is already being reimbursed. This program seems to be highly applicable for this setting. A first example has recently started in the Netherlands ('BRCA1-like biomarker for stage III breast cancer').

Important to highlight is that integration of HTA into the biomarker development process requires communication between researchers, clinicians, health-economists and decision-makers. This cooperation is necessary to ensure that all the relevant questions to move forward the biomarker translation process are answered and that appropriate data and methods are used. Partnerships like the Center for Translational Molecular Medicine (CTMM) in the Netherlands [51] or the INterdisciplinary HEalth Research International Team on BReast CAncer susceptibility (INHERIT BRCA) in Canada [52] have demonstrated that collaborations result in solid scientific impact and accelerated translational research.

Box 1 provides a summary of the review in 7 key points.

Box 1

- Our investigations concluded that predictive biomarkers for neo-adjuvant treatment of breast cancer are in early stage evaluation and that (very) early HTA is hardly being used.
- There is no best investigational nor HTA framework for predictive biomarkers, and it is likely best to keep analyses case-specific
- Predictive biomarker research requires specific study design choices to characterize the treatment effect, prognostic effect and predictive effect in a biomarker-treatment-outcome combination
- Predictive biomarker research could be planned based on current evidence but taking into account future required investigations and associated investments that go with it.
- Use the HTA and study design methodology appropriate for the current investigational stage critically, to make explicit why or how a certain study contributes to reaching a specific target
- Consider early on research and during development the regulatory, organizational, patient-related and economic requirements of biomarker development and ask help for those considerations that you do not understand
- Different HTA methods can inform different decisions during biomarker research. While multiple choice decisions can be informed by using CA, AHP and MCDA methods, decisions on the continuity and design of further research can be informed by using the CE model together with CEA VOI, ROA methods.

Outlook

It is likely that the use of predictive biomarkers will become more prevalent. We will describe the advances in this field by using the previously mentioned components of a successful predictive biomarker: the biomarker, the treatments, the outcome and the relation between these three parameters. Regarding the biomarker, our understanding of tumor biology has greatly expanded due to the use of high throughput methods, allowing for simultaneous assessment of tumors at DNA, RNA and protein level [53]. In combination with experimental data, discovering mechanisms of action should improve the chances of finding predictive biomarkers. However, it has also become clear that tumors are more heterogeneous than often described before [54]. Evolutionary pressure exists both intrinsically as well as extrinsically, by applying selection through therapies. Under these pressures, multiple resistance mechanisms may be present or develop [55]. This heterogeneity should be taken into account for predictive biomarkers. For example, it could be that differential sensitivity between the primary tumor and occult systemic disease exists, especially when NACT is used in presence of occult systemic disease. Measuring biomarkers in the tumor is an invasive procedure and the development of bloodstream biomarkers is promising. Yet it has to be proven, first, whether the ease of assaying outweighs the uncertainty on which lesion is being investigated, and second, whether the bloodstream ("liquid biopsies") can be used sufficiently reliable to forego tumor sampling [56,57]. Focusing outside of the tumor, host factors

can affect the sensitivity of these, as they contribute significantly to varying drug responses. For instance, drug metabolism (pharmacodynamics) has been recognized to result in different levels of drugs exposure. The dose of drug (regimens) administered is widely optimized to be as high as possible while having acceptable toxicity for a large population. This results in the under-treatment of some patients, whereas other patients develop unacceptable toxicity [58–61]. Another host factor currently being investigated is the immune/tumor microenvironment system, which also seems to contribute or shape drug response [62]. First, the immune system may be sensitized to attack tumor cells or already work to keep the tumor from expanding in a balance between tumor growth and immune cell killing. Contrary to this tumor-suppressing role, the immune system's tumor promoting role may be important. Both the immune system and micro-environment may act as protective factors against therapy. The compromised or tumor-recruited microenvironment could therefore be predictive for response [63].

Regarding the drugs a range of new drugs targeted at specific proteins are being developed aiming for a more specific killing of tumor cells [64,65]. With this increased target specificity, developing companion diagnostic may become more straightforward or even already available from outset. These targeted therapies are increasingly added to drug regimens used in the NACT setting [66]. Although currently used chemotherapy drugs were identified in screening efforts the identification of its mechanism of action to improve efficacy, reduce toxicity, and predict their resistance/sensitivity is an ongoing effort [67–72]. This knowledge and new biomarkers could make 'untargeted' drugs similar to newly mechanistically developed targeted drugs. Both old and new drugs may have unexpected efficacy in certain subsets of tumors that was previously overlooked due to the then current standard of developing drugs for the whole tumor populations rather than a more targeted approach. Linking the improved tumor characterizations to better characterized cohorts likely will improve understanding of reliable endpoints [73–77]. It will also facilitate the translation to clinical practice of biomarker-drug combinations that meaningfully improve treatment outcomes.

The use of early HTA is still not incorporated into routine practice, yet it is expected to become more common [78]. Especially in the predictive biomarker field, as expensive medicines like nivolumab are increasingly used for the total population and the urge for biomarkers is huge. Early HTA can help making the biomarker research process more efficient, so as to prevent futile investments and delays in patient access. With the raise of multiple testing, the use of panels and whole genome testing, the construction of CEA models will become more complex, the amount of effectiveness data originating from studies that are not RCTs (e.g., practice based studies) will increase and we will be facing so far unaddressed ethical and organizational concerns. This will require the development of innovative evaluation frameworks outside the traditional model-based CEA, where the remaining HTA aspects have more weight in decision-making. Furthermore,

these assessments will be required to be more iterative, rapidly incorporating new evidence and re-calculating outcomes.

Concluding, we found that research on biomarkers (in NACT) is methodologically weak and provided suggestions for improvement that are of a rather basic methodological nature. Early stage HTA can be more fully exploited in assisting in- and preparing for bringing the findings to the next translational development stage (or falsifying developments in a timely way). Closer interaction between clinical researchers and HTA experts may smoothen these processes. With the lessons from the past, the current possibilities of techniques, exciting times are ahead that may improve therapy choices for patients by optimizing existing applications and discovery of new options.

References

- [1] Hayes DF. Biomarker validation and testing. *Mol Oncol* 2015;9:960–6. doi:10.1016/j.molonc.2014.10.004.
- [2] Ptolemy AS, Rifai N. What is a biomarker? Research investments and lack of clinical integration necessitate a review of biomarker terminology and validation schema. *Scand J Clin Lab Investig Suppl* 2010;242:6–14. doi:10.3109/00365513.2010.493354.
- [3] Poste G. Bring on the biomarkers. *Nature* 2011;469:156–7. doi:10.1038/469156a.
- [4] European Network for Health technology Assessment (eunetha). What is Health Technology Assessment (HTA) n.d. <http://www.eunetha.eu/about-us/faq#t287n73>.
- [5] Ijzerman MJ, Steuten LMG. Early assessment of medical technologies to inform product development and market access: a review of methods and applications. *Appl Health Econ Health Policy* 2011;9:331–47. doi:10.2165/11593380-000000000-00000.
- [6] Claxton K, Griffin S, Koffijberg H, McKenna C. How to estimate the health benefits of additional research and changing clinical practice. *BMJ* 2015;351:h5987.
- [7] Şardaş S, Endrenyi L, Gürsoy UK, Hutz M, Lin B, Patrinos GP, et al. A call for pharmacogenovigilance and rapid falsification in the age of big data: why not first road test your biomarker? *Omics J Integr Biol* 2014;18:663–5. doi:10.1089/omi.2014.0132.
- [8] Steuten LM, Ramsey SD. Improving early cycle economic evaluation of diagnostic technologies. *Expert Rev Pharmacoecon Outcomes Res* 2014;14:491–8. doi:10.1586/14737167.2014.914435.
- [9] Markiewicz K, van Til JA, Ijzerman MJ. MEDICAL DEVICES EARLY ASSESSMENT METHODS: SYSTEMATIC LITERATURE REVIEW. *Int J Technol Assess Health Care* 2014;30:137–46. doi:10.1017/S0266462314000026.
- [10] Simon R. Clinical trial designs for evaluating the medical utility of prognostic and predictive biomarkers in oncology. *Pers Med* 2010;7:33–47. doi:10.2217/pme.09.49.
- [11] Simon R. Sensitivity, Specificity, PPV, and NPV for Predictive Biomarkers. *J Natl Cancer Inst* 2015;107. doi:10.1093/jnci/djv153.
- [12] Simon RM, Paik S, Hayes DF. Use of archived specimens in evaluation of prognostic and predictive biomarkers. *J Natl Cancer Inst* 2009;101:1446–52. doi:10.1093/jnci/djp335.
- [13] Mandrekas SJ, Sargent DJ. Clinical trial designs for predictive biomarker validation: theoretical considerations and practical challenges. *J Clin Oncol Off J Am Soc Clin Oncol* 2009;27:4027–34. doi:10.1200/JCO.2009.22.3701.
- [14] Sargent DJ, Conley BA, Allegra C, Collette L. Clinical trial designs for predictive marker validation in cancer treatment trials. *J Clin Oncol Off J Am Soc Clin Oncol* 2005;23:2020–7. doi:10.1200/JCO.2005.01.112.
- [15] Janes H, Pepe MS, McShane LM, Sargent DJ, Heagerty PJ. The Fundamental Difficulty With Evaluating the Accuracy of Biomarkers for Guiding Treatment. *J Natl Cancer Inst* 2015;107. doi:10.1093/jnci/djv157.
- [16] Fine JP, Pencina M. On the quantitative assessment of predictive biomarkers. *J Natl Cancer Inst* 2015;107. doi:10.1093/jnci/djv187.
- [17] Henry NL, Hayes DF. Uses and abuses of tumor markers in the diagnosis, monitoring, and treatment of primary and metastatic breast cancer. *The Oncologist* 2006;11:541–52. doi:10.1634/theoncologist.11-6-541.
- [18] Sheiner LB. Learning versus confirming in clinical drug development. *Clin Pharmacol Ther* 1997;61:275–91. doi:10.1016/S0009-9236(97)90160-0.
- [19] Parkinson DR, McCormack RT, Keating SM, Gutman SI, Hamilton SR, Mansfield EA, et al. Evidence of clinical utility: an unmet need in molecular diagnostics for patients with cancer. *Clin Cancer Res Off J Am Assoc Cancer Res* 2014;20:1428–44. doi:10.1158/1078-0432.CCR-13-2961.
- [20] Duffy MJ, Sturgeon CM, Sölétormos G, Barak V, Molina R, Hayes DF, et al. Validation of new cancer biomarkers: a position statement from the European group on tumor markers. *Clin Chem* 2015;61:809–20. doi:10.1373/clinchem.2015.239863.
- [21] Polley M-YC, Freidlin B, Korn EL, Conley BA, Abrams JS, McShane LM. Statistical and practical considerations for clinical evaluation of predictive biomarkers. *J Natl Cancer Inst* 2013;105:1677–83. doi:10.1093/jnci/djt282.

- [22] Simon R. Lost in translation: problems and pitfalls in translating laboratory observations to clinical utility. *Eur J Cancer Oxf Engl* 1990;44:2707–13. doi:10.1016/j.ejca.2008.09.009.
- [23] Henry NL, Hayes DF. Uses and abuses of tumor markers in the diagnosis, monitoring, and treatment of primary and metastatic breast cancer. *The Oncologist* 2006;11:541–52. doi:10.1634/theoncologist.11-6-541.
- [24] Parkinson DR, McCormack RT, Keating SM, Gutman SI, Hamilton SR, Mansfield EA, et al. Evidence of clinical utility: an unmet need in molecular diagnostics for patients with cancer. *Clin Cancer Res Off J Am Assoc Cancer Res* 2014;20:1428–44. doi:10.1158/1078-0432.CCR-13-2961.
- [25] Simon R. Clinical trials for predictive medicine: new challenges and paradigms. *Clin Trials Lond Engl* 2010;7:516–24. doi:10.1177/1740774510366454.
- [26] Kelloff GJ, Sigman CC. Cancer biomarkers: selecting the right drug for the right patient. *Nat Rev Drug Discov* 2012;11:201–14. doi:10.1038/nrd3651.
- [27] Schully SD, Carrick DM, Mechanic LE, Srivastava S, Anderson GL, Baron JA, et al. Leveraging Biospecimen Resources for Discovery or Validation of Markers for Early Cancer Detection. *JNCI J Natl Cancer Inst* 2015;107:djv012–djv012. doi:10.1093/jnci/djv012.
- [28] Hayes DF. Biomarker validation and testing. *Mol Oncol* 2015;9:960–6. doi:10.1016/j.molonc.2014.10.004.
- [29] Simon RM, Paik S, Hayes DF. Use of archived specimens in evaluation of prognostic and predictive biomarkers. *J Natl Cancer Inst* 2009;101:1446–52. doi:10.1093/jnci/djp335.
- [30] McShane LM, Cavenagh MM, Lively TG, Eberhard DA, Bigbee WL, Williams PM, et al. Criteria for the use of omics-based predictors in clinical trials. *Nature* 2013;502:317–20. doi:10.1038/nature12564.
- [31] Simon R, Roychowdhury S. Implementing personalized cancer genomics in clinical trials. *Nat Rev Drug Discov* 2013;12:358–69. doi:10.1038/nrd3979.
- [32] Beelen K, Zwart W, Linn SC. Can predictive biomarkers in breast cancer guide adjuvant endocrine therapy? *Nat Rev Clin Oncol* 2012;9:529–41. doi:10.1038/nrclinonc.2012.121.
- [33] European Commission. Regulation of the European Parliament and of the Council on medical devices, and amending directive 2001/83/EC, Regulation (EC) No 178/2002 and Regulation (EC) No 1223/2009. n.d.
- [34] U.S. Food and Drug Administration. Premarket Approval (PMA) n.d. <http://www.fda.gov/Medicaldevices/Deviceregulationandguidance/%20Howtomarketyourdevice/Premarketsubmissions/Premarketapprovalpma/>.
- [35] Annex IX of the EU Directive 93/42/EEC. n.d.
- [36] U.S. Food and Drug Administration. Clinical Laboratory Improvement Amendments (CLIA) n.d. <http://www.fda.gov/medicaldevices/deviceregulationandguidance/ivdregulatoryassistance/ucm124105.htm>.
- [37] European Commission. Revisions of Medical Device Directives n.d. http://ec.europa.eu/growth/sectors/medical-devices/regulatory-framework/revision/index_en.htm.
- [38] Fisher B, Bryant J, Wolmark N, Mamounas E, Brown A, Fisher ER, et al. Effect of preoperative chemotherapy on the outcome of women with operable breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol* 1998;16:2672–85.
- [39] Kuerer HM, Newman LA, Smith TL, Ames FC, Hunt KK, Dhingra K, et al. Clinical course of breast cancer patients with complete pathologic primary tumor and axillary lymph node response to doxorubicin-based neoadjuvant chemotherapy. *J Clin Oncol Off J Am Soc Clin Oncol* 1999;17:460–9.
- [40] Jones A, O'Brien M, Sommer H, Nowara E, Welt A, Pienkowski T, et al. Phase II study of oral vinorelbine in combination with capecitabine as second line chemotherapy in metastatic breast cancer patients previously treated with anthracyclines and taxanes. *Cancer Chemother Pharmacol* 2010;65:755–63. doi:10.1007/s00280-009-1081-y.
- [41] McShane LM, Hayes DF. Publication of tumor marker research results: the necessity for complete and transparent reporting. *J Clin Oncol Off J Am Soc Clin Oncol* 2012;30:4223–32. doi:10.1200/JCO.2012.42.6858.
- [42] Martin JL, Murphy E, Crowe JA, Norris BJ. Capturing user requirements in medical device development: the role of ergonomics. *Physiol Meas* 2006;27:R49–62. doi:10.1088/0967-3334/27/8/R01.
- [43] Cockburn A, Yan J, Rahardja D, Euhus D, Peng Y, Fang Y, et al. Modulatory effect of neoadjuvant chemotherapy on biomarkers expression; assessment by digital image analysis and relationship to residual cancer burden in patients with invasive breast cancer. *Hum Pathol* 2014;45:249–58. doi:10.1016/j.humpath.2013.09.002.

- [44] García-Caballero T, Prieto O, Vázquez-Boquete A, Gude F, Víaño P, Otero M, et al. Dual-colour CISH is a reliable alternative to FISH for assessment of topoisomerase 2-alpha amplification in breast carcinomas. *Breast Cancer Res Treat* 2014;143:81–9. doi:10.1007/s10549-013-2791-8.
- [45] Castle J, Shaker H, Morris K, Tugwood JD, Kirwan CC. The significance of circulating tumour cells in breast cancer: a review. *Breast Edinb Scotl* 2014;23:552–60. doi:10.1016/j.breast.2014.07.002.
- [46] Kuo S-J, Wang BB-T, Chang C-S, Chen T-H, Yeh K-T, Lee D-J, et al. Comparison of immunohistochemical and fluorescence in situ hybridization assessment for HER-2/neu status in Taiwanese breast cancer patients. *Taiwan J Obstet Gynecol* 2007;46:146–51. doi:10.1016/S1028-4559(07)60008-4.
- [47] Logullo AF, de Moura RP, Nonogaki S, Kowalski LP, Nagai MA, Simpson AJ. A proposal for the integration of immunohistochemical staining and DNA-based techniques for the determination of TP53 mutations in human carcinomas. *Diagn Mol Pathol Am J Surg Pathol Part B* 2000;9:35–40.
- [48] Miquel-Cases A, Steuten LMG, Retèl VP, van Harten WH. Early stage cost-effectiveness analysis of a BRCA1-like test to detect triple negative breast cancers responsive to high dose alkylating chemotherapy. *Breast Edinb Scotl* 2015;24:397–405. doi:10.1016/j.breast.2015.03.002.
- [49] Moreno L, Linossi C, Esteban I, Gadea N, Carrasco E, Bonache S, et al. Germline BRCA testing is moving from cancer risk assessment to a predictive biomarker for targeting cancer therapeutics. *Clin Transl Oncol Off Publ Fed Span Oncol Soc Natl Cancer Inst Mex* 2016. doi:10.1007/s12094-015-1470-0.
- [50] Lobb EA, Barlow-Stewart K, Suthers G, Hallowell N. Treatment-focused DNA testing for newly diagnosed breast cancer patients: some implications for clinical practice. *Clin Genet* 2010;77:350–4. doi:10.1111/j.1399-0004.2009.01307.x.
- [51] Steuten LMG, Schutten M, Kip M, IJzerman MJ. Assessing The Impact Of A Multi-Million Public-Private Partnership For Translational Research: The Centre For Translational Molecular Medicine (Ctmm). *Value Health* 2014;17:A31. doi:10.1016/j.jval.2014.03.192.
- [52] Avar D, Bridge P, Buccì LM, Chiquette J, Dorval M, Durocher F, et al. Partnering in oncogenetic research--the INHERIT BRCA's experience: opportunities and challenges. *Fam Cancer* 2006;5:3–13. doi:10.1007/s10689-005-2570-8.
- [53] Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* 2012;490:61–70. doi:10.1038/nature11412.
- [54] Zardavas D, Irrthum A, Swanton C, Piccart M. Clinical management of breast cancer heterogeneity. *Nat Rev Clin Oncol* 2015;12:381–94. doi:10.1038/nrclinonc.2015.73.
- [55] Zardavas D, Irrthum A, Swanton C, Piccart M. Clinical management of breast cancer heterogeneity. *Nat Rev Clin Oncol* 2015;12:381–94. doi:10.1038/nrclinonc.2015.73.
- [56] Beije N, Jager A, Sleijfer S. Circulating tumor cell enumeration by the CellSearch system: the clinician's guide to breast cancer treatment? *Cancer Treat Rev* 2015;41:144–50. doi:10.1016/j.ctrv.2014.12.008.
- [57] Patel KM, Tsui DWY. The translational potential of circulating tumour DNA in oncology. *Clin Biochem* 2015;48:957–61. doi:10.1016/j.clinbiochem.2015.04.005.
- [58] Deenen MJ, Cats A, Beijnen JH, Schellens JHM. Part 4: pharmacogenetic variability in anticancer pharmacodynamic drug effects. *The Oncologist* 2011;16:1006–20. doi:10.1634/theoncologist.2010-0261.
- [59] Deenen MJ, Cats A, Beijnen JH, Schellens JHM. Part 3: Pharmacogenetic variability in phase II anticancer drug metabolism. *The Oncologist* 2011;16:992–1005. doi:10.1634/theoncologist.2010-0260.
- [60] Deenen MJ, Cats A, Beijnen JH, Schellens JHM. Part 2: pharmacogenetic variability in drug transport and phase I anticancer drug metabolism. *The Oncologist* 2011;16:820–34. doi:10.1634/theoncologist.2010-0259.
- [61] Deenen MJ, Cats A, Beijnen JH, Schellens JHM. Part 1: background, methodology, and clinical adoption of pharmacogenetics. *The Oncologist* 2011;16:811–9. doi:10.1634/theoncologist.2010-0258.
- [62] Galluzzi L, Vacchelli E, Bravo-San Pedro J-M, Buqué A, Senovilla L, Baracco EE, et al. Classification of current anticancer immunotherapies. *Oncotarget* 2014;5:12472–508.
- [63] Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011;144:646–74. doi:10.1016/j.cell.2011.02.013.
- [64] Zardavas D, Baselga J, Piccart M. Emerging targeted agents in metastatic breast cancer. *Nat Rev Clin Oncol* 2013;10:191–210. doi:10.1038/nrclinonc.2013.29.

- [65] Arnedos M, Vicier C, Loi S, Lefebvre C, Michiels S, Bonnefoi H, et al. Precision medicine for metastatic breast cancer: limitations and solutions. *Nat Rev Clin Oncol* 2015;12:693–704. doi:10.1038/nrclinonc.2015.123.
- [66] Barker AD, Sigman CC, Kelloff GJ, Hylton NM, Berry DA, Esserman LJ. I-SPY 2: an adaptive breast cancer trial design in the setting of neoadjuvant chemotherapy. *Clin Pharmacol Ther* 2009;86:97–100. doi:10.1038/clpt.2009.68.
- [67] Prota AE, Bargsten K, Zurwerra D, Field JJ, Díaz JF, Altmann K-H, et al. Molecular mechanism of action of microtubule-stabilizing anticancer agents. *Science* 2013;339:587–90. doi:10.1126/science.1230582.
- [68] Dumontet C, Jordan MA. Microtubule-binding agents: a dynamic field of cancer therapeutics. *Nat Rev Drug Discov* 2010;9:790–803. doi:10.1038/nrd3253.
- [69] Minotti G, Menna P, Salvatorelli E, Cairo G, Gianni L. Anthracyclines: molecular advances and pharmacologic developments in antitumor activity and cardiotoxicity. *Pharmacol Rev* 2004;56:185–229. doi:10.1124/pr.56.2.6.
- [70] Pang B, Qiao X, Janssen L, Velds A, Groothuis T, Kerkhoven R, et al. Drug-induced histone eviction from open chromatin contributes to the chemotherapeutic effects of doxorubicin. *Nat Commun* 2013;4:1908. doi:10.1038/ncomms2921.
- [71] Rottenberg S, Nygren AOH, Pajic M, van Leeuwen FWB, van der Heijden I, van de Wetering K, et al. Selective induction of chemotherapy resistance of mammary tumors in a conditional mouse model for hereditary breast cancer. *Proc Natl Acad Sci U S A* 2007;104:12117–22. doi:10.1073/pnas.0702955104.
- [72] Borst P, Wessels L. Do predictive signatures really predict response to cancer chemotherapy? *Cell Cycle Georget Tex* 2010;9:4836–40.
- [73] Fontanella C, Loibl S, von Minckwitz G. Clinical usefulness and relevance of intermediate endpoints for cytotoxic neoadjuvant therapy. *Breast Edinb Scotl* 2015;24 Suppl 2:S84–7. doi:10.1016/j.breast.2015.07.020.
- [74] von Minckwitz G, Fontanella C. Comprehensive Review on the Surrogate Endpoints of Efficacy Proposed or Hypothesized in the Scientific Community Today. *J Natl Cancer Inst Monogr* 2015;2015:29–31. doi:10.1093/jncimonographs/lgv007.
- [75] Hudis CA, Barlow WE, Costantino JP, Gray RJ, Pritchard KI, Chapman J-AW, et al. Proposal for standardized definitions for efficacy end points in adjuvant breast cancer trials: the STEEP system. *J Clin Oncol Off J Am Soc Clin Oncol* 2007;25:2127–32. doi:10.1200/JCO.2006.10.3523.
- [76] Gourgou-Bourgade S, Cameron D, Poortmans P, Asselain B, Azria D, Cardoso F, et al. Guidelines for time-to-event end point definitions in breast cancer trials: results of the DATECAN initiative (Definition for the Assessment of Time-to-event Endpoints in CANcer trials)†. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2015;26:873–9. doi:10.1093/annonc/mdv106.
- [77] Berruti A, Amoroso V, Gallo F, Bertaglia V, Simoncini E, Pedersini R, et al. Pathologic complete response as a potential surrogate for the clinical outcome in patients with breast cancer after neoadjuvant therapy: a meta-regression of 29 randomized prospective studies. *J Clin Oncol Off J Am Soc Clin Oncol* 2014;32:3883–91. doi:10.1200/JCO.2014.55.2836.
- [78] Buchanan J, Wordsworth S, Schuh A. Issues surrounding the health economic evaluation of genomic technologies. *Pharmacogenomics* 2013;14:1833–47. doi:10.2217/pgs.13.183.
- [79] Holtby H, Skowno JJ, Kor DJ, Flick RP, Uezono S. New technologies in pediatric anesthesia. *Paediatr Anaesth* 2012;22:952–61. doi:10.1111/pan.12007.
- [80] Kittel A, Di MA, Stewart H. Factors influencing the decision to abandon manual wheelchairs for three individuals with a spinal cord injury. *Disabil Rehabil* 2002;24:106–14.
- [81] Hardisty AR, Peirce SC, Preece A, Bolton CE, Conley EC, Gray WA, et al. Bridging two translation gaps: A new informatics research agenda for telemonitoring of chronic disease. *Int J Med Inf* 2011;80:734–44. doi:10.1016/j.ijmedinf.2011.07.002.
- [82] Cytryn KN, Patel VL. Reasoning about diabetes and its relationship to the use of telecommunication technology by patients and physicians. *Int J Med Inf* 1998;51:137–51.
- [83] Retèl VP, Joore MA, Linn SC, Rutgers EJT, van Harten WH. Scenario drafting to anticipate future developments in technology assessment. *BMC Res Notes* 2012;5:442. doi:10.1186/1756-0500-5-442.
- [84] Sanders PMH, Ijzerman MJ, Roach MJ, Gustafson KJ. Patient preferences for next generation neural prostheses to restore bladder function. *Spinal Cord* 2011;49:113–9. doi:10.1038/sc.2010.65.

- [85] Pecchia L, Martin JL, Ragozzino A, Vanzanella C, Scognamiglio A, Mirarchi L, et al. User needs elicitation via analytic hierarchy process (AHP). A case study on a Computed Tomography (CT) scanner. *BMC Med Inform Decis Mak* 2013;13:2. doi:10.1186/1472-6947-13-2.
- [86] Hummel JM, van Rossum W, Verkerke GJ, Rakhorst G. Medical technology assessment: the use of the analytic hierarchy process as a tool for multidisciplinary evaluation of medical devices. *Int J Artif Organs* 2000;23:782–7.
- [87] Hummel MJ, van Rossum W, Verkerke GJ, Rakhorst G. Assessing medical technologies in development. A new paradigm of medical technology assessment. *Int J Technol Assess Health Care* 2000;16:1214–9.
- [88] Haakma W, Steuten LMG, Bojke L, IJzerman MJ. Belief elicitation to populate health economic models of medical diagnostic devices in development. *Appl Health Econ Health Policy* 2014;12:327–34. doi:10.1007/s40258-014-0092-y.
- [89] Fukamachi K. New technologies for mechanical circulatory support: current status and future prospects of CorAide and MagScrew technologies. *J Artif Organs Off J Jpn Soc Artif Organs* 2004;7:45–57. doi:10.1007/s10047-004-0256-x.
- [90] Carrara S. Nano-bio-technology and sensing chips: new systems for detection in personalized therapies and cell biology. *Sensors* 2010;10:526–43. doi:10.3390/s100100526.
- [91] Hovorka R. Closed-loop insulin delivery: from bench to clinical practice. *Nat Rev Endocrinol* 2011;7:385–95. doi:10.1038/nrendo.2011.32.
- [92] Maruthur NM, Joy SM, Dolan JG, Shihab HM, Singh S. Use of the analytic hierarchy process for medication decision-making in type 2 diabetes. *PloS One* 2015;10:e0126625. doi:10.1371/journal.pone.0126625.
- [93] van Til JA, Dolan JG, Stiggelbout AM, Groothuis KCGM, IJzerman MJ. The use of multi-criteria decision analysis weight elicitation techniques in patients with mild cognitive impairment: a pilot study. *The Patient* 2008;1:127–35.
- [94] Ahn MJ, Zwikael O, Bednarek R. Technological invention to product innovation: A project management approach. *Int J Proj Manag* 2010;28:559–68. doi:10.1016/j.jproman.2009.11.001.
- [95] Holford N, Ma SC, Ploeger BA. Clinical trial simulation: a review. *Clin Pharmacol Ther* 2010;88:166–82. doi:10.1038/clpt.2010.114.
- [96] LeRouge C, Ma J, Sneha S, Tolle K. User profiles and personas in the design and development of consumer health technologies. *Int J Med Inf* 2013;82:e251–68. doi:10.1016/j.ijmedinf.2011.03.006.
- [97] Garmer K, Liljegren E, Osvalder A-L, Dahlman S. Application of usability testing to the development of medical equipment. Usability testing of a frequently used infusion pump and a new user interface for an infusion pump developed with a Human Factors approach. *Int J Ind Ergon* 2002;29:145–59. doi:10.1016/S0169-8141(01)00060-9.
- [98] Cavalcanti A, Shirinzadeh B, Kretly LC. Medical nanorobotics for diabetes control. *Nanomedicine Nanotechnol Biol Med* 2008;4:127–38. doi:10.1016/j.nano.2008.03.001.
- [99] Chávez-Santiago R, Balasingham I, Bergsland J. Ultrawideband Technology in Medicine: A Survey. *J Electr Comput Eng* 2012;2012:1–9. doi:10.1155/2012/716973.
- [100] Taketani F, Hara Y. Characteristics of spherical aberrations in 3 aspheric intraocular lens models measured in a model eye. *J Cataract Refract Surg* 2011;37:931–6. doi:10.1016/j.jcrs.2010.12.044.
- [101] van de Wetering G, Steuten LMG, von Birgelen C, Adang EMM, IJzerman MJ. Early Bayesian modeling of a potassium lab-on-a-chip for monitoring of heart failure patients at increased risk of hyperkalaemia. *Technol Forecast Soc Change* 2012;79:1268–79. doi:10.1016/j.techfore.2012.02.004.
- [102] Dong H, Buxton M. Early assessment of the likely cost-effectiveness of a new technology: A Markov model with probabilistic sensitivity analysis of computer-assisted total knee replacement. *Int J Technol Assess Health Care* 2006;22:191–202. doi:10.1017/S0266462306051014.
- [103] Hummel JM, Boomkamp ISM, Steuten LMG, Verkerke BGJ, IJzerman MJ. Predicting the health economic performance of new non-fusion surgery in adolescent idiopathic scoliosis. *J Orthop Res* 2012;30:1453–8. doi:10.1002/jor.22104.
- [104] McAteer H, Cosh E, Freeman G, Pandit A, Wood P, Lilford R. Cost-effectiveness analysis at the development phase of a potential health technology: examples based on tissue engineering of bladder and urethra. *J Tissue Eng Regen Med* 2007;1:343–9. doi:10.1002/term.36.

- [105] Wong WB, Ramsey SD, Barlow WE, Garrison LP, Veenstra DL. The value of comparative effectiveness research: projected return on investment of the RxPONDER trial (SWOG S1007). *Contemp Clin Trials* 2012;33:1117–23. doi:10.1016/j.cct.2012.08.006.
- [106] Sadatsafavi H, Niknejad B, Zadeh R, Sadatsafavi M. Do cost savings from reductions in nosocomial infections justify additional costs of single-bed rooms in intensive care units? A simulation case study. *J Crit Care* 2015. doi:10.1016/j.jcrc.2015.10.010.
- [107] Retèl VP, Grutters JPC, van Harten WH, Joore MA. Value of research and value of development in early assessments of new medical technologies. *Value Health J Int Soc Pharmacoeconomics Outcomes Res* 2013;16:720–8. doi:10.1016/j.jval.2013.04.013.
- [108] Grutters JPC, Abrams KR, de Ruyscher D, Pijls-Johannesma M, Peters HJM, Beutner E, et al. When to wait for more evidence? Real options analysis in proton therapy. *The Oncologist* 2011;16:1752–61. doi:10.1634/theoncologist.2011-0029.
- [109] Shell. Shell scenarios n.d. <http://www.shell.com/global/future-energy/scenarios/what-are-scenarios.html>.
- [110] Saaty TL. How to make a decision: The analytic hierarchy process. *Eur J Oper Res* 1990;48:9–26. doi:10.1016/0377-2217(90)90057-I.
- [111] Business dictionary n.d. <http://www.businessdictionary.com>.
- [112] Investopedia n.d. <http://www.investopedia.com/terms/r/returnoninvestment.asp>.

Supplementary material

Predictive biomarkers review

2

The evaluation of biomarkers for neoadjuvant chemotherapy is a complex issue. In particular the translation from preclinical work to enter early studies involves expertise from a wide background. As discussed the reasons for low rate of validated and used biomarkers may vary widely. The literature review allowed us to demonstrate and quantify particular issues in the literature for further discussion. Unfortunately, this quantification in itself is not perfect and neither are the choices that have to be made to obtain the database. Here we aim to specify these choices and particular issues that we were unable to solve 'objectively'. Some examples:

- To overcome issues in identification of biomarker studies that do not mention the word biomarker, one would need to come up with ways to find studies that actually contribute to the evidence for a specific interesting biomarker or broadly include studies that may yield biomarkers but that on average do not include clear evidence.
- To evaluate pre-clinical evidence, which (if present) is usually briefly described, one would need to dive deeply into the underlying studies. Conversely, when one wants to assess the validity of the study at hand, does one follow the line of thought of the authors or rely on re-analysis or re-interpretation of the presented data and how does one weigh studies when analysis and reporting vary. E.g. given 3 studies, one lacking pre-clinical evidence, 1 with small sample size and 1 without control group; do none of them qualify or is there something to learn while complete evidence has not been gathered/reported.

Systematic search

We searched in Pubmed and Embase using the search terms "breast cancer", "biological markers", "predictive", "and neoadjuvant" and "human". Only full-text articles published in English by 15 July 2015 were selected. The full search identified 1029 papers, of which we excluded 892 for not involving biomarkers for NACT measured in pre-treatment tissue (i.e., imaging, serum and/or post-treatment biomarkers), for being already accepted measures ER, PR, HER2 (subtyping) and ki67, for being prognostic biomarkers, for being non-interventional studies, or due to lack of access.

Database construction and analysis

To describe the studies we particularly focused on large issues that may make the studies less reliable. We described the biomarker, drug, study design and outcome (as reported by identification/validation of particular biomarker). We summarized on the gene/signature level. We did not go deeply into preclinical evidence or the particulars of the statistical analysis, other than noting that studies that investigate a predictive biomarker should contain a control group without the treatment of interest and that preferably interaction tests should be performed. We did not weigh the particular statistical analyses against each other nor did we judge the analysis or interpretation based on "expert opinion", but simply report whether a conclusion was drawn that a specific biomarker was interesting based on the reported statistics.

Interpretation

We tried to use objectively testable measures to describe the studies and identify issues. Studies may be interesting or contribute in other ways than strict analysis as predictive biomarker. Furthermore, the search may have missed advances occurring at the frontlines of Phase III RCTs.

(very) early HTA review*Systematic search*

The search for studies that used (very) early HTA applied to predictive biomarkers was also performed systematically in Pubmed. We decided to start this search by using the names of the biomarker that were investigated in more than 10 studies (according to our predictive biomarker review results). We expected that if very (early) HTA was used, it would be in biomarkers with the biggest bulk of clinical evidence. These biomarkers were p53, Topo2A, BCL2, BRCA1, EGFR. Each of these names (and other synonyms) were searched in combination with the terms "breast", "costs", "assessment", "users", "scenario", "experts", "cost-effectiveness. Furthermore, we performed additional searches with the term "multigene" and "predictive biomarker" instead of the particular biomarker names. These allowed exploring whether our initial search terms were narrowing the results. All the searches were performed also by including the term neoadjuvant. Only full-text articles published in English by 15 January 2016 were selected.

Database construction and analysis

Hits for each biomarker specific search were p53 (n=147), Topo2A (n=15), BCL2 (n=14), BRCA1 (n=110), multigene (n=50) and predictive biomarker (n=22). Papers published prior to 2000, that reported on risk prediction biomarkers or on the already established biomarkers ER/PR/HER2 were excluded. Furthermore, papers reporting on biomarkers' effectiveness, clinical expert guidelines or clinical reviews were also removed. These were already captured in the prior review or already common practice.

This resulted in 31 included studies. These were classified on 1) whether they described clinical, economic, ethical, organizational and/or patient/doctor related aspects, and 2) whether they were on very (early), early or mainstream HTA according to Figure 3. Most papers were clinical and reported on the comparison between different technologies to detect a biomarker, except one paper that reported a method to determine of cut-off values for the biomarker. These papers were classified as early HTA as they informed on biomarker and test development characteristics. One paper reported on organizational aspects like biomarker test uptake. These papers were considered early and very early HTA respectively. One study presented a cost-effectiveness analysis in early stages of biomarker development. Furthermore, in the BRCA1-like search we identified one study where key opinion leaders perceptions were collected. This study was considered early HTA. Last, we identified several reviews touching on all HTA aspects. Most of the identified papers used semi-qualitative HTA methods (reviews, surveys) and few quantitative methods (cost-effectiveness analysis) were used. As our main intention was to report on the use of (very) early HTA rather than systematically quantifying the number of studies on it, we did not count all studies on each type of application, but rather provided

examples of the type of studies found (see section *(Very) early HTA is not yet used to assess predictive biomarkers* on the main manuscript). As we found a very low numbers of relevant studies, we did not gather all results in a database.

Interpretation

Our objective was to provide an oversight of the use of (very) early HTA in predictive biomarker research by using NACT as a case study. Nonetheless our search covered other research settings than NACT as well. We are aware that our search may be limited to research performed in the academic setting as the use of (very) early HTA in private companies is not publicly available.

Brief summary on the HTA methods presented in table 1

HTA methods can be divided in those that help characterizing the candidates on a variety of aspects, and those that specifically inform on end users, effectiveness and cost-effectiveness aspects (see table 1).

Qualitative methods that inform on *various aspects* and are relevant at different stages of research are literature review, interviews, discussions, focus groups and surveys. Scenario analysis, which is a structured way to explore likely futures for the alternatives based on expectations that one has for the future, can hypothesize on ethical concerns on the use of a specific biomarker test. Scenario analysis can be combined with other methods, for instance economic methods and explore the cost-effectiveness consequences of those. Additional methods that are relevant in the biomarker translation phase are SWOT (strengths weakness opportunities and threats) & PEST (political economical social and technological) analysis, which are business tools developed to explore the capabilities and external influences in the development of a product, and the multi-path mapping tool, which helps understanding and drawing on the potential development paths of the tests' technology. Furthermore, the clinical trial simulator (CTS) method can explore the effects of specific design assumptions to the expected outcomes. Quantitative methods that inform on *various aspects* are the analytical hierarchical process (AHP) and the conjoint analysis (CA), which prioritize alternatives in a step-wise approach and via software that provides interactive support for group deliberations.

Specific methods to derive information on *end users* (patients and/or doctors) are user profile building, which may be more useful in the biomarker identification phase because is a method whereby looking at epidemiological data or using direct observation identifies expectations from end-users on a new application, and usability testing, which is expected more useful at the translation phases as it's a method that assesses experienced end-users opinions.

Table 1: Methods to gather data on HTA aspects (for specific definitions of each method see supplementary table 2).

Early HTA methods to gather data on:				Biomarker identification phase Very early HTA	Biomarker translation phase Early HTA	Biomarker validation phase Mainstream HTA
Various aspects	End users	Effectiveness	Cost/ cost-effectiveness	-Literature review[79] Interviews [79] [80], discussions/focus groups[81], surveys[82] with doctors or/and patients -Scenario analysis [83] -Conjoint analysis [84] on end-users -AHP [85] on end-users -AHP [86,87] on doctors -Other elicitation [88] on doctors	-Literature review [89–91] -Interviews, discussions/ focus groups, surveys with doctors or/and patients (same references as previous cell) -Conjoint analysis [84] on end-users -AHP [92] [93] on end-users -Scenario building [83] -SWOT & PEST [94] -Multi-Path Mapping [81] -Clinical trial simulator [95]	Most aspects will have already been assessed in previous steps. Only if emerging trends that were not taken into account emerge, a combination of prior methods can be used. -Clinical trial simulator [95]
				-User profile building [96]	-Usability testing [97]	
				-Effectiveness data from a similar -technology[79] -Computer simulation models [98–100]	-POP data expected available	-Trial data expected available
				-Early CEA [101–103] -Headroom analysis [101, 104]	-Early CEA [48] -ROI of RC Ts [105]/ of implementation [106] -Sensitivity analysis [16,24] -VOI [107] -ROA [108]	-Final CEA

Quantitative, Quantitative/Qualitative

Abbreviations: CEA= cost-effectiveness analysis; AHP= hierarchical analytical process; VOI= value of information analysis; ROA= real options analysis; RCT= randomized clinical trial; ROI= return on investment; HTA= health technology assessment; SWOT= Strength, Weaknesses, Opportunities and Threats; PEST= Political, Economic, Social and Technological; POP= Proof of principle; End users= doctors / patients. By review we also mean meta-analysis.

In very early and early stages of biomarker research data on *effectiveness* may not be available, or only pre-clinical evidence without link to patient outcomes. In translational stages, it may be that effectiveness data on alternative technologies to detect the biomarker is missing. Methods to specifically derive estimates on effectiveness are computer simulations models, which require the construction of complex models that link technological features with clinical outcomes, and simple extrapolation, which requires assuming the same effectiveness to that of a similar technology already used for a similar application.

Evidence on *economic* grounds can be gathered by a range of quantitative methods. In very early stages and early stages of research the headroom method can be used to determine the greatest price at which the healthcare provider might fund the biomarker test under study, and the health economic (HE) model can be used to calculate the expected cost-effectiveness of this. In very early stages, these information will be derived by using early expectations of health impact, derived from the previously cited methods, and costs, derived from similar technologies or expert elicitation, and of effectiveness. In early stages of development, when the first in-human studies data is available, this can be used as an estimate for effectiveness and to derive cost data. At this stage, calculating the return on investment (ROI) from a specific part of the research or even for biomarker implementation can be interesting, as one of the most expensive parts of research still has to follow. This consists of simple arithmetic calculations, on the expected monetary gains from the use of the biomarker test when deducted by the required investment.

Table 2: Definitions of complex HTA methods presented in Figure 3 and supplementary table 1

User profiles	"Structured ways of typifying a group of users in text and pictorial formats (i.e., conceptually modeling the end users). They attempt to "capture" the users' mental model comprising of their expectations, prior experience and anticipated behavior" [96]over-worked health care professionals and a growing patient base suffering from multiple chronic diseases, one of which is diabetes. Consumer health technologies (CHT).
Scenario analysis	Long term research planning by exploring alternatives views of the future and create plausible stories around them [109].
Usability testing	Tests to assess whether the design of a new device would increase usability compared to the existing one. It is also used to explore whether there are any further user requirements [97]" container-title": "International Journal of Industrial Ergonomics", "page": "145-159", "volume": "29", "issue": "3", "source": "CrossRef", "DOI": "10.1016/S0169-8141(01.
Multi-Path Mapping	Combines the understanding of the potential of the technology with creative thinking about possible futures. It is a graphic illustration of the step-wise developmental pathways of technology over time, accounting for uncertainty about how the future may unfold.
Conjoint analysis	Reveals trends in consumer preferences for competing products by presenting them as bundles of attributes [84].
Analytical hierarchical process	Prioritizes alternatives when multiple criteria must be considered by arranging its characteristics in a hierarchic structure. It Thus it helps capturing both subjective and objective aspects of a decision [110].
SWOT & PEST	SWOT analysis is a situation analysis in which internal strengths (S) and weaknesses (W) of a organization/product, and external opportunities (O) and threats (T) faced by it are closely examined to chart a strategy. PEST analysis is a situation analysis in which political-legal (government stability, spending, taxation), economic (inflation, interest rates, unemployment), socio-cultural (demographics, education, income distribution), and technological (knowledge generation, conversion of discoveries into products, rates of obsolescence) factors are examined to chart an organization's long-term plans [111].
Health impact assessment	A combination of methods and tools by which a product or intervention may be judged for its potential effects on the health of a population [5]" container-title": "Applied Health Economics and Health Policy", "page": "331-347", "volume": "9", "issue": "5", "source": "NCBI PubMed", "abstract": "Worldwide, billions of dollars are invested in medical product development and there is an increasing pressure to maximize the revenues of these investments. That is, governments need to be informed about the benefits of spending public resources, companies need more information to manage their product development portfolios and even universities may need to direct their research programmes in order to maximize societal benefits. Assuming that all medical products need to be adopted by the heavily regulated healthcare market at one point in time, it is worthwhile to look at the logic behind healthcare decision making, specifically, decisions on the coverage of medical products and decisions on the use of these products under competing and uncertain conditions. With the growing tension between leveraging economic growth through R&D spending on the one hand and stricter control of healthcare budgets on the other, several attempts have been made to apply the health technology assessment (HTA).
Early health economics modeling	A model that structures evidence on clinical and economic outcomes in a form that can help to inform decisions about clinical practices and healthcare resource allocations.
Headroom analysis	The incremental cost of the technology where it could still be cost-effective [104].
Real options analysis	Technique that provides guidance as to whether to adopt a technology now or postpone the decision to when additional evidence is available.
Value of information analysis	Method that provides guidance as to whether conduct an adoption now vs conducting it later after undertake further research to increase certainty around the decision. Furthermore, if further research is worthwhile, it guides towards the design and sample size of the trial give the best value for money.
Sensitivity analysis	Simulation analysis in which key quantitative assumptions (i.e., biomarker test performance) are changed systematically to assess their effect on the final outcome (i.e., cost-effectiveness) [111].
Return on investment	Profitability ratio that measures the effectiveness of an investment by measuring the amount of return of an investment relative to the investments' costs [112].

CHAPTER 3

Early stage cost-effectiveness analysis of a BRCA1-like test to detect triple negative breast cancers responsive to high dose alkylating chemotherapy

**Anna Miquel-Cases
Lotte MG Steuten
Valesca P Retèl
Wim H van Harten**

The Breast 2015, Aug;24(4):397-405.

Abstract

Purpose: Triple negative breast cancers (TNBC) with a BRCA1-like profile may benefit from high dose alkylating chemotherapy (HDAC). This study examines whether BRCA1-like testing to target effective HDAC in TNBC patients can be more cost-effective than treating all patients with standard chemotherapy. Additionally, we estimated the minimum required prevalence of BRCA1-like and the required positive predictive value (PPV) for a BRCA1-like test to become cost-effective.

Methods: Our Markov model compared 1) the incremental costs; 2) the incremental number of respondents; 3) the incremental number of Quality Adjusted Life Years (QALYs); and 4) the incremental cost-effectiveness ratio (ICER) of treating TNBC women with personalized HDAC based on BRCA1-like testing vs. standard chemotherapy, from a Dutch societal perspective and a 20-year time horizon, using probabilistic sensitivity analysis. Furthermore, we performed one-way sensitivity analysis (SA) to all model parameters, and two-way SA to prevalence and PPV. Data were obtained from a current trial (NCT01057069), published literature and expert opinions.

Results: BRCA1-like testing to target effective HDAC would presently not be cost-effective at a willingness-to-pay threshold of €80.000/QALY (€81.981/QALY). SAs show that PPV drives the ICER changes. Lower bounds for the prevalence and the PPV were found to be 58.5% and 73.0% respectively.

Conclusion: BRCA1-like testing to target effective HDAC treatment in TNBC patients is currently not cost-effective at a willingness-to-pay of €80.000/QALY, but it can be when a minimum PPV of 73% is obtained in clinical practice. This information can help test developers and clinicians in decisions on further research and development of BRCA1-like tests.

Introduction

The human and economic consequences of resistant triple negative breast cancer (TNBC) are substantial. In the Netherlands, first-line anthracycline-based treatment is ineffective in approximately 40% [1] of 2.797 TNBC women [2], generating additional therapy costs of ~17 Million (when treated, for instance, with Erbulin) [3]. Increasing first-line treatment effectiveness seems a promising way forward to decrease both patient morbidity and healthcare costs.

As TNBC is a heterogeneous disease [4], treatment effectiveness could possibly be increased by basing its therapeutic management on sub-classifications. One important example is the absence of BRCA1 gene functionality, also known as BRCA1-like tumors [5]. Approximately 68% of TNBC have this defect, which seems to confer them sensitivity to alkylating agent-based regimens. The largest published study so far (using carboplatin, thiopeta and cyclophosphamide) reports a protective effect of the alkylating regimen vs. standard (anthracyclines-based) chemotherapy (SC) in these tumors, yielding a hazard ratio of relapse free survival (RFS) of 0.17 (95% CI: 0.05-0.60, $p = 0.05$) [6]. Whether this positive result is due to the chemo-sensitivity of BRCA1-like tumors to one specific agent (e.g., carboplatin), the combination, or the fact that the drugs were given at high doses is not known. Yet, a similar patient series treated with high dose ifosfamide, carboplatin and epirubicine (a different intensive regimen containing two alkylators) and retrospectively tested for BRCA1-like, yielded similar promising results (hazard ratio of disease free survival (DFS) of 0.05, 95% CI: 0.01-0.38, $p = 0.003$)[7]. Thus, it seems that the BRCA1-like profile could serve as a predictive biomarker for high dose alkylating chemotherapy (HDAC) in TNBC.

Prevalence of BRCA1-like is approximated to be 68.000 per 100.000 TNBC [8]. Targeted use of HDAC in this subgroup could substantially improve health outcomes and reduce healthcare spending on ineffective treatment. Yet, HDAC requires peripheral blood progenitor cell transplant (PBPC) with mean costs per patient of €53.600 [9]. Added to the BRCA1-like testing costs, these represent the additional direct medical costs to society of testing and treating one BRCA1-like patient with personalized HDAC compared to SC. The question therefore is whether these additional costs are offset by the health benefits and the reduction in spending on ineffective treatments. A timely investigation of the relationship between the expected test performance characteristics, its potential clinical consequences and potential cost-effectiveness, is thus warranted.

In order to inform clinicians and developers of BRCA1-like tests that predict response to HDAC in TNBC, we performed an exploratory cost-effectiveness analysis to examine whether BRCA1-like testing to personalize HDAC can be cost-effective compared to current clinical practice.

Additionally, we estimated the minimum prevalence of BRCA1-like and the positive predictive value (PPV) required for a BRCA1-like test to render this strategy cost-effective.

Methods

Model overview and structure

3

We developed a Markov model (2010; Microsoft Corporation, Redmond, WA) to compare the health economic consequences of treating two identical cohorts of TNBC women aged 40 [8] by one of the following strategies: BRCA1-like testing followed by targeted treatment with HDAC (i.e., "BRCA1-like strategy") or no testing and standard (anthracycline based) chemotherapy treatment (i.e., "current practice"), from a Dutch societal perspective over a 20-year time horizon. Costs were calculated in 2013 Euros (€). Future costs and effects were discounted at a rate of 4% and 1.5% per year respectively, according to Dutch pharmacoeconomics guidelines [10].

BRCA1-like strategy: Patients were initially tested for BRCA1-like. Those with the biomarker were assigned to HDAC (4*FEC: Fluorouracil, epirubicin and cyclophosphamide, followed by 1*CTC: Cyclophosphamide, thiotepa and carboplatin), and those without the biomarker to SC (5*FEC). **Current practice:** All patients received 5*FEC. The mean duration of the intervention was of one year. Regimens were based on a previously published randomized clinical trial (RCT) comparing HDAC and SC efficacy in high risk breast cancer (BC) patients [11].

Patients were classified as "respondents" to the assigned chemotherapy when no relapse or death occurred within the first 5-years, and "non-respondents" in the case such an event occurred within the first 5-years. This time-frame was considered a reasonable limit to include all events related to chemotherapy response [1,12,13].

After the intervention, patients entered in the DFS health state of the Markov model (Fig. 1). From this state, transitions to the relapse (R, including local, regional, and distant relapse), death (D) and the same DFS health state were modeled. In year one, patients were assigned the costs and the health related quality of life (HRQoL) weights of the administered chemotherapy. During this year patients could die from toxic events (septicemia and heart failure [11]) or non-BC related events, but they could not relapse. From this year onwards, disease-free patients could relapse or die from a non-BC related event. Patients with a relapse received treatment and could 1) remain in this state and accrue the costs and HRQoL weights of the DFS health state, representing a "cured" relapse; or 2) die from BC or other unrelated cause. We assumed that patients could only develop one relapse.

Figure 1: Decision tree, Markov model and potential health economic consequences of BRCA1-like testing followed by personalized HDAC vs. current clinical practice. The decision analytic tree illustrates the two treatment pathways under study: 1) BRCA1-like testing followed by personalized HDAC and 2) treating all patients with (anthracycline based) SC. After the intervention, all patients enter the Markov model in the DFS state and they accumulate life years, QALYs and costs over a 20-year period based on the assigned transition probabilities. In the end, we expect the main health economic consequences to be driven by the costs and effectiveness of the treatment received in each patient subgroup. TNBC = triple negative breast cancer; HDAC = high dose alkylating chemotherapy, SC = standard chemotherapy; DFS = disease free survival; R = relapse.

Model input parameters

Model inputs for clinical effectiveness, transition probabilities (tp), and HRQoL-weights are presented in Table 1.

The BRCA1-like baseline prevalence was assumed 68%, as presented in literature [8]. The test's PPV (proportion of BRCA1-like patients responding to HDAC within the first 5-years) was assumed 72%. This was the average PPV of the BRCA1-like array comparative genomic hybridization (aCGH) test and the BRCA1-like multiplex ligation-dependent probe amplification (MLPA) tests. Both tests have been tested in the 60 TNBC samples from the publication of Vollebergh et al. [6]. The MLPA data is still internal data from the Netherlands Cancer Institute-Antoni van Leeuwenhoek Hospital (NKI-AVL). Based on patient level data from the same publication, we estimated the proportion of non-BRCA1-like patients and unselected TNBC patients respondents to SC to be 35%. The proportion of patients with toxic deaths after HDAC were derived from the previously mentioned RCT, which compared HDAC and SC efficacy in high risk BC [11].

The tp of RFS, the tp of BC specific survival (BCSS) and the tps of all-cause mortality for years 1, 2, 5, 10 and 20 were estimated as follows:

- tp of RFS for respondents were considered zero over the 20-year time horizon reflecting that respondents, by definition, do not relapse during the first 5-years, and having a relapse later on is unlikely [12].
- tp of RFS for non-respondents and the tp of BCSS for all patients were derived from two hypothetical survival curves of RFS and BCSS. These were constructed by making use of an exponential model and the assumption that at 5 years, 95% of the patients had an event, relapse or BC death respectively; $S(t) = \exp\{-kt\}$, where k is the hazard rate and t is time. This assumption was confirmed by an experienced oncologist of the NKI-AVL.
- tp of all-cause mortality on the survival curve of the cohort were modeled using Dutch life tables [14].

HRQoL weights were obtained from sources using the EuroQoL-5D questionnaire, and attributed to the DFS and R health states [15,16]. The HRQoL-weight for R is the average of local and distant relapse. We assumed that HRQoL was not affected by BRCA1-like testing.

Model costs include testing, chemotherapy, and health state specific costs, all calculated accounting for direct medical, direct non-medical - (i.e., traveling expenses), and productivity losses. Direct medical and direct non-medical costs were derived from literature, the NKI financial

department, and Dutch sources on resource use and unit prices [9,10,17,18]. Productivity losses were calculated using the friction cost method [19]. Foreign currencies were exchanged to 2013 euros [20], and the consumer price index was used to account for inflation [21]. A detailed cost break-down is presented in Table 2 and a textual description in the annex.

Outcomes

Model outcomes are: 1) the incremental costs; 2) the incremental number of respondents; 3) the incremental number of Quality Adjusted Life Years (QALYs); and 4) the incremental cost-effectiveness ratio (ICER). Incremental cost-effectiveness was assessed against a Willingness-to-Pay threshold (WTP) of €80.000 per QALY, as recommended in the Dutch pharmacoeconomics guidelines [22].

Sensitivity analyses

Probabilistic sensitivity analysis (PSA) was performed in order to quantify the decision uncertainty around the base case scenario by assigning distributions to all stochastic input parameters (see Tables 1 and 2). A beta distribution was assigned to clinical effectiveness parameters and transition probabilities, a normal distribution to utilities, and a log-normal distribution to costs. For costs parameters, we assumed 25% variance of the mean when empirical estimates of variance were not available. We run the analysis by using Monte Carlo simulation with 10.000 random samples from the pre-defined distributions. Cost-effectiveness acceptability curves (CEACs) were derived from these, to show the decision uncertainty surrounding the expected incremental cost-effectiveness. CEACs are presented at a range (€0 to €100.000) of WTP values for one additional QALY. Furthermore, we plotted the net benefit probability map (NBPM) [23] which shows the evolution of net health benefit over time.

Subsequently, a threshold SA was used to estimate 1) the minimum required prevalence, 2) the minimum required PPV, and 3) the combination, for the BRCA1-like strategy to be cost-effective. The values were initially varied in 20% intervals from 0 to a 100%. Finally, we narrowed the intervals until we found the prevalence (with one decimal place) where the ICER was €80.000/QALY. Furthermore, one-way SA was performed to all parameters, by varying them within one standard deviation of error, or a 25% of their base case value if this information was missing.

Table 1: Baseline values for clinical effectiveness parameters, transition probabilities and HRQoL-weights included in the Markov model.

Parameter	Baseline	SE	Distribution parameters	Distribution	Source
Clinical effectiveness					
Positive predictive value (PPV) of the BRCA1-like test	72%	23.00%	2.12, 1.01	Beta	[6]/NKI-AVL
Prevalence of BRCA1-like in TNBC	68%	23.00%	2.01, 0.77	Beta	[8]
Non BRCA1-like respondents to standard chemotherapy	35%	23.00%	1.13, 2.14	Beta	[6]
TNBC respondents to standard chemotherapy	35%	9.00%	9, 17	Beta	[6]
Toxic deaths due to high dose alkylating chemotherapy					
Septicemia	0.004	0.32%	2, 441	Beta	[11]
Heart failure	0.004	0.32%	2, 441	Beta	[11]
Transition probabilities					
Relapse free survival (RFS)					
Respondents	0	-	-	-	Expert opinion
Transition probability 1, 2, 5, 10 and 20 years					Expert opinion
Transition probability 1 year	0.451	8.00%	18.31, 22.31	Beta	Expert opinion
Transition probability 2 year	0.248	0.40%	2800.35, 8511.93	Beta	Expert opinion
Transition probability 5 year	0.092	1.00%	80.50, 793.22	Beta	Expert opinion
Transition probability 10 year	0.010	0.50%	4.40, 449.57	Beta	Expert opinion
Transition probability 20 year	0.0002	0.04%	0.43, 1727.13	Beta	Expert opinion
Breast cancer specific survival (BCSS)					
Transition probability 1 year	0	-	-	-	Expert opinion
Transition probability 2 year	0.451	8.00%	18.31, 22.31	Beta	Expert opinion
Transition probability 5 year	0.112	0.84%	157.11, 12510.14	Beta	Expert opinion
Transition probability 10 year	0.018	0.63%	7.89, 432.11	Beta	Expert opinion
Transition probability 20 year	0.0005	0.06%	0.63, 1377.88	Beta	Expert opinion
Utilities					
High dose alkylating chemotherapy	0.610	29.00%	0.61, 0.084	Normal ^b	[15]
Standard chemotherapy	0.620	3.93%	0.62, 0.002	Normal	[16]
Relapse ^a	0.732	1.63%	0.732, 0.0003	Normal	[16]
Disease free survival	0.779	3.06%	0.778, 0.001	Normal	[16]

SE = standard error

^a Calculated as an average of the utility of local relapse and the utility of distant relapse. ^b Truncated normal distribution bounded between 0 and 1.

Table 2: Baseline costs included in the Markov model

Input parameters	Unit costs	Unit measure	Mean resource use	Mean cost	SE	Distribution parameters (ln scale)	Source
BRCA1-like MLPA test	Direct medical costs						
	€9	Per sample ^a	24 ^b	€219	-	-	[33]
	€62	Per 7 samples	3.4	€212	-	-	NKI-AVL
	€25	Per hour	5.5	€137	-	-	[34]
	€40	Per hour	1	€40	-	-	[34]
	€15	Per run	1	€15	-	-	NKI-AVL
	€40	Per run	1	€40	-	-	NKI-AVL
	€3	Day	0	€0	-	-	[10]
	€251	Day	0	€0	-	-	[10]
	-	-	-	€664	-	-	-
	-	-	-	€37	10% ^c	3.61, 0.01	-
	Total per sample						
	-	-	-	€3556	-	-	-
Standard chemotherapy (5* FEC)	Direct medical costs						
	€176	1800 mg	2.2	€390	-	-	[35]
	€147	100 mg	7.2	€1062	-	-	[35]
	€45	1080 mg	3.7	€167	-	-	[35]
	€279	Day	5	€1393	-	-	[10]
	€109	Visit	5	€544	-	-	[35]
	€3	Day	5	€15	-	-	[10]
	€251	Day	25	€6272	-	-	[10]
	-	-	-	€9844	25%	9.19, 0.69	-
	Total						

High dose alkylating chemotherapy (4*FEC +1CTC)	4*FEC	Direct medical costs	-	-	-	€59901	-	-	-	[35]
		Fluorouracil	€176	1800 mg	-	€312	1.8	-	-	[35]
		Epirubicine	€147	100 mg	-	€850	5.8	-	-	[35]
		Cyclophosphamide	€45	1080 mg	-	€134	3	-	-	[35]
		Day care	€279	Day	-	€1114	4	-	-	[10]
		Oncologist visit	€109	Visit	-	€435	4	-	-	[35]
		Cyclophosphamide	€45	1080 mg	-	€401	8.9	-	-	[35]
	1*CTC	Carboplatin	€117	150 mg	-	€1996	17.1	-	-	[35]
		Thiotepa	€1021	1000 mg	-	€784	0.8	-	-	[18]
		Day care	€279	Day	-	€279	1	-	-	[10]
		PBPCT ^d harvesting	€13440	per patient	-	€13440	1	-	-	[9]
		PBPCT	€24682	per patient	-	€24682	1	-	-	[9]
		Post PBPCT ^e	€15476	per patient	-	€15476	1	-	-	[9]
		Direct non-medical costs ^f	€3	Day	-	€18	6	-	-	[10]
Other	Other	Loss of productivity costs ^g	€251	Day	-	€15555	62	-	-	[10]
		Total	-	-	-	€75472	25%	11.23, 1.07	-	-
Septicemia	Septicemia	Direct medical costs	€27330	Episode	-	€27330	1	-	-	[36]
		Direct non-medical costs	€3	Day	-	€3	1	-	-	[10]
		Loss of productivity costs	€251	Day	-	€5018	20	-	-	[10]
		Total	-	-	-	€32351	25%	10.38, 0.91	-	-
Heart failure	Heart failure	Direct medical costs	€31528	Episode	-	€31528	1	-	-	[38]
		Direct non-medical costs	€3	Day	-	€3	1	-	-	[10]
		Loss of productivity costs	€251	Day	-	€1505	6	-	-	[38]
		Total	-	-	-	€33036	25%	10.41, 0.91	-	-
Disease free state ^h	Disease free state ^h	Direct medical costs	-	-	-	€2872	-	-	-	[39]
		In & out -patient	€2793	Episode	-	€2793	1	20%	7.93, 0.03	[39]
		Drugs	€79	Episode	-	€79	1	25%	4.37, 0.01	[39]
		Loss of productivity costs ⁱ	€251	Day	-	€2352	9.4	25%	7.76, 0.44	[39]
		Total	-	-	-	€5225	-	-	-	[39]

Relapse state ^h	Local relapse	-	-	-	€22987	-	-	[39]
	Direct medical costs	-	-	-	€14833	-	-	[39]
	In & out -patient	€12497	Episode	1	€12497	14%	9.43, 0.01	[39]
	Drugs	€2336	Episode	1	€2336	25%	7.76, 0.44	[41]
	Loss of productivity costs ⁱ	€251	Day	32.5	€8154	25%	9.01, 0.66	[41]
	Distant relapse	-	-	-	€23313	-	-	[41]
	Direct medical costs	-	-	-	€17417	-	-	[41]
	In & out -patient	€11645	Episode	1	€11645	11%	9.36, 0.01	[41]
	Drugs	€5772	Episode	1	€5772	25%	8.66, 0.01	[41]
	Loss of productivity costs ⁱ	€251	Day	23.5	€5896	25%	8.68, 0.60	[41]
Breast cancer death state ^h	Total	-	-	-	€23150	-	-	-
	Direct medical costs	€8296	Episode	1	€8296	25%	9.02, 0.66	[41]
	Loss of productivity costs ⁱ	€251	Day	23.5	€5896	25%	8.68, 0.60	[41]
	Total	-	-	-	€14192	-	-	-

SE = standard error.

^a Each BRCA1-like MLPA test requires both patient and control samples, each of them costing V9 of MLPA kit (enzymes and reagents).

^b 6 control samples are added in each run. With an optimal sample size of 18 samples, this results in 24 samples.

^c Using the assumption of 25% variance of the mean reported value in a logarithmic scale resulted in a negative value, thus we used 10% instead.

^d Abbreviation for peripheral blood progenitor cell transplant.

^e Follow up period were the patient is controlled until recovery of blood activity.

^f Includes one trip to the hospital for each FEC cycle, and one trip to the hospital for PBPCT (admission and discharge).

^g We assumed patients did not work during chemotherapy (n = 20), during PBPCT procedures (n = 21) and during the post- PBPCT program (n = 20).

^h Source did not report travelling expenses, and thus, they were not added.

ⁱ Indirect costs were calculated by using resource use of Lidgren et al [39] and the friction method as recommended by the Dutch guidelines.

^j Loss of productivity was assumed to be the same as in the distant relapse health state

Results

Outcomes

Based on our PSA, the BRCA1-like strategy would cost an additional €76.369 per patient while increasing QALYs by 0.93 and the number of respondents by 25%, over a 20-year time horizon. Over this time-horizon, this strategy is expected to have an ICER of €81.981, which is not considered cost-effective. Yet decision uncertainty surrounding the ICER is substantial, with a 62% probability that the BRCA1-testing strategy is cost-effective (Fig. 2). The NBPM illustrates that the BRCA1-like strategy becomes cost-effective only after 20-years (Fig. 3).

Sensitivity analysis

The threshold SA demonstrated that the PPV, but not the prevalence, drives the ICER changes. Only when the PPV and prevalence values are well above 60% the strategy becomes cost-effective (Fig. 4). The minimum prevalence and PPV values at which BRCA1-like testing is expected to be just about cost-effective are 58.5% and 73.0% respectively. The one-way SA on the remaining model parameters indicated that the effectiveness parameters, the costs of HDAC and the utility of HDAC had the strongest impact on the ICER (Fig. 5) and can change the expectation of cost-effectiveness.

Figure 2: Cost effectiveness acceptability curves. The BRCA1-like strategy has a 62% probability to be cost-effective when compared to current practice.

Figure 3: Net benefit probability map. The BRCA1-like strategy becomes cost-effective only after 20 years, when the cost-effectiveness threshold is met.

Figure 4: Threshold sensitivity analysis (SA). a) one-way sensitivity analysis to the prevalence; b) one way SA to the PPV and the prevalence. The baseline values for the PPV (72%) and prevalence (67%) were derived from the 10,000 Monte Carlo simulations. The dots falling on the right side of the €80,000 per QALY threshold line are cost-effective results and those falling in the left side of the line are non-cost-effective results. The minimum prevalence and PPV values at which BRCA1-like testing is expected to be just about cost-effective are 58.5% and 73.0% respectively.

Figure 5: Tornado plot of one-way sensitivity analyses. The main drivers of the ICER are the effectiveness parameters, the costs of high dose alkylating chemotherapy and the utility of high dose alkylating chemotherapy.

Discussion

This study explored the costs and benefits of BRCA1-like testing followed by targeted treatment with HDAC in TNBC, in order to inform clinicians and developers of BRCA1-like tests on the requirements for this test to potentially become a cost-effective alternative to current clinical practice.

Our base case analysis indicates that the BRCA1-like strategy likely increases the number of respondents by 25% and the number of QALYs by 0.93 over a time horizon of 20-years. However, as indicated by the NBPM, these health benefits are only expected to outweigh the additional €76.369 costs per patient after 20-years, as the costs for testing and HDAC are made in the short term, and the health and financial benefits are recouped in the longer term. Furthermore, decision uncertainty around the ICER remains, and the BRCA1-like strategy is expected to be cost-effective at 20-years with a 62% probability. Threshold SA demonstrated that the PPV, but not the prevalence, drives the ICER, and the lower bounds for these two parameters for the strategy to be cost-effective are 58.5% (prevalence) and 73.0% (PPV).

Furthermore, we observed that the effectiveness parameters, the costs of HDAC and the utility of HDAC parameters can affect the cost-effectiveness of the BRCA1-like strategy.

To the best of our knowledge, this is the first exploratory analysis of the potential cost-effectiveness of BRCA1-like testing to target HDAC treatment in TNBC. The results can therefore not yet be compared to other cost-effectiveness estimations. However, key factors that drive economic value of stratified medicine have been described before and our findings are largely in line with those. Notably, as Trusheim et al. [24], we observed that the therapeutic effect within the biomarker positive population, the prevalence of the predictive biomarker and the clinical performance of the test drive stratified medicine's economic value. Specifically, we observed that with good therapeutic effect (tps of respondents) and clinical performance of the test (PPV) (note that in our model therapeutic effect in respondents was always good), the BRCA1-like strategy is expected to be cost-effective at a minimum required prevalence (in our study 58.5%). Furthermore, with low test performance, even if prevalence and therapeutic effect are perfect, no good economic value can be derived (Fig. 4).

Given that test performance is crucial for attaining economic value, it is important to realize that several tests for BRCA1-like detection are available [5]. Each test uses different aberrations to characterize the profile, which means that they may yield different results in terms of clinical effectiveness for specific applications. To our knowledge, the only tests used as predictors of sensitivity to HDAC in TNBC are the aCGH [6,25] and the MLPA [8,26], whose performance data we used in our PSA. Both tests are presently being validated, and from the few available data of these studies (internal NKI-AVL data) it seems that the PPVs for both tests are close to the lower bound of 73.0%.

From a policymaker's perspective, we highlight two important points. First, although incorporating HDAC treatment for TNBC is costly, if based on a BRCA1-like predictive test, the overall strategy costs can be justified by its long-term health benefits. This is of particular relevance to countries such as the United States, in which there is hesitance to cover HD chemotherapy [27,28]. Emergence of clinical and cost-effectiveness data on tests that can better target the usage of such costly treatment, may provide evidence to support coverage for those patients likely to respond. Risk sharing agreements and other reimbursement models might be needed to incentivize this appropriately for both the developers, the care providers and health insurers [29]. However, to support this scenario, further studies on this topic should be performed especially under a United States perspective. Second, although the adoption of a BRCA1-like test requires equipment and expertise to PBPCT, in the majority of Dutch centers that qualify, this would imply practice changes, but no monetary investments would be needed.

Our analysis indicated that the cost-effectiveness of the BRCA1-like strategy is affected by effectiveness parameters and costs. We therefore expect that further analysis of our model with

data from other studies using different HDAC regimens and different doses (i.e., the recently published cohort by Schouten et al. [7]) could result in different outcomes.

There are two important limitations of our study. First, we used assumptions for survival based on the TNBC subset of Vollebergh et al. [6]. Second, calculations of per test costs assumed optimal sample turnaround time, i.e. 18 samples per 10 days. Given the prevalence of TNBC in the BC population (2.797/year in the Netherlands [2]), this may be an optimistic assumption. That said, one-way SA reveals that test costs have little influence on the ICER.

Since we present an exploratory cost-effectiveness study performed in early stages of test development, we recommend subsequent cost-effectiveness analyses [30e32] to be performed once new data becomes available from clinical studies. For instance, from the on-going prospective validation study of the BRCA1-like MLPA test (NCT01057069). This study aims at providing evidence on the effectiveness of the BRCA1-like MLPA test to personalize HDAC (using the same regimen as the one used in this study) in TNBC. It can thus contribute information on transition probabilities, on BRCA1-like prevalence, MLPA test' PPV and costs.

Acknowledgments

The authors acknowledge the Center for Translational Molecular Medicine (CTMM, project Breast CARE, grant no.03O-104), source of funding for this project, and Prof. Dr. Sjoerd Rodenhuis, Mr. Philip Schouten, Dr. Petra M Nederlof and Dr. Esther H Lips for sharing their valuable insights regarding BRCA1-like testing in clinical practice.

References

- [1] Liedtke C, Mazouni C, Hess KR, Andre F, Tordai A, Mejia JA, et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol* 2008;26:1275e81. <http://dx.doi.org/10.1200/JCO.2007.14.4147>.
- [2] iKNL (Integraal Kankercentrum Nederland) Nederlandse Kankerregistratie. <http://www.cijfersoverkanker.nl/over-de-registratie-12.html>.
- [3] Cortes J, O'Shaughnessy J, Loesch D, Blum JL, Vahdat LT, Petrakova K, et al. Eribulin monotherapy versus treatment of physician's choice in patients with metastatic breast cancer (EMBRACE): a phase 3 open-label randomised study. *Lancet* 2011; 377:914e23. [http://dx.doi.org/10.1016/S0140-6736\(11\)60070-6](http://dx.doi.org/10.1016/S0140-6736(11)60070-6).
- [4] Metzger-Filho O, Tutt A, de Azambuja E, Saini KS, Viale G, Loi S, et al. Dissecting the heterogeneity of triple-negative breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol* 2012;30:1879e87. <http://dx.doi.org/10.1200/JCO.2011.38.2010>.
- [5] Chalasani P, Livingston R. Differential chemotherapeutic sensitivity for breast tumors with "BRCAness": a review. *Oncol* 2013;18:909e16. <http://dx.doi.org/10.1634/theoncologist.2013-0039>.
- [6] Vollebergh MA, Lips EH, Nederlof PM, Wessels LFA, Schmidt MK, van Beers EH, et al. An aCGH classifier derived from BRCA1-mutated breast cancer and benefit of high-dose platinum based chemotherapy in HER2-negative breast cancer patients. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2011;22:1561e70. <http://dx.doi.org/10.1093/annonc/mdq624>.
- [7] Schouten PC, Marme F, Aulmann S, Sinn H-P, Van Essen DF, Ylstra B, et al. Breast cancers with a BRCA1-like DNA copy number profile recur less often than expected after high-dose alkylating chemotherapy. *Clin Cancer Res Off J Am Assoc Cancer Res* 2015;21(4):763e70. <http://dx.doi.org/10.1158/1078-0432.CCR-14-1894>.
- [8] Lips EH, Mulder L, Oonk A, van der Kolk LE, Hogervorst FBL, Imholz ALT, et al. Triple-negative breast cancer: BRCAness and concordance of clinical features with BRCA1-mutation carriers. *Br J Cancer* 2013;108:2172e7. <http://dx.doi.org/10.1038/bjc.2013.144>.
- [9] Dutch Healthcare Authority (NZa.nl). DBC product-finder for tariffs [accessed 27.02.14], <http://www.nza.nl/organisatie/>; 2014.
- [10] Hakkaart L, Roijen van, Tan SS, Brouwmans CAM. Guide for research costs methods and standard cost prices for economic evaluations in healthcare \commissioned by the Health Care Insurance Board. Rotterdam. 2010.
- [11] Rodenhuis S, Bontenbal M, Beex LVAM, Wagstaff J, Richel DJ, Nooij MA, et al. High-dose chemotherapy with hematopoietic stem-cell rescue for high-risk breast cancer. *N Engl J Med* 2003;349:7e16. <http://dx.doi.org/10.1056/NEJMoa022794>.
- [12] Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, Sawka CA, et al. Triple negative breast cancer: clinical features and patterns of recurrence. *Clin Cancer Res* 2007;13:4429e34. <http://dx.doi.org/10.1158/1078-0432.CCR-06-3045>.
- [13] Pogoda K, Niwińska A, Murawska M, Pieńkowski T. Analysis of pattern, time and risk factors influencing recurrence in triple-negative breast cancer patients. *Med Oncol N Lond Engl* 2013;30:388. <http://dx.doi.org/10.1007/s12032-012-0388-4>.
- [14] Dutch National Center for Health Statistics, Centraal Bureau voor de Statistiek DH (in Dutch). <http://statline.cbs.nl>
- [15] Conner-Spady BL, Cumming C, Nabholz J-M, Jacobs P, Stewart D. A longitudinal prospective study of health-related quality of life in breast cancer patients following high-dose chemotherapy with autologous blood stem cell transplantation. *Bone Marrow Transpl* 2005;36:251e9. <http://dx.doi.org/10.1038/sj.bmt.1705032>.
- [16] Lidgren M, Wilking N, Jönsson B, Rehnberg C. Health related quality of life in different states of breast cancer. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil* 2007;16:1073e81. <http://dx.doi.org/10.1007/s1136-007-9202-8>.
- [17] Zorginstituut Nederland. Farmacotherapeutisch Kompas. 2013 [accessed 06.09.13], <http://www.fk.cvz.nl/>.
- [18] Zorginstituut Nederland. Medicijnkosten. 2013 [accessed 09.06.13], <http://medicijnkosten.nl/>.
- [19] Koopmanschap MA, Rutten FF, van Ineveld BM, van Roijen L. The friction cost method for measuring indirect costs of disease. *J Health Econ* 1995;14:171e89.
- [20] XE™. <http://www.xe.com/company/>. [accessed 11.11.13].

- [21] StatExtracts OECD. Consumer prices (MEI) e annual inflation [accessed 11.11.13], <http://stats.oecd.org/index.aspx?queryid%22519>; 2013.
- [22] Zorginstituut Nederland Het College voor Zorgverzekeringen. cvz.nl.
- [23] McCabe C, Edlin R, Hall P. Navigating time and uncertainty in health technology appraisal: would a map help? *PharmacoEconomics* 2013;31:731e7. <http://dx.doi.org/10.1007/s40273-013-0077-y>.
- [24] Trusheim MR, Burgess B, Hu SX, Long T, Averbuch SD, Flynn AA, et al. Quantifying factors for the success of stratified medicine. *Nat Rev Drug Discov* 2011;10:817e33. <http://dx.doi.org/10.1038/nrd3557>.
- [25] Joosse SA, van Beers EH, Tielen IHG, Horlings H, Peterse JL, Hoogerbrugge N, et al. prediction of BRCA1-association in hereditary non-BRCA1/2 breast carcinomas with array-CGH. *Breast Cancer Res Treat* 2009;116:479e89. <http://dx.doi.org/10.1007/s10549-008-0117-z>.
- [26] Lips EH, Laddach N, Savola SP, Vollebergh MA, Oonk AMM, Imholz ALT, et al. Quantitative copy number analysis by multiplex ligation-dependent probe amplification (MLPA) of BRCA1- associated breast cancer regions identifies BRCAness. *Breast Cancer Res BCR* 2011;13:R107. <http://dx.doi.org/10.1186/bcr3049>.
- [27] Weiss RB, Gill GG, Hudis CA. An on-site audit of the South African trial of high dose chemotherapy for metastatic breast cancer and associated publications. *J Clin Oncol Off J Am Soc Clin Oncol* 2001;19:2771e7.
- [28] Mello MM, Brennan TA. The controversy over high-dose chemotherapy with autologous bone marrow transplant for breast cancer. *Health Aff (Millwood)* 2001;20:101e17. <http://dx.doi.org/10.1377/hlthaff.20.5.101>.
- [29] Ramsey SD, Sullivan SD. A new model for reimbursing genome-based cancer care. *Oncol* 2014;19:1e4.<http://dx.doi.org/10.1634/theoncologist.2013-0392>.
- [30] Steuten LM, Ramsey SD. Improving early cycle economic evaluation of diagnostic technologies. *Expert Rev Pharmacoecon Outcomes Res* 2014;14:491e8. <http://dx.doi.org/10.1586/14737167.2014.914435>.
- [31] Vallejo-Torres L, Steuten L, Parkinson B, Girling AJ, Buxton MJ. Integrating health economics into the product development cycle: a case study of absorbable pins for treating hallux valgus. *Med Decis Mak Int J Soc Med Decis Mak* 2011;31:596e610. <http://dx.doi.org/10.1177/0272989X10388041>.
- [32] Vallejo-Torres L, Steuten LMG, Buxton MJ, Girling AJ, Lilford RJ, Young T. Integrating health economics modeling in the product development cycle of medical devices: a Bayesian approach. *Int J Technol Assess Health Care* 2008;24:459e64. <http://dx.doi.org/10.1017/S0266462308080604>.
- [33] MRC-Holland e MLPA Technology e MLPA an introduction. <https://mlpa.com/WebForms/WebFormMain.aspx?Tag%2fzjCZBtdOUyAt3KF3EwRZhNWLtcfv9pVl/tHJIM%5Cfa9FWO8KMqctOGloqYwxaGF9Y>. [accessed 01.07.14].
- [34] VSNU. Collective labour agreement dutch universities, 1 September 2007 to 1 March 2010. The Hague. 2008.
- [35] Frederix GW. Disease specific methods for economic evaluations of breast cancer therapies. University of Utrecht; 2013.
- [36] Davies A, Ridley S, Hutton J, Chinn C, Barber B, Angus DC. Cost effectiveness of drotrecogin alfa (activated) for the treatment of severe sepsis in the United Kingdom. *Anaesthesia* 2005;60:155e62. <http://dx.doi.org/10.1111/j.1365-2044.2004.04068.x>.
- [37] Schilling MB. Costs and outcomes associated with hospitalized cancer patients with neutropenic complications: a retrospective study. *Exp Ther Med* 2011;2(5):859e66. <http://dx.doi.org/10.3892/etm.2011.312>.
- [38] Wang G, Zhang Z, Ayala C, Wall HK, Fang J. Costs of heart failure-related hospitalizations in patients aged 18 to 64 years. *AmJManag Care* 2010;16:769e76.
- [39] Lidgren M, Wilking N, J€onsson B, Rehnberg C. Resource use and costs associated with different states of breast cancer. *Int J Technol Assess Health Care* 2007;23:223e31. <http://dx.doi.org/10.1017/S0266462307070328>.

Supplementary material

Testing costs

The costs of BRCA1-like testing were calculated based on the multiplex ligation-dependent probe amplification (MLPA) test, used in the NKI as part of prospective validation study (TNM study; NCT01057069). This test is suitable for clinical routine practice as it is robust, user-friendly, rapid and commercially available [15]. Costs of testing included (1) technician and laboratory costs to perform the test (material and overheads), (2) molecular biologist costs to interpret the results and generate reports, and (3) administration and depreciation costs. The costs of running the tests were calculated with the optimal test batching of 18 samples per 10 days. The purchasing costs for the MLPA kit were obtained from the MRC- Holland (Amsterdam, the Netherlands) website (SALSA MLPA P376 BRCA1ness probemix [26]). Other laboratory costs, administration and depreciation costs were derived from the financial department of the NKI-AVL, and the personnel costs from the collective labour agreement for Dutch hospitals [35].

Chemotherapy related costs

Medical direct costs of chemotherapy consisted of drug costs, day care costs and medical visit costs. We did not include the costs of radiotherapy because they were assumed equal under both regimens. The costs of chemotherapy were derived from and based on Dutch prices [12,36]. The costs associated to peripheral blood progenitor cell transplant (PBPC) procedures and subsequent follow up (in the HDAC arm) were derived from the Dutch Healthcare Authority's tariffs [11]. For both regimens we made two assumptions: (1) patients did not work during chemotherapy and (2) visits to the oncologist were scheduled during the chemotherapy days. Therefore, direct non-medical and productivity costs in the conventional regimen included the traveling costs on the days of chemotherapy and the 25 days missed at work. The direct non-medical and productivity costs in the HDAC regimen included one day of traveling costs for admission to the hospital, and productivity losses for 20 days of 4*FEC, 21 days hospitalized for 1*CTC/ PBPC and 21 days post-transplant were the patient is controlled until recovery of blood activity. The costs associated with toxic deaths under the HDAC regimen were obtained from literature [37-39].

Health states costs

The costs of the health states disease free survival (DFS) and relapse (R) were based on Lidgren et al. [39]. Cost of relapse was calculated as an average of local and distant relapse costs. The costs of death were excluded, unless it was consequence of treatment toxicity or breast cancer. In those situations we accounted for the specific costs to treat the toxicity (mentioned in the previous section) and for the palliative treatment.

CHAPTER 4

Decisions on further research for predictive biomarkers of high dose alkylating chemotherapy in triple negative breast cancer: A value of information analysis

Anna Miquel-Cases

Valesca P Retèl

Wim H van Harten

Lotte MG Steuten

Abstract

Objectives: Informing decisions about the design and priority of further studies of emerging predictive biomarkers of high-dose alkylating chemotherapy (HDAC) in triple negative breast cancer (TNBC), using Value of Information (VOI) analysis.

Methods: A state transition model compared treating TNBC women with current clinical practice and four biomarker strategies to personalize HDAC: 1) BRCA1-like by aCGH testing; 2) BRCA1-like by MLPA testing, 3) strategy-1 followed by XIST and 53BP1 testing; and 4) strategy-2 followed by XIST and 53BP1 testing, from a Dutch societal perspective and a 20-year time horizon. Input data came from literature and expert opinions. We assessed the expected value of (partial) perfect information (EV(P)PI), the expected value of sample information (EVSPI) and the expected net benefit of sampling (ENBS) for potential ancillary studies of an on-going randomized clinical trial (RCT; NCT01057069).

Results: EVPPIs indicate that further research should be prioritized to the parameter group including “biomarkers’ prevalence, positive predictive value (PPV), and treatment response rates (TRRs) in biomarker negative and TNBC patients” (€639M), followed by utilities (€48M), costs (€40M) and transition probabilities (tp) (€30M). By setting-up four ancillary studies to the on-going RCT, data on 1) tp and MLPA prevalence, PPV and TRR; 2) aCGH and aCGH /MLPA plus XIST and 53BP1 prevalence, PPV and TRR; 3) utilities; and 4) costs, could be simultaneously collected (optimal size =3000).

Conclusions: Further research on predictive biomarkers for HDAC should focus on gathering data on tps, prevalence, PPV, TRRs, utilities and costs from four ancillary studies to the on-going RCT.

Introduction

Triple negative breast cancer (TNBC) accounts for 15% to 20% of newly diagnosed breast cancer cases [1]. Currently, no targeted treatment exists for this subtype and standard chemotherapy is the guideline recommended treatment ([2–6]. While standard chemotherapy can be effective, 40% of TNBC patients suffer from early relapses and short post-recurrence survival [7,8]. Although second and third line treatments exist, these typically increase overall costs but do not contribute sufficiently to improve long term health outcomes [9–11]. Thereby, improving first-line treatment seems a promising way forward to decrease both patient morbidity and healthcare costs in this population.

As TNBC is a heterogeneous disease [12], treatment effectiveness could possibly be increased by basing its therapeutic management on sub-classifications. Pre-clinical data [13–15], and clinical data from a retrospective study conducted alongside a prospective randomized clinical trial (RCT) in our centre (the Netherlands Cancer Institute – Antoni van Leeuwenhoek hospital, NKI) [16], indicate that high-dose alkylating chemotherapy may be an effective treatment option for TNBC tumors without functional BRCA1, also known as BRCA1-like tumors. Furthermore, in an extension of this study, it was found that by further characterizing BRCA1-like tumors with two other biomarkers, XIST (X-inactive specific transcript gene) [20] and 53BP1 (tumor suppressor p-53 binding protein) [14,21,22], responses to high-dose alkylating chemotherapy treatment increase by 30%, i.e., patients with a BRCA-like profile, expression of 53BP1 (53BP1+) and low-expression of XIST (XIST-) have a 100% response rate compared to the 70% yielded with the BRCA1-like biomarker alone. Based on these results, a prospective RCT to test the survival advantage of treating TNBCs based on the BRCA1-like biomarker and high-dose alkylating chemotherapy was started (TNM-trial, NCT01057069). The trial started in 2010, and is currently on-going.

As the research on BRCA1-like, XIST and 53BP1 biomarkers is now progressing from initial clinical studies towards “pivotal” studies to determine its diagnostic, patient and societal value, early phase economic evaluation can be applied to improve the efficiency of the research and development process. Early phase economic evaluations are a decision analytic approach to iteratively evaluate technologies in development so as to increase their return on investment as well as patient and societal impact, when the technology becomes available [23]. For instance, value of information (VOI) methods quantify the potential benefit of additional information in the face of uncertainty. VOI is based on the idea that information is valuable because it reduces the expected costs of uncertainty surrounding a decision. A detailed explanation of the VOI methodology can be found elsewhere [24].

As decisions on emerging technologies with scarce clinical studies will inevitably be uncertain, research is expected to be worthwhile but only up to a certain cost of research. VOI methods allow to estimate an upper bound to the returns of further research expenditures and are particularly helpful in setting research priorities for specific model parameters as well as for specific research designs and sample sizes [25]. The data gathered in and the research infrastructure of the ongoing TNM-trial provides an opportunity to reduce uncertainty in a range of parameters that inform the decision problem, against additional costs. Therefore, this study aims to identify for which specific ancillary study designs further research is most valuable, and to inform future decisions on emerging predictive biomarkers for the selection of high-dose alkylating chemotherapy in TNBC.

4

Methods

A Markov model was constructed with three mutually exclusive health-states: disease free survival (DFS), relapse (R) (including local, regional, and distant relapses), and death (D). Our analysis took a Dutch societal perspective and a time horizon of 20-years, as the occurrence of relapses and deaths are expected within this time-frame [7,26–28]. Effectiveness was assessed in terms of quality-adjusted life-years (QALY) and costs in 2013 Euros (€). Future costs and effects were discounted to their present value by a rate of 4% and 1.5% per year respectively [29].

Patient population studied and strategies compared

We modelled five identical cohorts of 40-year old TNBC women, four treated with personalized high-dose alkylating chemotherapy as dictated by biomarkers and one treated according to current practice, with mean duration of 1-year (see figure 1 and description below). Drug regimens were based on a published RCT comparing high-dose alkylating chemotherapy and standard chemotherapy efficacy in breast cancer [30].

- 1) BRCA1-like tested by aCGH (array comparative genomic hybridisation) (BRCA1-like-aCGH): Women are initially tested for BRCA1-like by aCGH. Those who have a BRCA1-like profile are assigned to the high-dose alkylating chemotherapy arm (4*FEC: Fluorouracil, epirubicin and cyclophosphamide, followed by 1*CTC: Cyclophosphamide, thiotepa and Carboplatin), and those absent of the profile are assigned to standard chemotherapy (5*FEC);
- 2) BRCA1-like tested by MLPA (Multiplex Ligation-dependent Probe Amplification) (BRCA1-like-MLPA): MLPA was developed to be more time-efficient, cheaper, and technically less complicated than the aCGH [31]. We modelled this strategy exactly as the previous one;

- 3) BRCA1-like-aCGH followed by XIST and 53BP1 (BRCA1-like-aCGH/XIST-53BP1): Women are initially tested with the BRCA1-like-aCGH classifier, as above. Patients with a BRCA1-like profile are further tested for XIST and 53BP1 expression, and patients with a non-BRCA1-like profile receive standard chemotherapy. XIST expression is detected with a MLPA assay and 53BP1 by immunochemistry. These markers are interpreted together; BRCA1-like patients with a low expression of XIST and presence of 53BP1 are considered sensitive for high-dose alkylating chemotherapy and thus assigned to high-dose alkylating chemotherapy, and patients with any other combination of the markers are considered resistant and are assigned to standard chemotherapy;
- 4) BRCA1-like-MLPA followed by XIST and 53BP1 (BRCA1-like-MLPA/XIST-53BP1): This strategy was modelled exactly as the previous, but assessing BRCA1-like status by MLPA;
- 5) Current clinical practice: All women are treated with standard chemotherapy.

Patients are classified as “respondents” to the assigned chemotherapy when no relapse occurred within the first 5-years, and “non-respondents” in the case such an event occurred within the first 5-years. This time-frame was considered a reasonable limit to include all events related to chemotherapy response [7,8,33].

After the intervention, patients enter in the DFS health-state of the model, where they will remain for the 1st-year, accruing the costs and the health related quality of life (HRQoL) weights of the administered chemotherapy. During this year, patients can die from chemotherapy-related toxic events (septicemia and heart failure [30]) or non- breast cancer related events. Patients can move to the R health-state from the 1st-year onwards. Patients with a relapse receive treatment and can 1) remain in the R health-state and accrue the costs and HRQoL weights of the DFS health-state, representing a “cured” relapse; or 2) die from breast cancer or other unrelated cause. We assumed that patients could only develop one relapse during the time horizon of the model.

◀ Terminal node, patients enter the Markov process; MLPA, Multiplex Ligation-dependent Probe Amplification; aCGH, array Comparative Genomic Hybridization; XIST, X-inactive specific transcript gene; 53BP1, tumor suppressor p-53 binding protein; HDAC, High dose alkylating chemotherapy; Stand. Chemo, Standard chemotherapy.

Figure 1: Decision tree

Model input parameters

The baseline prevalence of BRCA1-like was derived from three patient series (n=377) in our hospital [34], including patients enrolled in the TNM-trial, and it was considered equal for both MLPA and aCGH tests. The baseline prevalence of BRCA1-like/XIST-/53BP1+ was determined from the existing retrospective study from a prospective RCT in our institute [16] (n=60), separately for the MLPA and the aCGH tests. This patient series was also used to derive 1) the PPV (proportion of biomarker positive patients responding to high-dose alkylating chemotherapy as determined by the MLPA and aCGH BRCA1-like tests alone, and by its combination with the XIST and 53BP1

tests; 2) the treatment response rates (TRRs) of biomarker negative patients as determined by the MLPA and aCGH BRCA1-like tests alone, and by its combination with the XIST and 53BP1 tests; and 3) the TRRs of TNBC patients.

The transition probabilities (tp) of relapse free survival (RFS) and breast cancer specific survival (BCSS) were estimated from Lester-Coll et al [35], in turn derived from survival data of Kennecke et al [27]. Using this data required making the assumption that most relapses in TNBC are metastatic, which is a plausible assumption given that in this subtype 1) metastatic disease is rarely preceded by other recurrences (Dent et al, Clin Cancer Res, 2007), and 2) there is low post-recurrence survival (Liedtke, JCO, 2008). All-cause mortality on the survival curve of the cohort was modelled using Dutch life tables [36].

The HRQoL weights were obtained from two studies reporting EuroQoL-5D utility weights [37,38]. During the 1st-year of the DFS health-state, patients were attributed the utility of the chemotherapy received (i.e., standard chemotherapy or high-dose alkylating chemotherapy and during the following 4-years, the HRQoL of DFS. In the 1st-year of the R health-state, patients were attributed the utility of R, and in subsequent years, the utility of DFS. We assumed that HRQoL was not affected by BRCA1-like testing itself.

Model costs include costs for biomarker testing, chemotherapy, and breast cancer health-states, each of them calculated as a sum of direct medical costs, indirect medical costs (e.g. patient travel expenses) and productivity losses. Direct medical and indirect medical costs were derived from literature, the NKI financial department, and Dutch sources on resource use and unit prices [29,39,40]. Productivity losses were calculated using the friction cost method [41]. Foreign currencies were exchanged to 2013 euros [42], and the consumer price index was used to account for inflation [43].

An overview of model parameters and sources are presented in table 1, and a detailed breakdown of the model costs can be found in the annex.

Table 1: Baseline prevalence, clinical effectiveness, tp, utilities and costs included in the markov model

Prevalence, clinical effectiveness, tp and utilities parameters				Source	Distribution	Parameters
Prevalence						
Prevalence BRCA1-like based on MLPA	68%	[31]	23%	[31,64]	Beta	(2.01, 1.01)
Prevalence BRCA1-like based on aCGH	68%	[31]	9%	[64]	Beta	(17.60, 8.41)
Prevalence BRCA1-like/XIST/53BP1+ based on MLPA	45%	[16]	11%	[16]	Beta	(9,11)
Prevalence BRCA1-like/XIST/53BP1+ based on aCGH	39%	[16]	10%	[16]	Beta	(9,14)
Clinical effectiveness						
PPV of the MLPA BRCA1-like test	72%	[16]	23%	[31,64]	Beta	(2.01, 0.77)
PPV of the aCGH BRCA1-like test	72%	[16]	9%	[64]	Beta	(17.14, 6.54)
PPV of the MLPA BRCA1-like test together with XIST and 53BP1 tests	100%	[16]	11%	[16]	Beta	(7,1)
PPV of the aCGH BRCA1-like test together with XIST and 53BP1 tests	100%	[16]	9%	[16]	Beta	(9,1)
TRR in non BRCA1-like respondents to SC by MLPA	35%	[16]	23%	[31,64]	Beta	(1.15, 2.14)
TRR in non BRCA1-like respondents to SC by aCGH	35%	[16]	9%	[64]	Beta	(9.42, 17.61)
TRR rates in TNBC respondents to SC	35%	[16]	9%	[16]	Beta	(9, 17)
Toxic deaths due to HDAC						
Septicemia	0.45%	[30]	0.32 %	[30]	Beta	(2,44)
Heart failure	0.45%	[30]	0.32 %	[30]	Beta	(2,44)
Transition probabilities^a						
Relapse free survival						
Respondents	0	Assum.	-	-	Fixed	-
Non-respondents	0.096	[35]	0.021	[35]	Beta	(19.37, 183.38)
	0.042	[35]	0.009	[35]	Beta	(18.96, 431.25)
Breast cancer specific survival						
Respondents & non-respondents	0	Assum.	-	-	Fixed	-
	0.681	[35]	0.042	[35]	Beta	(83.55, 39.09)
Utilities						
HDAC	0.610	[38]	29%	[38]	Normal truncated	(0.61, 0.08)
SC	0.620	[37]	4%	[37]	Normal	(0.62, 0.002)
Relapse ^b	0.732	[37]	3%	[37]	Normal	(0.73, 0)
Disease free survival	0.779	[37]	2%	[37]	Normal	(0.77, 0.001)

Cost parameters (log normal distribution)	Unit costs	Unit measure	Mean resource use	Mean cost	Source	SD (ln scale)	Source	Parameters (ln scale)
MLPA BRCA1-like test ^e	Direct medical costs							
	€ 9	Per sample ^c	24 ^d	€ 219	[32]	-	-	-
	€ 62	Per 7 samples	3.4	€ 212	NKI	-	-	-
	€25	Per hour	5.5	€137	[65]	-	-	-
	€40	Per hour	1	€40	[65]	-	-	-
	-	-	-	€609	-	-	-	-
aCGH BRCA1-like test ^e	-	-	-	€34	-	0.10	Assum. ^f	(3.52, 0.10)
	Direct medical costs							
	€26	1 reaction	13 ^g	€342	[66]	-	-	-
	€ 62	Per 1 sample	12 ^h	€750	NKI	-	-	-
	€25	Per hour	3.4	€137	[65]	-	-	-
	€40	Per hour	5.5	€40	[65]	-	-	-
MLPA XIST test ^e	-	-	-	€1,270	-	-	-	-
	-	-	-	€106	-	0.16	Assum.	(4.66, 0.03)
	Direct medical costs							
	€ 6	Per sample ^c	24 ^d	€ 153	[32]	-	-	-
	€ 62	Per 7 samples	3.4	€ 212	NKI	-	-	-
	€25	Per hour	5.5	€137	[65]	-	-	-
IHC 53BP1 test ^e	€40	Per hour	1	€40	[65]	-	-	-
	-	-	-	€543	-	-	-	-
	-	-	-	€30	-	0.10	Assum. ^f	(3.41, 0.01)
	Direct medical costs							
	€21.72	Per run	1	€21.72	[40]	-	-	-
	€0.71	Per run	1	€0.71	[40]	-	-	-
SC (5* FEC)	-	-	-	€22	-	0.10	Assum. ^f	(3.11, 0.01)
	Direct medical costs							
	€176	1800 mg	2.2	€3,556	-	-	-	-
	€147	100 mg	7.2	€390	[67]	-	-	-
	€45	1080 mg	3.7	€1,062	[67]	-	-	-
	€279	Day	5	€167	[67]	-	-	-
Day care				€1,393	[29]	-	-	-

HDAC (4*FEC +1CTC)	Oncologist visit	€109	Visit	5	€544	[67]	-	-	-
	Direct non-medical costs	€3	Day	5	€15	[29]	-	-	-
	Loss of productivity costs	€251	Day	25	€6.272	-	-	-	-
	Total	-	-	-	€9.844	-	0.83	Assum.	(9.19, 0.69)
	Direct medical costs	-	-	-	59.901€	-	-	-	-
	Fluorouracil	€176	1800 mg	1.8	€312	[67]	-	-	-
	Epirubicine	€147	100 mg	5.8	€850	[67]	-	-	-
	Cyclophosphamide	€45	1080 mg	3	€134	[67]	-	-	-
	Day care	€279	Day	4	€1.114	[29]	-	-	-
	Oncologist visit	€109	Visit	4	€435	[67]	-	-	-
1*CTC	Cyclophosphamide	€45	1080 mg	8.9	€401	[67]	-	-	-
	Carboplatin	€117	150 mg	17.1	€1.996	[67]	-	-	-
	Thiotepa	€1.021	1000 mg	0.8	€784	[68]	-	-	-
	Day care	€279	Day	1	€279	[29]	-	-	-
	PBPCT harvesting	€13.440	per patient	1	€13.440	[40]	-	-	-
	PBPCT	€24.682	per patient	1	€24.682	[40]	-	-	-
	Post PBPCT ⁱ	€15.476	per patient	1	€15.476	[40]	-	-	-
	Direct non-medical costs ^j	€3	Day	6	€18	[29]	-	-	-
	Loss of productivity costs ^k	€251	Day	62	€15.555	[29]	-	-	-
	Total	-	-	-	€75.472	-	1.03	Assum.	(11.23, 1.07)
Septicemia	Direct medical costs	€27.330	Episode	1	€27.330	[69]	-	-	-
	Direct non-medical costs	€3	Day	1	€3	[29]	-	-	-
	Indirect costs	€251	Day	20	€5.018	[29]	-	-	-
	Total	-	-	-	€32.501	-	0.95	Assum.	(10.34, 0.91)
	Direct medical costs	€31.528	Episode	1	€31.528	[71]	-	-	-
Heart failure	Direct non-medical costs	€3	Day	1	€3	[29]	-	-	-
	Loss of productivity costs	€251	Day	6	€1.505	[71]	-	-	-
	Total	-	-	-	€33.036	[29,71]	0.96	Assum.	(10.40, 0.91)
	Direct medical costs	-	-	-	€2.872	-	-	-	-
	In & out –patient	€2.793	Episode	1	€2.793	[72]	0.17	[72]	(7.93, 0.03)
Disease free state ^l	Drugs	€79	Episode	1	€79	[72]	0.09	Assum.	(4.37, 0.01)
	Loss of productivity costs ^m	€251	Day	9.4	€2.352	[72]	0.66	Assum.	(7.76, 0.44)

Relapse state ¹	Total	-	-	-	€5.225	[72]	-	-	-
	Local relapse	-	-	-	€22.987	[72]	-	-	-
	Direct medical costs	-	-	-	€14.833	[72]	-	-	-
	In & out -patient	€12.497	Episode	1	€12.497	[72]	0.12	[72]	(9.43, 0.01)
	Drugs	€2.336	Episode	1	€2.336	[72]	0.66	Assum.	(7.76, 0.44)
	Loss of productivity costs ^m	€251	Day	32.5	€8.154	[72]	0.81	Assum.	
Distant relapse	Direct medical costs	-	-	-	€23.313	[72]	-	-	-
	In & out -patient	€11.645	Episode	1	€11.645	[72]	0.10	[72]	(9.36, 0.01)
	Drugs	€5.772	Episode	1	€5.772	[72]	0.77	Assum.	(8.66, 0.01)
	Loss of productivity costs ^m	€251	Day	23.5	€5.896	[72]	0.77	Assum.	(8.68, 0.60)
Breast cancer death state ¹	Direct medical costs	€8.296	Episode	1	€8.296	[72]	0.81	Assum.	(9.02, 0.66)
	Loss of productivity costs ⁿ	€251	Day	23.5	€5.896	[72]	0.77	Assum.	(8.68, 0.60)
	Total	-	-	-	€14.192	[72]	-	-	-

Abbreviations: SD, standard deviation; MLPA, Multiplex Ligation-dependent Probe Amplification; aCGH, array Comparative Genomic Hybridization; XIST, X-inactive specific transcript gene; 53BP1, tumor suppressor p-53 binding protein; IHC, Immunohistochemistry; PPV, positive predictive value; TRR, treatment response rates; SC, standard chemotherapy; TNBC, triple negative breast cancer; HDAC, high dose alkylating chemotherapy; PBPC, Peripheral Blood Progenitor Cell Transplantation. Assum, standard deviation is equal to 25% of the mean.

Parameters for the distributions: Beta distribution: α/β , Normal distribution: mean/variance, Log-normal distribution: Log mean/log SD

^a Based on expert opinion, the 5-years RFS and 5 years -BCSS were assumed to vary from a minimum of 0% to a maximum of 10% , with 5% as baseline.

^b Calculated as an average of the utility of local relapse and the utility of distant relapse.

^c Each BRCA1-like MLPA test requires both patient and control samples, each of them costing € 9 of MLPA kit (enzymes and reagents).

^d The MLPA test requires 6 control samples and 1 patient sample in each run. With an optimal sample size of 18 samples, this results in 24 samples.

^e Indirect costs in test are zero.

^f Using the assumption of 25% standard deviation of the mean reported value in a logarithmic scale resulted in a negative value, thus we used 10% instead.

^g The aCGH test requires labelling of 12 patient samples and 1 control sample in each run.

^h We assumed optimal test batching of 12 patient samples in each run.

ⁱ Follow up period were the patient is controlled until recovery of blood activity

^j Includes one trip to the hospital for each FEC cycle, and one trip for the hospital for PBPC (admission & discharge).

^k We assumed patients did not work during chemotherapy (n=20), during PBPC procedures (n=21) and during the post- PBPC program (n=20).

^l Source did not report travelling expenses thus were not added.

^m Indirect costs were calculated by using resource use of Lidgren and the friction method as recommended by the Dutch guidelines.

ⁿ Loss of productivity was assumed to be the same as in the distant relapse health state.

Estimating decision uncertainty

Parameter uncertainty was quantified in the decision model by assigning distributions to all parameters that are subject to sampling uncertainty. Following the recommendations by Briggs et al [24], a beta distribution was assigned to binomial data, such as biomarkers' prevalence, PPVs, tps and TRRs in biomarker negative and TNBC patients, and a log-normal distribution to right-skewed data, such as costs. For uncertainty in mean utilities, we followed Brennan et al [44], suggesting the use of a normal distribution. As sampling from one utility distribution (HDAC) occasionally produced a parameter value below zero, this was truncated. The parameterization of each distribution can be derived from table 1. Uncertainty ranges for BRCA1-like-MLPA and BRCA1-like-aCGH prevalence, and for TRR in non-BRCA-1 like patients under both tests came from literature on the tests' development. This reported a 14% error of the MLPA vs. aCGH test [34] and an 11% of the aCGH test vs mutation status (gold standard) [45]. Uncertainty in the remaining binomial parameters were derived from the patient series of Vollebergh et al [16], except for tp. For these, alpha and beta parameters were derived from Lester-Coll et al [35], which were in turn derived by applying the method of moments to Kennecke et al survival data [27]. For the utility data, either the standard error, or the 95% confidence intervals of the mean, were derived from literature. As limited information regarding parameter uncertainty is available for costs, we assumed that standard errors of the aggregate costs were equal to 25% of the mean. However, if on the logarithmic scale this resulted in negative values, 10% was used. As literature to characterize uncertainty on specific items of the health state aggregate costs existed, this was used accordingly in these separate items, with the former assumptions being made for the remaining items of the aggregate value. The joint parameter uncertainty was then propagated through the model using Monte Carlo simulation with 10.000 random samples from the pre-defined distributions. Cost-effectiveness acceptability curves (CEACs) were estimated to show the joint decision uncertainty surrounding the expected incremental cost-effectiveness across €0 to €80.000 willingness-to-pay values for one additional QALY.

Value of further research and research priorities

The EVPI was calculated for the population expected to benefit from a reduction of uncertainty, TNBC patients eligible for high-dose alkylating chemotherapy i.e., patients below 60 years old with stage II-IV treatable cancers. The model assumes that the entire affected population will receive the optimal strategy. In the Netherlands the affected population amounts to 662 patients per annum (of the 6619 breast cancer women below 60 years in the Netherlands [46], 20% are expected to be TNBC [28,47–50], of these, 30% are stage II-III [51] and 20% are oligometastatic cancers [52] i.e., treatable metastatic cases). To this figure, an annual discount rate of 4% was applied over a 10-year time horizon of the technology, assumed to be the period during which

the information is relevant to inform the decision. The EVPPI requires two-level Monte Carlo simulation [24], beginning with an outer loop (100) sampling values from the distribution of the parameters of interest, and an inner loop (1000) sampling the remaining parameters from their conditional distribution [44]. The parameters groups of interests were determined based on the type of study design required for further research: 1) RCT to inform the tp, 2) QoL survey to provide further information regarding utility weights associated with chemotherapy and breast cancer health-states, 3) longitudinal costing study to provide more information on resource use of the tests, the chemotherapy and the health-states, and 4) longitudinal study to provide more information on the biomarkers' prevalence, PPVs, and the TRRs of biomarker negative and TNBC patients [24].

Research designs for further research

In this study we prioritize specific further research designs, designs depending on what type of data are needed and their vulnerability to specific risks of bias, and on the research infrastructure that is available from the TNM-trial, an on-going Dutch RCT aiming to provide evidence on the survival advantage (in terms of RFS and overall survival) of treating TNBC BRCA1-like patients as detected by MLPA with high-dose alkylating chemotherapy vs. standard chemotherapy. Thereby, further research was proposed as follows:

Further data on tp, BRCA1-like prevalence, BRCA1-like PPV and TTRs in biomarker negative and TNBC as identified by MLPA were assumed to come at the expenses of the TNM-trial, with the only additional costs of more advanced statistical analysis methods than planned for the original trial (this was defined as study1). Evidence on BRCA1-like prevalence as determined by aCGH, BRCA1-like/XIST/53BP1+ prevalence as determined by MLPA and aCGH, and TTRs in biomarker negative and TNBC as identified by aCGH could be derived from undertaking a retrospective study using the TNM-trial samples. To determine the prevalence, patient samples would first be tested by aCGH. Subsequently, those resulting BRCA1-like would be tested by 53BP1 and XIST. To determine the PPV and TTR in each case, additional statistical analysis correlating presence/absence of biomarker with survival data would be performed. The costs for this study would include re-testing patient samples and additional statistical analysis (study2). Evidence on direct medical costs could also be gathered from a retrospective study to the TNM-trial. In this study resource use and unit costs for the relevant parameters would be determined, incurring costs for data collection and statistical analysis (study3). Evidence on QoL could be derived from an ancillary prospective survey to the TNM-trial. Expenses resulting from this trial would be distributing, collecting and analyzing the QoL surveys' (study4).

Testing costs for the aCGH, 53BP1 and XIST biomarkers were derived from the financial department of the NKI (€30 for XIST testing; €22 for 53BP1 testing; and €106 for aCGH testing). The costs of performing statistical analysis only, of performing additional data collection and statistical analysis, and of performing a QoL survey were based on the costs of data management and analysis of a mock RCT presented in literature [53]. From this source, we specifically used the average of ‘academic medical and cancer centers’ costs and ‘oncology group practices’ costs. The total costs per patient were estimated at €1.325 for study 1, at €1.466 for study 2 (including €141 for XIST and 53BP1 testing in 68% BRCA1-like patients and aCGH testing to all patients, and €1.325 for the statistical analysis), and at €1.325 for each study 3 and 4. The EVSI was calculated for each of the four studies for a range of sample sizes, starting from 100, using a two-level Monte Carlo simulation with 5.000 inner and 5.000 outer loops (the number of loops was increased sequentially to check for convergence i.e., that increasing simulation size (both inner and outer) would not change estimates). The expected net benefit of sampling (ENBS) was subsequently calculated for each study design and n , by subtracting the corresponding costs of research. The n where the ENBS is maximized is the optimal sample size for each proposed study¹. Furthermore, we calculated the optimal sample size for the portfolio of studies, by assuming that these are undertaken simultaneously and results of one cannot inform results of others. Under this assumption, the optimal sample size is the combination of sample sizes across studies that maximizes the ENBS [24].

Results

Uncertainty in cost-effectiveness

The BRCA1-like-MLPA/XIST-53BP1, the BRCA1-like-aCGH/XIST-53BP1 and the BRCA1-like-aCGH strategies are expected to be cost-effective at a WTP of €80.000/QALY, when compared to current clinical practice, the BRCA1-like-MLPA/XIST-53BP1 and the BRCA1-like-MLPA strategy respectively. On the contrary, the additional costs of the BRCA1-like-MLPA strategy were not balanced by the gain in health outcomes, when compared to the BRCA1-like-aCGH/XIST-53BP1, resulting in an ICER of €94.310/QALY. The CEACs show that at a willingness-to-pay threshold of €80.000/QALY the decision as to which strategy is most cost-effective is uncertain. The base case results and the CEACs are presented in figure 2.

¹ Note that the costs of research always accounted for the same costs, even for sample sizes larger than the TNM-trial ($n=270$). It was assumed that other future RCTs with similar characteristics to the TNM-trial could be used to continue deriving the required data via equally designed retrospective studies.

Value of further research and research priorities

Results of the EVPI and EVPPIs are presented in figure 3. The EVPI was estimated at €693M at the prevailing threshold of €80.000/QALY. The EVPPI identified the group of parameters including the “biomarkers’ prevalence, the PPVs, and TRRs in biomarker negative and TNBC patients” to be most uncertain (€639M), followed by utilities (€48M), cost-related parameters (€40M) and tp (€30M).

Research designs for further research

In figure 4 we present graphically the ENBS and optimal sample size for the four proposed studies separately. These were €600M and 9000 for study 1, €440M and 1000 for study 2, €597M and 200 for study 3 and €446M and 1000 for study 4. The optimal sample size for the portfolio of studies was 3000, with an ENBS of €2074M.

	Life years (LY)	Quality adjusted life years (QALYs)	Costs (€)	ICER (€/QALY)
Current clinical practice	12.23	9.38	78.311	
BRCA1-like-MLPA/XIST-53BP1	13.23	10.14	122.032	57.673
BRCA1-like-aCGH/XIST-53BP1	13.47	10.33	126.831	25.384
BRCA1-like-MLPA	13.91	10.66	157.706	94.310
BRCA1-like-aCGH	13.93	10.67	159.080	74.643

Figure 2: Base case results and cost-effectiveness acceptability curves. The strategies are listed in order of increasing costs. In evaluating the ICERs, each strategy's costs and effects where compared with those of the strategy just slightly more expensive.

Figure 3: EVPI and EVPI estimates

Figure 4: ENBS and optimal sample size for each of the four ancillary study to the on-going RCT.

Discussion

This study found that testing for BRCA1-like alone with the aCGH test, and testing for BRCA1-like in combination with the biomarkers XIST and 53BP1, with the aCGH and the MLPA tests, may be cost-effective, and that there is substantial value in investing in further research for these diagnostic tests. VOI analysis showed that setting up four ancillary studies to the current TNM-trial to collect data on: 1) tp and MLPA prevalence, PPV and TRR; 2) aCGH and aCGH /MLPA plus XIST and 53BP1 prevalence, PPV and TRR; 3) utilities; and 4) costs, would be most efficient in generating information that decreases decision uncertainty around the test and treat strategies. The optimal sample size to simultaneously collect data from these four groups of parameters was 3000 patients, with and ENBS of €2074M.

This paper contributes to the literature on real-time applications of EVSI analysis to design and prioritize further research, which is under-represented [54–58]. Groot Koerkamp et al [55]

previously presented an EVSI application in a diagnostic procedure, but most EVSI analyses are applied to treatment interventions. Enhancing the literature on the expected value of further information about diagnostics is relevant for manufacturers, because current regulations incentivize research and development of diagnostics relatively poorly [59]. In the meantime, EVSI examples can illustrate how diagnostics' R&D can be steered more efficiently to increase the returns on investments from a healthcare and societal perspective. While many articles indicate the RCT to be the preferred study design to conduct any further research by default, we contribute to the literature in presenting the value of further research for various study designs, depending on what type of data are needed, the risk of bias and existing research infrastructure.

Apart from the fact that requiring RCTs for all forms of further data collections cannot inherently be justified in a rational way, there are two external motivations to consider the ENBS of non-RCT designs: 1) the evidence requirements for market approval and reimbursement of diagnostics, which are generally less rigidly defined compared to pharmaceuticals, therefore allowing to utilize valuable other sources of evidence; and 2) lower levels of evidence than RCTs are increasingly acceptable to decision-makers, as for example recently stated by the FDA [60].

When calculating the EVSI of study designs other than RCTs, parameter vulnerability to selection bias needs to be assessed. While this may be of less concern for costs and health-states utility data, selection bias in retrospective and/or observational studies can severely affect effectiveness parameters (like TRRs, and PPVs) and should be prevented or statistically accounted for. The use of retrospective studies alongside RCTs are increasingly promoted as these can generate high-quality evidence while being fast and inexpensive [61]. This is however only possible for diagnostics of already existing chemotherapeutic regimens, where data on efficacy is already available from RCTs.

Our study was not exempt of limitations. First, by nature of the early stage analysis, the input data on biomarkers' prevalence, biomarkers' PPV and TRRs in biomarker negative and TNBC patients was derived from several small retrospective studies. Indeed, EVPPI analysis showed high value in collecting further information on these, and our ENBS analysis suggest how this could be done most efficiently. Second, the TNM-trial uses intensified alkylating chemotherapy instead of high-dose alkylating chemotherapy. Although this means that the therapy is administered more frequently (2x) and at lower doses (half), it results in equal cumulative doses and equal need for stem-cell transplantation. Thereby, the survival advantage is expected to be similar. Third, the costs of testing were estimated by using optimal test batching; probably an optimistic assumption considering the prevalence of TNBC in the breast cancer population. However, it is not expected that this would markedly alter the conclusions of the analysis, as in a previous analysis of our model [62] testing costs were not a key driver of outcomes. Fourth, the research costs used for the ENBS calculations are derived from the published costs of a typical though hypothetical RCT [53]. While these estimates seem reasonable for a real trial, the use of actual

costs may change the results. Fifth, the estimated costs of study2 ignore the different accuracy of the aCGH and MLPA tests. Although this could translate in additional XIST and 53BP1 testing to derive the prevalence and PPV under the BRCA1-like-aCGH/XIST-53BP1 strategy, we expected these costs to be minimal. Sixth, the EVPI is dependent on estimates of population size, the time horizon, and the discount rate. We based these parameters on the Dutch situation, yet results to other countries requires reconsideration of these inputs. Seventh, it is possible that other biomarkers to predict sensitivity to high-dose alkylating chemotherapy will be identified in the future. This would add additional comparator(s) to the decision problem, thus increasing EVPI and probably the need for further research. Thereby, this type of analysis needs to be repeated over time (iterative process), in order to keep up with the latest developments. Furthermore, biases in early phase evidence are expected, when their design and conduct are not as rigorous as a large RCT. In this situation it is important to characterize the extent of uncertainty, as VOI is highly sensitive to this [63]. While we justified our data sources for both mean values and their variance, and explained data assumptions thoroughly, we did not conduct additional sensitivity analyses on the resulting parameter distributions [63]. Last, while we accounted for the correlation between the most important cost-effectiveness drivers sensitivity and specificity by using the Dirichlet distribution, we acknowledge that correlations may be present in other input parameters. This could impact the EVPI results and hence the EVSI estimates, with a magnitude depending on the strength of input correlation ([64]). We suggest that sophisticated methods that explicitly quantify joint distributions of correlated parameters are considered in further VOI analysis.

To conclude, this study illustrated the use of full Bayesian VOI analysis in a set of diagnostic tests, where further research was designed depending on the type of data needed and its vulnerability to specific risks of bias, and on the research infrastructure available from an on-going RCT.

References

- [1] Bauer KR, Brown M, Cress RD, Parise CA, Caggiano V. Descriptive analysis of estrogen receptor (ER)-negative, progesterone receptor (PR)-negative, and HER2-negative invasive breast cancer, the so-called triple-negative phenotype: a population-based study from the California cancer Registry. *Cancer* 2007;109:1721–8. doi:10.1002/cncr.22618.
- [2] National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology: Breast Cancer v2. 2011.
- [3] Goldhirsch A, Wood WC, Coates AS, Gelber RD, Thurlimann B, Senn H-J, et al. Strategies for subtypes—dealing with the diversity of breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2011. *Ann Oncol* 2011;22:1736–47. doi:10.1093/annonc/mdr304.
- [4] Aebi S, Davidson T, Gruber G, Castiglione M, ESMO Guidelines Working Group. Primary breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2010;21 Suppl 5:v9–14. doi:10.1093/annonc/mdq159.
- [5] Prise en charge du cancer du sein infiltrant de la femme non ménopausée. *Oncologie* 2009;11:507–32. doi:10.1007/s10269-009-1818-6.
- [6] Integraal Kankercentrum Nederland (IKNL). Breast Cancer Guideline, NABON 2012. n.d.
- [7] Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, Sawka CA, et al. Triple-Negative Breast Cancer: Clinical Features and Patterns of Recurrence. *Clin Cancer Res* 2007;13:4429–34. doi:10.1158/1078-0432.CCR-06-3045.
- [8] Liedtke C, Mazouni C, Hess KR, Andre F, Tordai A, Mejia JA, et al. Response to Neoadjuvant Therapy and Long-Term Survival in Patients With Triple-Negative Breast Cancer. *J Clin Oncol* 2008;26:1275–81. doi:10.1200/JCO.2007.14.4147.
- [9] Montero AJ, Avancha K, Glück S, Lopes G. A cost-benefit analysis of bevacizumab in combination with paclitaxel in the first-line treatment of patients with metastatic breast cancer. *Breast Cancer Res Treat* 2012;132:747–51. doi:10.1007/s10549-011-1919-y.
- [10] Lopes G, Glück S, Avancha K, Montero AJ. A cost effectiveness study of eribulin versus standard single-agent cytotoxic chemotherapy for women with previously treated metastatic breast cancer. *Breast Cancer Res Treat* 2013;137:187–93. doi:10.1007/s10549-012-2326-8.
- [11] Reed SD, Li Y, Anstrom KJ, Schulman KA. Cost effectiveness of ixabepilone plus capecitabine for metastatic breast cancer progressing after anthracycline and taxane treatment. *J Clin Oncol Off J Am Soc Clin Oncol* 2009;27:2185–91. doi:10.1200/JCO.2008.19.6352.
- [12] Metzger-Filho O, Tutt A, de Azambuja E, Saini KS, Viale G, Loi S, et al. Dissecting the heterogeneity of triple-negative breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol* 2012;30:1879–87. doi:10.1200/JCO.2011.38.2010.
- [13] Quinn JE, Kennedy RD, Mullan PB, Gilmore PM, Carty M, Johnston PG, et al. BRCA1 functions as a differential modulator of chemotherapy-induced apoptosis. *Cancer Res* 2003;63:6221–8.
- [14] Rottenberg S, Nygren AOH, Pajic M, van Leeuwen FWB, van der Heijden I, van de Wetering K, et al. Selective induction of chemotherapy resistance of mammary tumors in a conditional mouse model for hereditary breast cancer. *Proc Natl Acad Sci U S A* 2007;104:12117–22. doi:10.1073/pnas.0702955104.
- [15] Fong PC, Boss DS, Yap TA, Tutt A, Wu P, Mergui-Roelvink M, et al. Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med* 2009;361:123–34. doi:10.1056/NEJMoa0900212.
- [16] Vollebergh MA, Lips EH, Nederlof PM, Wessels LFA, Schmidt MK, van Beers EH, et al. An aCGH classifier derived from BRCA1-mutated breast cancer and benefit of high-dose platinum-based chemotherapy in HER2-negative breast cancer patients. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2011;22:1561–70. doi:10.1093/annonc/mdq624.
- [17] Philip C, Schouten. BRCA1 like copy number profiles to predict benefit of intensified alkylating chemotherapy in breast cancer. *ASCO Annu. Meet.* 2013 *J Clin Oncol* 31 2013 Suppl Abstr 11023, 2013.
- [18] Fasano J, Muggia F. Breast cancer arising in a BRCA-mutated background: therapeutic implications from an animal model and drug development. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2009;20:609–14. doi:10.1093/annonc/mdn669.

- [19] Chalasani P, Livingston R. Differential Chemotherapeutic Sensitivity for Breast Tumors With “BRCAness”: A Review. *The Oncologist* 2013;18:909–16. doi:10.1634/theoncologist.2013-0039.
- [20] Rottenberg S, Vollebergh MA, de Hoon B, de Ronde J, Schouten PC, Kersbergen A, et al. Impact of intertumoral heterogeneity on predicting chemotherapy response of BRCA1-deficient mammary tumors. *Cancer Res* 2012;72:2350–61. doi:10.1158/0008-5472.CAN-11-4201.
- [21] Bunting SF, Calln E, Wong N, Chen H-T, Polato F, Gunn A, et al. 53BP1 inhibits homologous recombination in Brca1-deficient cells by blocking resection of DNA breaks. *Cell* 2010;141:243–54. doi:10.1016/j.cell.2010.03.012.
- [22] Bouwman P, Aly A, Escandell JM, Pieterse M, Bartkova J, van der Gulden H, et al. 53BP1 loss rescues BRCA1 deficiency and is associated with triple-negative and BRCA-mutated breast cancers. *Nat Struct Mol Biol* 2010;17:688–95. doi:10.1038/nsmb.1831.
- [23] Ijzerman MJ, Steuten LMG. Early assessment of medical technologies to inform product development and market access: a review of methods and applications. *Appl Health Econ Health Policy* 2011;9:331–47. doi:10.2165/11593380-000000000-00000.
- [24] Briggs A, Claxton K, Sculpher M. Decision Modelling for Health Economic Evaluation. *Handbooks in Health Economic Evaluation*. Oxford, UK: Oxford University Press; n.d.
- [25] Steuten LM, Ramsey SD. Improving early cycle economic evaluation of diagnostic technologies. *Expert Rev Pharmacoecon Outcomes Res* 2014;14:491–8. doi:10.1586/14737167.2014.914435.
- [26] Cheang MCU, Voduc D, Bajdik C, Leung S, McKinney S, Chia SK, et al. Basal-like breast cancer defined by five biomarkers has superior prognostic value than triple-negative phenotype. *Clin Cancer Res Off J Am Assoc Cancer Res* 2008;14:1368–76. doi:10.1158/1078-0432.CCR-07-1658.
- [27] Kennecke H, Yerushalmi R, Woods R, Cheang MCU, Voduc D, Speers CH, et al. Metastatic Behavior of Breast Cancer Subtypes. *J Clin Oncol* 2010;28:3271–7. doi:10.1200/JCO.2009.25.9820.
- [28] Voduc KD, Cheang MCU, Tyldesley S, Gelmon K, Nielsen TO, Kennecke H. Breast cancer subtypes and the risk of local and regional relapse. *J Clin Oncol Off J Am Soc Clin Oncol* 2010;28:1684–91. doi:10.1200/JCO.2009.24.9284.
- [29] L. Hakkaart - van Roijen, S.S Tan, CAM Brouwmans. Guide for research costs - Methods and standard cost prices for economic evaluations in healthcare \ commissioned by the Health Care Insurance Board. Rotterdam: 2010.
- [30] Rodenhuis S, Bontenbal M, Beex LVAM, Wagstaff J, Richel DJ, Nooij MA, et al. High-dose chemotherapy with hematopoietic stem-cell rescue for high-risk breast cancer. *N Engl J Med* 2003;349:7–16. doi:10.1056/NEJMoa022794.
- [31] Lips EH, Laddach N, Savola SP, Vollebergh MA, Oonk AMM, Imholz ALT, et al. Quantitative copy number analysis by Multiplex Ligation-dependent Probe Amplification (MLPA) of BRCA1-associated breast cancer regions identifies BRCAness. *Breast Cancer Res BCR* 2011;13:R107. doi:10.1186/bcr3049.
- [32] MRC-Holland. SALSA MLPA P376 BRCA1ness probemix 2013. <http://www.mlpa.com/WebForms/WebFormMain.aspx?Tag=fNPBLedDVp38pCxU2h0mQ> (accessed November 7, 2013).
- [33] Pogoda K, Niwiska A, Murawska M, Piekowski T. Analysis of pattern, time and risk factors influencing recurrence in triple-negative breast cancer patients. *Med Oncol Northwood Lond Engl* 2013;30:388. doi:10.1007/s12032-012-0388-4.
- [34] Lips EH, Mulder L, Oonk A, van der Kolk LE, Hogervorst FBL, Imholz ALT, et al. Triple-negative breast cancer: BRCAness and concordance of clinical features with BRCA1-mutation carriers. *Br J Cancer* 2013;108:2172–7. doi:10.1038/bjc.2013.144.
- [35] Lester-Coll NH, Lee JM, Gogineni K, Hwang W-T, Schwartz JS, Prosnitz RG. Benefits and risks of contralateral prophylactic mastectomy in women undergoing treatment for sporadic unilateral breast cancer: a decision analysis. *Breast Cancer Res Treat* 2015;152:217–26. doi:10.1007/s10549-015-3462-8.
- [36] Dutch National Center for Health Statistics n.d. http://statline.cbs.nl/StatWeb/publication/?VW=T&DM=SLNL&PA=7052_95&D1=0-1,7,30-31,34,38,42,49,56,62-63,66,69-71,75,79&D2=0&D3=0&D4=a,!0-28&HD=080509-0829&HDR=G2,G1,G3&STB=T.
- [37] Lidgren M, Wilking N, Jnsson B, Rehnberg C. Health related quality of life in different states of breast cancer. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil* 2007;16:1073–81. doi:10.1007/s11136-007-9202-8.

- [38] Conner-Spady BL, Cumming C, Nabholz J-M, Jacobs P, Stewart D. A longitudinal prospective study of health-related quality of life in breast cancer patients following high-dose chemotherapy with autologous blood stem cell transplantation. *Bone Marrow Transplant* 2005;36:251–9. doi:10.1038/sj.bmt.1705032.
- [39] Dutch National Health Care Institute. CVZ n.d. cvz.nl.
- [40] Dutch Healthcare Authority (NZA.nl). DBC product-finder for tariffs 2014. <http://www.nza.nl/organisatie/> (accessed February 27, 2014).
- [41] Koopmanschap MA, Rutten FF, van Ineveld BM, van Roijen L. The friction cost method for measuring indirect costs of disease. *J Health Econ* 1995;14:171–89.
- [42] XE currency Converter n.d. <http://www.xe.com/>.
- [43] OECD.Stat. OECD (2013) n.d. <http://stats.oecd.org/index.aspx?queryid=22519#>.
- [44] Brennan A, Kharroubi S, O'hagan A, Chilcott J. Calculating partial expected value of perfect information via Monte Carlo sampling algorithms. *Med Decis Mak Int J Soc Med Decis Mak* 2007;27:448–70. doi:10.1177/0272989X07302555.
- [45] Joosse SA, van Beers EH, Tielen IHG, Horlings H, Peterse JL, Hoogerbrugge N, et al. Prediction of BRCA1-association in hereditary non-BRCA1/2 breast carcinomas with array-CGH. *Breast Cancer Res Treat* 2009;116:479–89. doi:10.1007/s10549-008-0117-z.
- [46] iKNL (Integraal Kankercentrum Nederland). Nederlandse Kankerregistratie n.d. <http://www.cijfersoverkanker.nl/overde-registratie-12.html>.
- [47] Carey LA, Perou CM, Livasy CA, Dressler LG, Cowan D, Conway K, et al. Race, breast cancer subtypes, and survival in the Carolina Breast Cancer Study. *JAMA* 2006;295:2492–502. doi:10.1001/jama.295.21.2492.
- [48] Hu Z, Fan C, Oh DS, Marron JS, He X, Qaqish BF, et al. The molecular portraits of breast tumors are conserved across microarray platforms. *BMC Genomics* 2006;7:96. doi:10.1186/1471-2164-7-96.
- [49] Fan C, Oh DS, Wessels L, Weigelt B, Nuyten DSA, Nobel AB, et al. Concordance among gene-expression-based predictors for breast cancer. *N Engl J Med* 2006;355:560–9. doi:10.1056/NEJMoa052933.
- [50] Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* 2012;490:61–70. doi:10.1038/nature11412.
- [51] Ries L, Eisner M, Kosary C. SEER cancer statistics review, 1973-1999. *Natl Cancer Inst* 2002.
- [52] Jain SK, Dorn PL, Chmura SJ, Weichselbaum RR. Incidence and implications of oligometastatic breast cancer. vol. 30, 2012, p. Suppl; abstr e11512.
- [53] Emanuel EJ. The Costs of Conducting Clinical Research. *J Clin Oncol* 2003;21:4145–50. doi:10.1200/JCO.2003.08.156.
- [54] Girling AJ, Freeman G, Gordon JP, Poole-Wilson P, Scott DA, Lilford RJ. Modeling payback from research into the efficacy of left-ventricular assist devices as destination therapy. *Int J Technol Assess Health Care* 2007;23:269–77. doi:10.1017/S0266462307070365.
- [55] Groot Koerkamp B, Nikken JJ, Oei EH, Stijnen T, Ginai AZ, Hunink MGM. Value of information analysis used to determine the necessity of additional research: MR imaging in acute knee trauma as an example. *Radiology* 2008;246:420–5. doi:10.1148/radiol.2462070093.
- [56] Ramsey SD, Blough DK, Sullivan SD. A forensic evaluation of the National Emphysema Treatment Trial using the expected value of information approach. *Med Care* 2008;46:542–8. doi:10.1097/MLR.0b013e318160b479.
- [57] McKenna C, McDaid C, Suekarran S, Hawkins N, Claxton K, Light K, et al. Enhanced external counterpulsation for the treatment of stable angina and heart failure: a systematic review and economic analysis. *Health Technol Assess Winch Engl* 2009;13:iii – iv, ix – xi, 1–90. doi:10.3310/hta13240.
- [58] Stevenson MD, Oakley JE, Lloyd Jones M, Brennan A, Compston JE, McCloskey EV, et al. The cost-effectiveness of an RCT to establish whether 5 or 10 years of bisphosphonate treatment is the better duration for women with a prior fracture. *Med Decis Mak Int J Soc Med Decis Mak* 2009;29:678–89. doi:10.1177/0272989X09336077.
- [59] Towse A, Garrison LP. Economic Incentives for Evidence Generation: Promoting an Efficient Path to Personalized Medicine. *Value Health* 2013;16:S39–43. doi:10.1016/j.jval.2013.06.003.
- [60] US Food and Drug Administration (FDA). Guidance Documents (Medical Devices and Radiation-Emitting Products) > Guidance for Industry and Food and Drug Administration Staff - Factors to Consider When Making Benefit-Risk Determinations in Medical Device Premarket Approvals and De Novo Classifications 2012. <http://www.fda.gov/medicaldevices/deviceregulationandguidance/guidancedocuments/ucm267829.htm> (accessed September 23, 2014).

- [61] Simon RM, Paik S, Hayes DF. Use of archived specimens in evaluation of prognostic and predictive biomarkers. *J Natl Cancer Inst* 2009;101:1446–52. doi:10.1093/jnci/djp335.
- [62] Miquel-Cases A, Steuten LMG, Retèl VP, van Harten WH. Early stage cost-effectiveness analysis of a BRCA1-like test to detect triple negative breast cancers responsive to high dose alkylating chemotherapy. *Breast Edinb Scotl* 2015;24:397–405. doi:10.1016/j.breast.2015.03.002.
- [63] Madan J, Ades AE, Price M, Maitland K, Jemutai J, Revill P, et al. Strategies for efficient computation of the expected value of partial perfect information. *Med Decis Mak Int J Soc Med Decis Mak* 2014;34:327–42. doi:10.1177/0272989X13514774.

Supplementary material

Testing costs

The costs of testing with MLPA (for BRCA1-like and XIST) and aCGH (for BRCA1-like) were estimated from the local experience of the NKI. Those included (1) technician and laboratory costs to perform the test (material and overheads), and (2) molecular biologist costs to interpret the results. Non-personnel costs were derived from the financial department of the NKI-AVL, and personnel costs from the collective labour agreement for Dutch hospitals (CAO) [1]. The purchasing costs for the MLPA kit were obtained from the MLPA website (SALSA MLPA P376 BRCA1ness probemix [2]) and the purchasing costs for the labeling kit for aCGH from ENZO lifesciences [3]. In the case of 53BP1, which is tested with immunochemistry, we derived the personal and testing costs from the Dutch Healthcare Authority's tariffs. The costs of running the tests were calculated with the most optimal test batch, being 18 samples for the MLPA and 12 for the aCGH. Direct non-medical and productivity costs of testing were assumed negligible.

Chemotherapy related costs

Medical direct costs of chemotherapy consisted of drug costs, day care costs and medical visit costs. We did not include the costs of radiotherapy because they were assumed equal under both regimens. The costs of chemotherapy were derived from and based on Dutch prices [4,5]. The costs associated to peripheral blood progenitor cell transplant (PBPC) procedures and subsequent follow up (in the HDAC arm) were derived from the Dutch Healthcare Authority's tariffs [6]. For both regimens we made two assumptions: (1) patients did not work during chemotherapy and (2) visits to the oncologist were scheduled during the chemotherapy days. Therefore, direct non-medical and productivity costs in the conventional regimen included the travelling costs on the days of chemotherapy and the 25 days missed at work. The direct non-medical and productivity costs in the HDAC regimen included one day of travelling costs for admission to the hospital, and productivity losses for 20 days of 4*FEC, 21 days hospitalized for 1*CTC/PBPC and 21 days post-transplant were the patient is controlled until recovery of blood activity. The costs associated with toxic deaths under the HDAC regimen were obtained from literature [7–9]

Health states costs

The costs of the health states disease free survival (DFS) and relapse (R) were based on Lidgren et al [10]. Costs of relapse were calculated as an average of local and distant relapse costs. The costs of death were excluded, unless it was consequence of treatment toxicity or breast cancer. In those situations we accounted for the specific costs to treat the toxicity (mentioned in the previous section) and for the palliative treatment.

-
- [1] VSNU. Collective labour agreement dutch universities, 1 September 2007 to 1 March 2010. The Hague: 2008.
 - [2] MRC-Holland. SALSA MLPA P376 BRCA1ness problemix 2013. <http://www.mlpa.com/WebForms/WebFormMain.aspx?Tag=fNPBLedDVp38p\CxU2h0mQ||> (accessed November 7, 2013).
 - [3] Enzo Life Sciences n.d. <http://www.enzolifesciences.com/contact-us/> (accessed September 23, 2014).
 - [4] Frederix GW. Disease specific methods for economic evaluations of breast cancer therapies. University of Utrecht, 2013.
 - [5] L. Hakkaart - van Roijen, S.S Tan, CAM Brouwmans. Guide for research costs - Methods and standard cost prices for economic evaluations in healthcare \ commissioned by the Health Care Insurance Board. Rotterdam: 2010.
 - [6] Dutch Healthcare Authority (NZA.nl). DBC product-finder for tariffs 2014. <http://www.nza.nl/organisatie/> (accessed February 27, 2014).
 - [7] Davies A, Ridley S, Hutton J, Chinn C, Barber B, Angus DC. Cost effectiveness of drotrecogin alfa (activated) for the treatment of severe sepsis in the United Kingdom. *Anaesthesia* 2005;60:155–62. doi:10.1111/j.1365-2044.2004.04068.x.
 - [8] Schilling MB. Costs and outcomes associated with hospitalized cancer patients with neutropenic complications: A retrospective study. *Exp Ther Med* 2011. doi:10.3892/etm.2011.312.
 - [9] Wang G, Zhang Z, Ayala C, Wall HK, Fang J. Costs of heart failure-related hospitalizations in patients aged 18 to 64 years. *Am J Manag Care* 2010;16:769–76.
 - [10] Lidgren M, Wilking N, Jönsson B, Rehnberg C. Resource use and costs associated with different states of breast cancer. *Int J Technol Assess Health Care* 2007;23:223–31. doi:10.1017/S0266462307070328.

PART III

IMAGING TECHNIQUES: MONITORING SYSTEMIC TREATMENT

CHAPTER 5

Imaging performance in guiding response to neoadjuvant therapy according to breast cancer subtypes: A systematic literature review

Melanie A Lindenberg

Anna Miquel-Cases

Valesca P Retèl

Gabe S Sonke

Marcel PM Stokkel

Jelle Wesseling

Wim H van Harten

Submitted for publication

Abstract

Background: Monitoring early therapeutic response to neoadjuvant chemotherapy (NAC) by imaging allows for an adaptive treatment approach likely to improve NAC effectiveness. As imaging performance seems to vary per tumor subtype, we aimed to generate a literature overview on subtype specific imaging performance in monitoring NAC in breast cancer (BC).

Methods: We performed a subtype specific literature search (BC classified by ER and HER2 status) to identify studies reporting on the performance of various imaging techniques in predicting pCR. Articles' quality was assessed by 1) sample size, 2) study design and 3) risk of bias assessed by the QUADAS tool. For each included study, negative and positive predictive value, (pooled) sensitivity and specificity, area under the curve values (AUC) and accuracy values were derived.

Results: Out of 106 identified articles, 15 were included. In *ER-positive/HER2-negative* BCs, ^{18}F -FDG-PET/CT showed a pooled sensitivity/specificity of 55%/89% and an AUC between 0.61–0.81, while MRI showed a pooled sensitivity/specificity of 35%/85% and an AUC of 0,55 (0,45–0,65). In *triple negative* BCs, ^{18}F -FDG-PET/CT showed a pooled sensitivity/specificity of 73%/96% and MRI showed a correlation with BRI ($p < 0.0001$, BRI represents relative change in tumor stage). In the overall *HER2-positive* group, ^{18}F -FDG-PET/CT showed a pooled sensitivity/specificity of 71%/69% and an AUC between 0.41–0.86, while MRI showed a correlation with BRI ($p = 0.05$). In *ER-positive/HER2-positive* and *ER-negative/HER2-positive* BCs, ^{18}F -FDG-PET/CT showed sensitivity/specificity of 59%/80% and 27%/88% respectively.

Conclusions: Our review reveals that evidence on the performance of imaging in monitoring NAC according to BC subtypes is lacking. Prior to starting well-designed studies that generate higher levels of evidence, consensus on specific study design characteristics should be reached (i.e., pCR definitions, imaging protocols or time intervals between baseline and response monitoring).

Introduction

In 2012, 1.7 million new cases of breast cancer were diagnosed worldwide. Breast cancer is still one of the most prevalent cancers overall, the most prevalent cancer among women and one of the main causes of death [1]. Research on new treatment approaches is thus of evident interest.

Neoadjuvant chemotherapy (NAC) is a treatment modality that consists on providing the systemic treatment prior to surgical removal of the tumor. NAC is equally effective as adjuvant chemotherapy [2] while having the additional advantage that therapeutic response can be monitored during treatment [3,4,5]. Early monitoring of therapeutic response by imaging seems to be a predictor of pathologic complete response (pCR) [6], a predictor of long-term survival in HER2-positive, triple negative (TN) and some ER-positive/HER2-negative tumours [8,9].

By monitoring therapeutic response during NACT, one can guide systemic treatment i.e. responders continue with the same initial treatment, and non-responders can be switched to a presumably non-cross-resistant regimen (Figure 1)[10]. This approach to administering NACT can be called response-guided NAC [10].

Currently, there is no definite guideline that describes how therapeutic response during NAC should be monitored. Previous authors have proposed the use of physical examination plus mammography and ultrasound, but their performance seems to be limited [11–13]. Therefore, performance examination of more advanced techniques, i.e. magnetic resonance imaging (MRI) and PET – Computed Tomography (PET/CT) seems warranted. So far, meta-analyses have shown sensitivities and specificities of 68%-91%, 93%-82% and 84%-71% for dynamic contrast-enhanced (DCE)-MRI [14], diffusion-weighted (DW)-MRI [14] and ¹⁸F-FDG-PET/CT [15] respectively. On the basis of these findings, MRI is currently the technique mainly used in clinical practice. Recent studies have now shown that breast cancer subtype affects imaging performance [16–18]. This means that some techniques may be more suitable for monitoring some subtypes than others. This also means that if these imaging technique- BC subtype combinations are identified, imaging performance can further improve [16,19]. So far there is no subtype-specific guidance on imaging techniques to monitor therapeutic response during NAC. This paper aims to create an overview of current knowledge on the performance of imaging techniques in monitoring NACT for four different breast cancer subtypes (based on ER and HER2 expression).

Figure 1: Response-guided neoadjuvant (NAC) approach. Patients start with first-line NAC treatment and after a specific number of cycles, they are monitored by imaging. Patients considered responders of NAC at imaging (according to a pre-defined threshold) continue the same initial treatment, whereas non-responders are switched to a presumably non-cross resistant treatment. Upon NAC finalization, pathologic response is determined at surgery, which is used to determine whether there the imaging results were correct.

Methods

We performed a systematic literature search to find studies reporting on the performance of imaging techniques in predicting pCR during NAC, separately per breast cancer subtype.

Search strategy

We searched in PubMed with the terms: “breast cancer” (MeSH: Breast neoplasm); “imaging” (i.e. MRI, PET/CT); “outcome” (pathologic complete response, clinical response); “Neoadjuvant chemotherapy” and “breast cancer subtype” (oestrogen receptor (ER), progesterone receptor (PR), luminal, triple negative (TN) and human endocrine receptor 2 (HER2) (see the systematic search in supplementary material 1). Snowballing was used to find additional relevant publications.

Selection criteria

The search was limited to studies written in English and published between January 2000 and March 2015. Case studies were excluded. Studies were included if monitoring was performed: 1) before and during NAC, 2) specific to at least one receptor status (ER/HER2) and 3) using pCR as ‘gold standard’ for response. Alternative outcomes to pCR were the neoadjuvant response index (NRI) [20] and residual cancer burden [21]. Finally, studies using FDG-PET without CT were excluded, as this technology is no longer recommended in daily practice.

Table 1: Categorization of different pathologic complete response definitions (pCR).

Category	Classifications and scales used in literature
Category 1 <u>Complete absence of invasive tumour cells and ductal carcinoma in situ (DCIS) in breast and axillary lymph nodes</u> after completion of neoadjuvant chemotherapy	- Chevalier classification grade 1 [48] - ypT0 ypN0
Category 2 <u>Complete absence of invasive tumour cells in the breast and axillary lymph nodes</u> after completion of neoadjuvant chemotherapy	- Chevalier classification grade 2 [48] - ypT0/is ypN0 - ypT0/is ypN0/+ - Miller and Payne grade 5 and NRG A or D [49]
Category 3 <u>Complete absence of invasive tumour cells in the breast</u> after completion of neoadjuvant chemotherapy	- Miller and Payne grade 5 [49] - YpT0/is
Category 4 <u>Considerable or partial reduction in tumour cells in breast</u> after completion of neoadjuvant chemotherapy	- Sataloff classification T-A [50] - Sataloff classification T-B [50] - Miller and Payne grade 4 [49]

Data extraction

First selection was performed based on abstract information and following the in- and exclusion criteria by two independent reviewers (AMC and ML). The selected studies were fully read by the same reviewers and were again assessed based on the in- and exclusion criteria. Disagreements were first discussed between the two reviewers, and if no agreement was reached, a third reviewer was approached (VR). For each article, the following items were extracted: author, sample size, study design, treatment regimen, breast cancer subtype, clinical stage, age, monitoring technique, cut-off value or response definition at imaging, interval time between baseline and response monitoring, technical settings of the imaging technique, pCR definition, performance results, i.e. sensitivity, specificity, accuracy, negative and positive predictive values (NPV, PPV), Area Under the Curve (AUC) in a Receiver Operating Curve (ROC), and if available, the number of false/true positives/negatives cases. pCR was categorized in the 4 definitions shown in table 1. Authors of articles where imaging performance was stratified to only one receptor status were contacted. They were asked for the existence and access to performance data stratified by the two receptors.

Quality assessment

Three research design criteria were defined to assess the quality of the included articles: 1) absence of treatment switch during NAC administration; 2) score higher than 8 on the Quality Assessment of Diagnostic Accuracy Studies (QUADAS)[22]; and 3) sample size ≥ 20 . Articles were considered of sufficient quality if they satisfied two of the three criteria. If more than one subtype was presented in the article, criteria 2 and 3 were assessed per subtype.

Performance of imaging

We constructed 2 x 2 contingency tables for articles directly reporting on the number of true/false negative/positive (TN,FN,TP,FP) patients and for articles where these numbers could be derived. These tables were used to calculate sensitivity (ability of imaging to identify non-responders with residual tumor tissue after NAC i.e. TP/TP+FN), specificity (ability of imaging to identify responders achieving a pCR after NAC i.e. TN/TN+FP), NPV (TN/TN+FN), PPV (TP/TP+FP) and accuracy (TP+TN/all patients). The pooled sensitivity and specificity of an imaging technique for a defined subtype was calculated to compare performances of different imaging techniques. This was only calculated if there was information from ≥2 articles using the same outcome measure. Calculations were performed by Review Manager 5[23] and Meta-DiSc[24]. If the inconsistency parameter (I^2) determined was ≥50% we considered there was substantial heterogeneity between articles, while if this was ≤30% we considered there was no significant heterogeneity [25].

Preferred imaging technique per subtype

We developed a scale to score and compare the performance of the various imaging techniques. The scale runs from A (perfect performance) to E (insufficient performance) and was applied to the various performance concepts i.e., ROC-AUC value, accuracy and sensitivity/specificity (table 3). The performance results per breast cancer subtype were placed in order, and, if sufficient results were available, the preferred imaging technique was chosen.

Table 2: Scale to score diagnostic performance. Each performance concept has its sensitivity and specificity data described as (α), ROC-AUC values were presented as (β) and accuracy results as (γ). The performance scales used per concept are presented in the last three columns of the table, and these are in turn categorized from perfect (A) to insufficient (E) performance by the first column of the table. General abbreviations: ROC-AUC: Area Under the - Receiver operator curve.

Performance	Sensitivity / specificity (α)	ROC-AUC value (β)	Accuracy (γ)
A Perfect	Both > 80%	0.90 – 1.00	90% - 100%
B Good	Both > 60% and < 80% or one result > 60% and < 80% and one result > 80%	0.80 – < 0.90	70 % - < 90%
C Sufficient	Both > 40% and < 60% or one result > 40% and < 60% and one result >60%	0.70 – < 0.80	50% - < 70%
D Limited	Both > 20% and < 40% or one result > 20% and < 40% and one result > 40%	0.60 – < 0.70	30% - < 50%
E Insufficient	Both < 20% or one result < 20% and one result > 20%	< 0.60	< 30%

Results

Of the initially 229 identified articles, 30 were selected for full reading after removing duplicates. 16 articles were further excluded based on our selection criteria. After snowballing one extra article was included, which made a total of 15 included articles (figure 2).

Figure 2: Flow diagram of the selection process. Of the 106 identified articles through PubMed, 15 articles were finally included.

Study characteristics

Sample sizes ranged from 7 to 246 patients (median: 31) and the overall mean age was 50. Studies enrolled patients prospectively (8 studies) and retrospectively (7 studies). One of the five contacted authors replied with additional data [26]. Nine articles presented results for the subgroup of ER-positive/HER2-negative patients [16,26–33], nine for the group of TN patients [16,19,27,28,30,32–35], nine for the whole group of HER2-positive patients [16,19,27,28,30,32,33,36,37] and one for the group of HER2-positive patients stratified by ER receptor status [38]. The NAC regimen differed per subtype: 1) ER-positive/HER2-negative patients received doxorubicin and cyclophosphamide (AC) plus docetaxel and capecitabine (DC) in case

of an unfavourable intermediate response [16,26–28], 2) TNBC patients received epirubicin and cyclophosphamide followed by Docetaxel (EC-D)[29,33–36] or one of the following regimens: intensified EC-D (SIM) [34,35], fluorouracil plus EC (FEC) [19,39] and FEC-D [19,39], and 3) ER-negative/HER2-positive patients received EC(-D) followed by a combination of trastuzumab and paclitaxel or Docetaxel [33,37,38]. Of the included articles, 3 were on MRI and 12 on ^{18}F -FDG-PET/CT (a summary of the main technical settings used in response assessment are presented in table 3). Regarding the quality of the studies, 3 of the assessed subtypes showed a small sample size [30,32,33], 4 had a study design that allowed a switch in treatment during NAC [16,26–28], but no study showed a score below 8 on the QUADAS list (supplementary material 3). Since each subgroup of each article satisfied 2 of the 3 criteria, no study or subgroup was excluded from further analysis (table 4).

All collected study characteristics are presented in supplementary material 2.

Table 3: Main technical settings of imaging techniques used in response assessment summarized per imaging technique. More details are described in the study characteristics table (supplement 2). General abbreviations: MBq MegaBecquerel; mAs: milliampere /second; kV: Kilovolt; T: Tesla.

Imaging technique	Technology	Contrast (dosage)	Settings	Position
MRI [16,26,31]	Philips magnetom vision [16,26] 1.5T and 3.0T magnet [16,26,31]	Gadolinium (14ml of 0.1mmol/kg)[16,26]	-	Use of breast coils [16,26,31]
^{18}F-FDG-PET/CT [19,27–30,32–38]	Philips [19,27–29,33–36,38,42] GE medical [30,32,38] Siemens [38]	^{18}F -FDG (3.5 MBq/kg – 7.4 MBq/kg)[19,27–30,32–38] Fasted 6 hours before injection [19,27–30,32–38]	Scan performed 60 to 70 min after contrast injection CT: 120kV and 100mAs [19,27–30,32–38]	Hanging breast method [27,28]

Table 4: Quality assessment based on three criteria: 1. The treatment was not switched during NAC, 2. Study does not score below 8 on the quality assessment tool for diagnostic accuracy studies (QUADAS), 3. The sample size is above 20 patients.

Author (year) reference	Subtype	Sample size	Criteria 1 Treatment is not switched during NAC	Criteria 2 No risk of bias is present	Criteria 3 Sample size is ≥ 20 patients	Include?
Charehbil (2014) [31]	ER-positive/HER2- negative	194	+	+	+	Yes
Gebhart (2013) [38]	ER-negative/HER2- positive	43	+	+	+	Yes
	ER-positive/HER2- positive	34	+	+	+	Yes
Groheux (2012) [34]	TN	20	+	+	+	Yes
Groheux (2013) [29]	ER-positive/HER2- negative	64	+	+	+	Yes
Groheux (2013) [36]	HER2-positive	30	+		+	Yes
Groheux (2014) [35]	TN	50	+	+	+	Yes
Hatt (2013)[33]	ER-positive/HER2- negative	26	+	+	+	Yes
	TN	13	+	+	-	Yes
	HER2-positive	12	+	+	-	Yes
Humbert (2012) [19]	ER-positive/HER2- negative	53	+	+	+	Yes
	TN	25	+	+	+	Yes
	HER2-positive	37	+	+	+	Yes
Humbert (2014) [37]	HER2-positive	57	+	+	+	Yes
Koolen (2014)[27]	ER-positive/HER2- negative	50	-	+	+	Yes
	TN	31	+	+	+	Yes
	HER2-positive	26	+	+	+	Yes
Koolen (2013)[28]	ER-positive/HER2- negative	45	-	+	+	Yes
	TN	25	+	+	+	Yes
	HER2-positive	25	+	+	+	Yes
Loo (2011)[16]	ER-positive/HER2- negative	103	-	+	+	Yes
	TN	47	+	+	+	Yes
	HER2-positive	38	+	+	+	Yes
Martoni (2010)[32]	ER-positive/HER2- negative	16	+	+	-	Yes
	TN	9	+	+	-	Yes
	HER2-positive	7	+	+	-	Yes
Rigter (2013)[26]	ER-positive/HER2- negative	246	-	+	+	Yes
Zucchini (2013) [30]	ER-positive/HER2- negative	31	+	+	+	Yes
	TN	15	+	+	-	Yes
	HER2-positive	14	+	+	-	Yes

Performance of imaging techniques per subtype

Results on the performance of the various imaging techniques per breast cancer subtype are summarized in the section below and in table 5. For each study we determined the number of NAC cycles between baseline and response monitoring, the cut-off value of response and the pCR definition used.

ER-positive/HER2-negative

Six studies assessed the performance of ^{18}F -FDG-PET/CT. The use of ^{18}F -FDG-PET/CT showed AUC-ROC values of 0.61 (CI 0.37-0.86; after 1 NAC cycle; pCR category 2)[27], 0.87 (CI 0.69–1.00; after 3 NAC cycles; pCR category 2)[27], 0.77 (CI 0.68–0.87; after 3 NAC cycles; pCR category 3)[28] and 0.88 (after 2 NAC cycles; pCR category 4) in one study [33]. An Italian research group described the performance of ^{18}F -FDG-PET/CT in 2 articles. Both studies showed a sensitivity of 38% and specificity of 100% (cut-off value $\geq$ -50% Standardize Uptake Value ($\Delta\text{SUV max}$); after 2 NAC cycles; pCR category 4)[30,32]. Another study showed ^{18}F -FDG-PET/CT sensitivity of 62% and specificity of 78% (cut-off value $\geq$ -38% $\Delta\text{SUV max}$; after 2 NAC cycles; pCR category 4)[29]. When using the difference in Total Lesion Glycolysis (ΔTLG) as outcome measure at imaging, ^{18}F -FDG-PET/CT showed a sensitivity of 89% and sensitivity of 74%, and AUC values of 0.81 (cut-off value $\geq$ -71% ΔTLG ; after 2 NAC cycles; pCR category 4)[29] and 0.96 (after 2 NAC cycles; pCR definition 4)[33].

Three studies assessed the performance of MRI. One trial showed sensitivity of 35%, specificity of 89%, accuracy of 39%, NPV of 10% and PPV of 98% (cut-off value $\geq$ -25%; after 3 NAC cycles; pCR category 3)[26] and sensitivity of 37%, specificity of 87%, accuracy of 45%, NPV of 22% and PPV of 93% (cut-off value $\geq$ -30%; after 3 NAC cycles; pCR category 3)[31]. Although this trial results were reported for HER2-negative patients, as the majority of patients were ER-positive (187/222) we included them in this subtype [31]. One MRI study did not report specific performance results, but showed no significant association between tumour size decrease and Breast Response Index (BRI; part of the NRI outcome measure [20])($p=0.07$; after 3 NAC cycles; pCR definition 4)[16].

Triple negative

Eight studies assessed the performance of ^{18}F -FDG-PET/CT. The use of ^{18}F -FDG-PET/CT showed AUC values of 0.76 (CI 0.55-0.96; after 1 NAC cycle; pCR category 2)[27], 0.87 (CI 0.73-1.00; after 3 NAC cycles; pCR category 2)[27] and 0.85 (CI 0.68–1.00; after 3 NAC cycles; pCR category 3)[28]. The performance of ^{18}F -FDG-PET/CT was described in another 2 articles with sensitivity

of 71% and 79%, specificity of 95% and 100%, and accuracy of 80% and 85% (cut-off value $\geq$ -50% Δ SUV max; after 2 NAC cycles; pCR category 2)[34,35]. These studies showed that by lowering the Δ SUV max cut-off value to $\geq$ -42% specificity improved to 100%, but sensitivity decreased to 58% and 64% respectively [34,35]. Two additional studies of the same research group showed sensitivity of 0% and specificity of 100% (cut-off value $\geq$ -50% Δ SUV max; after 2 NAC cycles; pCR category 4) as in these studies neither true nor false non-responders were identified [30,32]. Furthermore, one study showed no significant association between Δ SUV and pCR ($p=0.50$ after 1 NAC cycle)[19], and another study showed no significant improvement in predictive value ($p>0.05$) by using Δ TLG as outcome measure[33].

One study assessed the performance of MRI. This study reported a significant association between tumour size decrease and BRI ($p < 0.001$)[16].

HER2-positive

Eight studies assessed the performance of ^{18}F -FDG-PET/CT. The use of ^{18}F -FDG-PET/CT showed AUC values of 0.61 (CI 0.33-0.89; after 3 NAC cycles; pCR category 2)[27], 0.59 (CI 0.34-0.85; after 8 NAC cycles; pCR category 2)[27] and 0.41 (CI 0.16-0.67; after 8 NAC cycles; pCR category 3)[28]. Two studies showed sensitivity of 17% and 20%, specificity of 100% [30,32], and accuracy of 29% [32](cut-off value $\geq$ -50% Δ SUV max; after 2 NAC cycles; pCR category 4). Three other studies showed sensitivities and specificities of ^{18}F -FDG-PET/C, 86% and 75% (cut-off value $\geq$ -62% Δ SUV max; after 2 NAC cycles; pCR category 2)[36], 86% and 63% (cut-off value $\geq$ -62% Δ SUV max; after 2 NAC cycles; pCR category 3)[36], 83% and 53% (cut-off value $\geq$ -60% Δ SUV max; after 1 NAC cycle; pCR category 2)[37] and 64%, 83% and accuracy of 76% (cut-off value $\geq$ -75%; after 1 NAC cycle; pCR category 2)[19]. In this subtype, using Δ TLG instead of Δ SUV max showed no improvement in predictive value [33].

One study assessed the performance of MRI. This study reported a significant association between response at imaging and BRI ($p=0.05$; after 8 cycles NAC)[16].

ER-positive/HER2-positive

One study assessed the performance of ^{18}F -FDG-PET/CT. This showed a sensitivity of 38%, specificity of 71%, accuracy of 44%, NPV of 20% and PPV of 86% (cut-off value $\geq$ -15% Δ SUV max; after 2 weeks; pCR category 3) and sensitivity of 59%, specificity of 80%, accuracy of 62%, NPV of 24% and PPV of 95%(cut-off value $\geq$ -25% Δ SUV max; after 6 weeks; pCR category 3) [38].

Table 5: Performance of imaging techniques per subtype. Response definition: I response category 1; II response category 2; III response category 3; IIII response category 4; Cut-off values: 1: cut-off value 25% size reduction; 2: cut-off value 30% size reduction; 3: cut-off value -15% Δ SUVmax; 4: cut-off value -25% Δ SUVmax; 5: cut-off value -38% Δ SUVmax; 6: cut-off value -42% Δ SUVmax; 7: cut-off value -50% Δ SUVmax; 8: cut-off value -60% Δ SUVmax; 9: cut-off value -62% Δ SUVmax; 10: cut-off value -75% Δ SUVmax; 11: cut-off value -71% Δ TLG; Outcome parameters: *: Δ SUVmax; Δ : Different outcome parameters; Performance: α : Sensitivity and specificity results; β : AUC values; γ : Accuracy values; Other: #=in the original article it was described as administrations instead of cycles; General abbreviations: AUC = Area Under the Curve; NPV: Negative Predictive Value; PPV: Positive Predictive Value; SUV: Standard Uptake Value; TLG: Total Lesion Glycolysis; MATV: Metabolic Active Tumour Value.

Article (reference) (PCR category)	Monitoring technique (cut-off value or outcome parameter)	Performance score (A – E) (type result)	sensitivity, specificity, accuracy, NPV, PPV	AUC	Monitoring interval
ER-positive/HER2-negative					
Hatt [33] ^(III)	¹⁸ F-FDG-PET/CT ^(Δ)	B ^(β) , A ^(β) , B ^(β)	-	Δ SUVmax: 0.88 Δ TLG: 0.96 Δ MATV: 0.98	After 2 cycles
Groheux [29] ^(III)	¹⁸ F-FDG-PET/CT ⁽¹¹⁾	B ^{(α)B^(β)}	89%, 74%, -, 31%, 98%	0.81	After 2 cycles
Koolen [27] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	B ^(β)	-	0.87 (0.69-1.00)	After 3 cycles
Groheux [29] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁵⁾	B ^{(α)C^(β)}	62%, 78%, -, 12%, 97%	0.73	After 2 cycles
Koolen [28] ^(III)	¹⁸ F-FDG-PET/CT ^(*)	C ^(β)	-	0.77 (0.68 – 0.87)	After 3 cycles
Koolen [27] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	D ^(β)	-	0.61 (0.37-0.86)	After 1 cycle
Zucchini [30] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	D ^(α)	38%, 100%, -, 24%, 100%	-	After 2 cycles
Martoni [32] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	D ^{(α)C^(γ)}	38%, 100%, 50%, 27%, 100%	-	After 2 cycles
Rigter [26] ^(III)	DCE MRI ⁽¹⁾	D ^{(α)D^(γ)}	35%, 89%, 39%, 10%, 98%	-	After 3 cycles
Charehbil [31] ^(III)	DCE MRI ⁽²⁾	D ^{(α)E^(β)} D ^(γ)	37%, 87%, 45%, 22%, 93%	0.55 (0.45 – 0.65)	After 3 cycles
Loo [16] ^(III)	DCE MRI ⁽²⁾	-	Association between BRI and tumor decrease was not significant (p = 0.07)	-	After 3 cycles
Triple negative					
Groheux [35] ^(II)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	B ^{(α)B^(γ)}	71%, 95%, 80%, 67%, 96%	-	After 2 cycles
Groheux [34] ^(II)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	B ^{(α)B^{(β)B^(γ)}}	79%, 100%, 85%, 67%, 100%	0.881	After 2 cycles
Groheux [34] ^(II)	¹⁸ F-FDG-PET/CT ⁽⁶⁾	B ^{(α)B^{(β)B^(γ)}}	64%, 100%, 75%, 55%, 100%	0.881	After 2 cycles
Koolen [28] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	B ^(β)	-	0.85 (0.69-1.00)	After 3 cycles
Koolen [27] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	B ^(β)	-	0.87 (0.73-1.00)	After 3 cycles
Groheux [35] ^(II)	¹⁸ F-FDG-PET/CT ⁽⁶⁾	C ^{(α)B^(γ)}	58%, 100%, 74%, 59%, 100%	-	After 2 cycles
Koolen [27] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	C ^(β)	-	0.76 (0.55-0.96)	After 1 cycle

Zucchini [30] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	E ^(α)	0%, 100%, -, 27%, 0%	-	After 2 cycles
Martoni [32] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	E ^{(α)D^(γ)}	0%, 100%, 33%, 33%, -	-	After 2 cycles
Humbert [19] ^(II)	¹⁸ F-FDG-PET/CT ⁽¹⁰⁾	-	No significant correlation between early metabolic response and pCR		After 1 cycle
Hatt [33] ^(III)	¹⁸ F-FDG-PET/CT ^(Δ)	-	Use of different parameters did not improve predictive value of ΔSUVmax		After 2 cycles
Loo et al [16] ^(III)	DCE MRI ⁽²⁾	-	Association between BRI and largest tumor diameter was significant (p= < 0.001)		After 3 cycles
HER2-positive					
Groheux [36] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁹⁾	B ^{(α)B^{(β)B^(γ)}}	86%, 63%, 73%, 84%, 67%	0.86	After 2 cycles
Groheux [36] ^(II)	¹⁸ F-FDG-PET/CT ⁽⁹⁾	B ^{(α)B^{(β)B^(γ)}}	86%, 75%, 80%, 86%, 75%	0.86	After 2 cycles
Humbert [19] ^(II)	¹⁸ F-FDG-PET/CT ⁽¹⁰⁾	B ^{(α)C^{(β)B^(γ)}}	64%, 83%, 76%, 79%, 69%	0.73	After 1 cycle
Humbert [37] ^(II)	¹⁸ F-FDG-PET/CT ⁽⁸⁾	C ^{(α)C^{(β)B^(γ)}}	83%, 52%, -, 84%, 50%	0.70 (0.55-0.85)	After 1 cycle
Koolen [27] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	D ^(β)	-	0.61 (0.33-0.89)	After 3 cycles [#]
Zucchini [30] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	D ^(α)	20%, 100%, -, 33%, 100%	-	After 2 cycles
Koolen [27] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	E ^(β)	-	0.59 (0.34-0.85)	After 8 cycles [#]
Koolen [28] ^(II)	¹⁸ F-FDG-PET/CT ^(*)	E ^(β)	-	0.41 (0.16-0.67)	After 8 cycles [#]
Martoni [32] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁷⁾	E ^{(α)E^(γ)}	17%, 100%, 29%, 17%, 100%	-	After 2 cycles
Hatt [33] ^(III)	¹⁸ F-FDG-PET/CT ^(Δ)	-	Use of different parameters did not improve predictive value of ΔSUVmax		After 2 cycles
Loo [16] ^(III)	DCE MRI ⁽²⁾	-	Association between BRI and largest tumor diameter was significant (p= 0.05)		After 8 cycles [#]
HER2-positive and ER-positive					
Gebhart [38] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁴⁾	C ^{(α)C^(γ)}	59%, 80%, 62%, 24%, 95%	-	After 6 weeks
Gebhart [38] ^(III)	¹⁸ F-FDG-PET/CT ⁽³⁾	D ^{(α)D^(γ)}	38%, 71%, 44%, 20%, 86%	-	After 2 weeks
HER2-positive and ER-negative					
Gebhart [38] ^(III)	¹⁸ F-FDG-PET/CT ⁽³⁾	D ^{(α)C^(γ)}	27%, 88%, 64%, 65%, 60%	-	After 2 weeks
Gebhart [38] ^(III)	¹⁸ F-FDG-PET/CT ⁽⁴⁾	E ^{(α)C^(γ)}	18%, 76%, 54%, 59%, 33%	-	After 6 weeks

ER-negative/HER2-positive

One study assessed the performance of ^{18}F -FDG-PET/CT. This showed a sensitivity of 27%, specificity of 88%, accuracy of 64%, NPV of 65% and PPV of 60% (cut-off value $\geq -15\%$ ΔSUV max; after 2 weeks; pCR category 3) and sensitivity of 18%, specificity of 76%, accuracy of 54%, NPV of 59% and PPV of 33% (cut-off value $\geq -25\%$ ΔSUV max; after 6 weeks; pCR category 3) [38].

Pooled performance of imaging

In ER-positive/HER2-negative patients the pooled sensitivity and specificity of ^{18}F -FDG-PET/CT was 55% (95% CI 0.44–0.65) and 89% (95% CI 0.52–1.00)[29,30] and for MRI it was 35% (95% CI 0.31 – 0.41) and 85% (95% CI 0.73–0.93)[26,31]. Two articles initially included in this pooled analysis used the same database, we thus only included the most recent results [30,32]. For TNBCs we constructed two pooled analyses for ^{18}F -FDG-PET/CT, one for $\geq -50\%$ ΔSUV max resulting in sensitivity and specificity of 73% (95% CI 0.58–0.85) and 96% (95% CI 0.80–1.00), and one for $\geq -42\%$ ΔSUV max resulting in sensitivity and specificity of 60% (95% CI 0.44–0.74) and 100% (95% CI 0.86–1.00) [34,35]. In the overall HER2-positive group, the pooled sensitivity and specificity of ^{18}F -FDG-PET/CT were 71% (95% CI 0.60–0.81) and 69% (95% CI 0.56–0.81)) [19,30,36,37]. Heterogeneity was present in the pooled sensitivity of ^{18}F -FDG-PET/CT in the ER-positive/HER2-negative and the HER2-positive groups (supplementary material 4).

Preferred imaging technique per subtype

Due to the limited number of studies reporting on the performance of imaging per subtype, we could not conclude on subtype preferred imaging techniques.

Discussion

In view of the potential of response-guided NAC to improve breast cancer survival, we aimed to generate a literature overview on subtype specific imaging performance in monitoring NAC in breast cancer (BC).

Our results suggest that due to the differences in imaging performance across subtypes, personalizing the monitoring step of response-guided NAC based on these is of relevance. However, after reviewing the 15 included articles, we revealed that there is lack of evidence with enough statistical power to conclude on the preferred imaging technique per subtype. Although

we did identify studies reporting on the performance of MRI and ^{18}F -FDG-PET/CT specified to breast cancer subtypes, all studies were observational and showed a lot of inter study variability. Thereby, our results should be seen as preliminary and thus be interpreted with caution. This information can nonetheless serve to pinpoint areas of further research.

In the ER-positive/HER2-negative subtype, the best performing technique was ^{18}F -FDG-PET/CT after 2 NAC cycles [29], while the use of MRI was limited. Furthermore, we saw that in this subtype the performance of ^{18}F -FDG-PET/CT improved when using the measures ΔTLG and Metabolic Active Tumour Volume instead of the standard $\Delta\text{SUV max}$ [29,33]. In TNBCs, ^{18}F -FDG-PET/CT showed also a good performance [27,28,34,35], with the best results seen after 2 NAC cycles using a cut-off value of $\geq 50\%$ $\Delta\text{SUV max}$ (performance: $(\alpha)\text{B}(\gamma)$) [35]. The use of MRI seems also promising in this subtype, as size decrease showed a correlation with BRI [16]. In the overall HER2-positive group, ^{18}F -FDG-PET/CT showed promising results [19,27,36,37], especially after 2 NAC cycles using a cut-off value $\geq 62\%$ ΔSUVmax (performance: $\text{B}(\alpha)\text{B}(\beta)\text{B}(\gamma)$) [36]. However, when these patients were split by ER status performance was limited [38]. We hypothesize that this may be consequence of the use of a lower cut-off value at imaging and a different monitoring interval vs. other ^{18}F -FDG-PET/CT studies. In the overall HER2-positive group, MRI showed an association between tumour size decrease and BRI [16]. Our study results thus suggest that further investigations on the performance of MRI in TNBC and HER2-positive breast cancer are relevant.

Previous publications that described and reviewed literature on subtype specific imaging performance in monitoring NAC are in line with our findings. For instance, Lobbes and colleagues showed that MRI was more accurate in HER2-positive tumours than in HER2-negative tumours [40]. Humbert et al. and Groheux et al. showed good performance of ^{18}F -FDG-PET/CT in HER2-positive breast cancer patients when using the difference in SUV uptake as measure [41,42]. ^{18}F -FDG-PET/CT showed promising performance results also in TNBC by both $\Delta\text{SUV max}$ and ΔTLG measurement (AUC values of 0.86 and 0.88 respectively [41] and overall accuracy of 75% [43]). The potential of ΔTLG as an outcome for ^{18}F -FDG-PET/CT was confirmed by other research groups, whom showed its correlation with survival [41,44]. In addition, the use of absolute values of SUVmax and SUV peak instead of their difference was also suggested for their better performance in predicting pCR [41]. Furthermore, FES-PET/CT, and DWI-MRI seem to be promising techniques; FES-PET/CT seems useful in ER-positive tumours [45] and DWI-MRI seems to be complementary to DCE-MRI [46]. Both techniques are currently being investigated in trials (NCT02398773; NCT01564368).

We identified two studies testing the effectiveness of the response-guided NAC approach. The first study was a RCT for ER-negative/HER2-positive patients in which patients were scanned by

^{18}F -FDG-PET after 1 NAC cycle, and Bevacizumab was randomly assigned to non-responders ($\leq -70\%$ $\Delta\text{SUV max}$) in a 2:1 ratio [47]. Unfortunately, response assessment in this study was based on PET alone and had to be excluded from our review. The second study was a non-randomized non-controlled prospective study for ER-positive/HER2-negative patients in which patients were scanned by MRI and in case of no response patients were switched from AC to DC. Patients that received AC and DC showed improved tumour size reduction [26]. The NPV of MRI in this study was 10%, meaning that only 10% of non-responders were correctly identified (assuming that 1) the switch to non-cross resistant would be beneficial, 2) pCR would correlate to survival in this subtype, and 3) the optimal way to predict therapeutic response had been chosen). Under these assumptions, the use of ^{18}F -FDG-PET/CT would increase the NPV to 31% (according to our results). These scenarios illustrate that improved effectiveness of the response-guided NAC approach can be achieved with improved imaging performance, more effective treatments or the combination of both.

5

This review included few studies, mainly underpowered, and of heterogeneous study designs and outcome measures. Variability mainly occurred due to 1) differences in interval time between imaging at baseline and monitoring, 2) cut-off values to define treatment response, and 3) pCR definitions. These variations are consequence of the lack of consensus on imaging settings and protocols. As we were aware of these and of its possible influence on results, we carefully described study differences in our results section. The inter-variability and the limited number of studies included in the review also limited the possibility of pooling. Another issue was the higher frequency of ^{18}F -FDG-PET/CT vs. MRI studies. This is consequence of many of the initially identified MRI studies combining performance results of response assessment during and after NAC in the same analysis. The lack of results on MRI in the majority of the subtypes made it impossible to compare its performance to ^{18}F -FDG-PET/CT and consequently to conclude on the preferred imaging technique per subtype. A last discussion point is the inclusion of studies only describing performance results according to one receptor status, as it is known that performance could be affected by the other unknown receptor status. Besides, in the ER-positive/HER2-negative group we did not differentiate into luminal A and B tumours, despite knowing that in luminal A tumours pCR does not correlate with survival [9]. Therefore, our conclusions for this subtype may be unlikely. Nonetheless, they serve to illustrate the urgency to reach consensus for a reliable alternative for pCR in this subgroup.

The major limitation of this study, which is the inclusion of few and insufficiently studies, has been also the guide to find what is needed to decide on the most effective imaging technique per subtype, which is consensus on several aspects that affect study comparability. Specifically, on 1) the definition of pathologic response, 2) the thresholds to define complete-, near-, partial-, or no- response during NAC in both ^{18}F -FDG-PET/CT and MRI, 3) the required interval time between

baseline and response monitoring per subtype and imaging technique, and 4) the imaging settings. Only then, meaningful well-designed studies which account for various breast cancer subtypes and imaging techniques can be conducted. Whereupon, RCTs such as the AVATACXER trial [47] which mimics the response-guided NAC approach, could be initiated. This type of trials will also inform on suitable treatment switches per subtype. Further, we suggest conducting further research to: 1) less investigated techniques such as FES-FDG/PET and DWI-MRI, 2) potential predictive biomarkers that could further personalize the response-guided NAC approach i.e. Ki67 and P53 and 3) the association between NAC treatments and imaging performance. Finally, a cost-effectiveness analysis could be interesting to explore the health-economic consequences of various scenarios of this response-guided NAC approach.

This literature review is unique in the way that it focuses on imaging performance of NAC monitoring specified to breast cancer subtypes. We conclude that the level of evidence of current studies is too low to be able to draw reliable subtype-specific imaging recommendations, and that these can only occur when consensus on imaging settings and work regulations are reached. Further research on these are necessary to eventually build protocols and use them to conceive comparable study outcomes.

References

1. WHO. Breast Cancer Estimated Incidence, Mortality and Prevalence Worldwide in 2012 [Internet]. [cited 2015 Nov 11]. Available from: http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx
2. Mauri D, Pavlidis N, Ioannidis JPA. Neoadjuvant versus adjuvant systemic treatment in breast cancer: A meta-analysis. *J Natl Cancer Inst.* 2005;97.
3. Fisher B, Brown A, Mamounas E, Wieand S, Robidoux A, Margolese RG, et al. Effect of preoperative chemotherapy on local-regional disease in women with operable breast cancer: findings from National Surgical Adjuvant Breast and Bowel Project B-18. *J Clin Oncol.* 1997;15.
4. van der Hage JA, van de Velde CJ, Julien JP, Tubiana-Hulin M, Vandervelden C, Duchateau L. Preoperative chemotherapy in primary operable breast cancer: results from the European Organization for Research and Treatment of Cancer trial 10902. *J Clin Oncol.* 2001;19.
5. Gralow JR, Burstein HJ, Wood W, Hortobagyi GN, Gianni L, Von Minckwitz G, et al. Preoperative therapy in invasive breast cancer: Pathologic assessment and systemic therapy issues in operable disease. *J Clin Oncol.* 2008;26.
6. Marinovich ML, Macaskill P, Irwig L, Sardanelli F, von Minckwitz G, Mamounas E, et al. Meta-analysis of agreement between MRI and pathologic breast tumour size after neoadjuvant chemotherapy. *Br J Cancer* [Internet]. 2013;109. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23963140>
7. Kaufmann M, Hortobagyi GN, Goldhirsch A, Scholl S, Makris A, Valagussa P, et al. Recommendations from an international expert panel on the use of neoadjuvant (primary) systemic treatment of operable breast cancer: an update. *J Clin Oncol.* 2006;24(12).
8. Chollet P, Amat S, Cure H, de Latour M, Le Bouedec G, Mouret-Reynier M-A, et al. Prognostic significance of a complete pathological response after induction chemotherapy in operable breast cancer. *Br J Cancer.* 2002;86(7).
9. Von Minckwitz G, Untch M, Blohmer JU, Costa SD, Eidtmann H, Fasching PA, et al. Definition and impact of pathologic complete response on prognosis after neoadjuvant chemotherapy in various intrinsic breast cancer subtypes. *J Clin Oncol.* 2012;30(15).
10. von Minckwitz G, Blohmer JU, Costa SD, Denkert C, Eidtmann H, Eiermann W, et al. Response-guided neoadjuvant chemotherapy for breast cancer. *J Clin Oncol.* 2013;31.
11. Yeh E, Slanetz P, Kopans DB, Rafferty E, Georgian-Smith D, Moy L, et al. Prospective comparison of mammography, sonography, and MRI in patients undergoing neoadjuvant chemotherapy for palpable breast cancer. *Am J Roentgenol.* 2005;184(3).
12. Hamisa M, Dabess N, Yosef R, Zakeria F, Hammed Q. Role of breast ultrasound, mammography, magnetic resonance imaging and diffusion weighted imaging in predicting pathologic response of breast cancer after neoadjuvant chemotherapy. *Egypt J Radiol Nucl Med.* 2015;46(1).
13. Londero V, Bazzocchi M, Frate C Del, Puglisi F, Loreto C Di, Francescutti G, et al. Locally advanced breast cancer: comparison of mammography, sonography and MR imaging in evaluation of residual disease in women receiving neoadjuvant chemotherapy. *Eur Radiol.* 2004;14(8).
14. Wu L-M, Hu J-N, Gu H-Y, Hua J, Chen J, Xu J-R. Can diffusion-weighted MR imaging and contrast-enhanced MR imaging precisely evaluate and predict pathological response to neoadjuvant chemotherapy in patients with breast cancer? *Breast Cancer Res Treat.* 2012;135(1).
15. Cheng X, Li Y, Liu B, Xu Z, Bao L, Wang J. 18F-FDG PET/CT and PET for evaluation of pathological response to neoadjuvant chemotherapy in breast cancer: a meta-analysis. *Acta Radiol.* 2012;53(6).
16. Loo CE, Straver ME, Rodenhuis S, Muller SH, Wesseling J, Vrancken Peeters MJTFD, et al. Magnetic resonance imaging response monitoring of breast cancer during neoadjuvant chemotherapy: Relevance of Breast Cancer Subtype. *J Clin Oncol.* 2011;29.
17. Hayashi Y, Takei H, Nozu S, Tochigi Y, Ichikawa A, Kobayashi N, et al. Analysis of complete response by MRI following neoadjuvant chemotherapy predicts pathological tumor responses differently for molecular subtypes of breast cancer. *Oncol Lett.* 2013 Jan;5(1).

18. Ko ES, Han B-K, Kim RB, Ko EY, Shin JH, Hahn SY, et al. Analysis of factors that influence the accuracy of magnetic resonance imaging for predicting response after neoadjuvant chemotherapy in locally advanced breast cancer. *Ann Surg Oncol*. United States; 2013 Aug;20(8).
19. Humbert O, Berriolo-Riedinger A, Riedinger JM, Coudert B, Arnould L, Cochet A, et al. Changes in 18F-FDG tumor metabolism after a first course of neoadjuvant chemotherapy in breast cancer: influence of tumor subtypes. *Ann Oncol*. England; 2012 Oct;23(10).
20. Rodenhuis S, Mandjes IAM, Wesseling J, van de Vijver MJ, Peeters M-JTFV, Sonke GS, et al. A simple system for grading the response of breast cancer to neoadjuvant chemotherapy. *Ann Oncol*. 2010;21(3).
21. Symmans WF, Peintinger F, Hatzis C, Rajan R, Kuerer H, Valero V, et al. Measurement of residual breast cancer burden to predict survival after neoadjuvant chemotherapy. *J Clin Oncol*. 2007;25(28).
22. Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM, et al. Towards complete and accurate reporting of studies of diagnostic accuracy: The STARD initiative. *Vet Clin Pathol*. 2007;36.
23. The Cochrane Collaboration. Review Manager (RevMan). Copenhagen: The Nordic Cochrane Centre; 2012.
24. Zamora J, Abraira V, Muriel A, Khan K, Coomarasamy A. Meta-DiSc: a software for meta-analysis of test accuracy data. *BMC Med Res Methodol*. 2006;6.
25. The Cochrane Collaboration. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. 2011. Available from: www.cochrane-handbook.org
26. Rigter LS, Loo CE, Linn SC, Sonke GS, van Werkhoven E, Lips EH, et al. Neoadjuvant chemotherapy adaptation and serial MRI response monitoring in ER-positive HER2-negative breast cancer. *Br J Cancer*. England; 2013 Dec;109(12).
27. Koolen BB, Pengel KE, Wesseling J, Vogel W V, Vrancken Peeters M-JTFD, Vincent AD, et al. Sequential (18)F-FDG PET/CT for early prediction of complete pathological response in breast and axilla during neoadjuvant chemotherapy. *Eur J Nucl Med Mol Imaging*. Germany; 2014 Jan;41(1).
28. Koolen BB, Pengel KE, Wesseling J, Vogel W V, Vrancken Peeters M-JTFD, Vincent AD, et al. FDG PET/CT during neoadjuvant chemotherapy may predict response in ER-positive/HER2-negative and triple negative, but not in HER2-positive breast cancer. *Breast*. 2013;22(5).
29. Groheux D, Hatt M, Hindié E, Giacchetti S, De Cremoux P, Lehmann-Che J, et al. Estrogen receptor-positive/human epidermal growth factor receptor 2-negative breast tumors: Early prediction of chemosensitivity with 18F-fluorodeoxyglucose positron emission tomography/computed tomography during neoadjuvant chemotherapy. *Cancer*. 2013;119(11).
30. Zucchini G, Quercia S, Zamagni C, Santini D, Taffurelli M, Fanti S, et al. Potential utility of early metabolic response by 18F-2-fluoro-2-deoxy-D-glucose-positron emission tomography/computed tomography in a selected group of breast cancer patients receiving preoperative chemotherapy. *Eur J Cancer*. 2013;49(7).
31. Charehbili A, Wasser MN, Smit VTHBM, Putter H, van Leeuwen-Stok AE, Meershoek-Klein Kranenbarg WM, et al. Accuracy of MRI for treatment response assessment after taxane- and anthracycline-based neoadjuvant chemotherapy in HER2-negative breast cancer. *Eur J Surg Oncol*. England; 2014 Oct;40(10).
32. Martoni AA, Zamagni C, Quercia S, Rosati M, Cacciari N, Bernardi A, et al. Early (18)F-2-fluoro-2-deoxy-d-glucose positron emission tomography may identify a subset of patients with estrogen receptor-positive breast cancer who will not respond optimally to preoperative chemotherapy. *Cancer*. United States; 2010 Feb;116(4).
33. Hatt M, Groheux D, Martineau A, Espié M, Hindié E, Giacchetti S, et al. Comparison between 18F-FDG PET image-derived indices for early prediction of response to neoadjuvant chemotherapy in breast cancer. *J Nucl Med*. 2013;54(3).
34. Groheux D, Hindié E, Giacchetti S, Delord M, Hamy A-S, de Roquancourt A, et al. Triple-negative breast cancer: early assessment with 18F-FDG PET/CT during neoadjuvant chemotherapy identifies patients who are unlikely to achieve a pathologic complete response and are at a high risk of early relapse. *J Nucl Med*. United States; 2012 Feb;53(2).
35. Groheux D, Hindié E, Giacchetti S, Hamy A-S, Berger F, Merlet P, et al. Early assessment with 18F-fluorodeoxyglucose positron emission tomography/computed tomography can help predict the outcome of neoadjuvant chemotherapy in triple negative breast cancer. *Eur J Cancer*. England; 2014 Jul;50(11).
36. Groheux D, Giacchetti S, Hatt M, Marty M, Vercellino L, de Roquancourt A, et al. HER2-overexpressing breast cancer: FDG uptake after two cycles of chemotherapy predicts the outcome of neoadjuvant treatment. *Br J Cancer*. 2013;109(5).

37. Humbert O, Cochet A, Riedinger J-M, Berriolo-Riedinger A, Arnould L, Coudert B, et al. HER2-positive breast cancer: (1)(8)F-FDG PET for early prediction of response to trastuzumab plus taxane-based neoadjuvant chemotherapy. *Eur J Nucl Med Mol Imaging*. Germany; 2014 Aug;41(8).
38. Gebhart G, Gamez C, Holmes E, Robles J, Garcia C, Cortes M, et al. 18F-FDG PET/CT for early prediction of response to neoadjuvant lapatinib, trastuzumab, and their combination in HER2-positive breast cancer: results from Neo-ALTO. *J Nucl Med*. 2013;54.
39. Humbert O, Berriolo-Riedinger A, Cochet A, Gauthier M, Charon-Barra C, Guiu S, et al. Prognostic relevance at 5 years of the early monitoring of neoadjuvant chemotherapy using 18F-FDG PET in luminal HER2-negative breast cancer. *Eur J Nucl Med Mol Imaging*. 2014;41.
40. Lobbes M, Prevos R, Smidt M. Response monitoring of breast cancer patients receiving neoadjuvant chemotherapy using breast MRI – a review of current knowledge. *J cancer Ther Res*. 2012;1(34).
41. Groheux D, Majdoub M, Sanna A, de Cremoux P, Hindié E, Giacchetti S, et al. Early Metabolic Response to Neoadjuvant Treatment: FDG PET/CT Criteria according to Breast Cancer Subtype. *Radiology*. Radiological Society of North America; 2015 Apr 27;
42. Humbert O, Cochet A, Coudert B, Berriolo-Riedinger A, Kanoun S, Brunotte F, et al. Role of Positron Emission Tomography for the Monitoring of Response to Therapy in Breast Cancer. *Oncologist*. 2015 Feb 1;20(2).
43. Humbert O, Riedinger J-M, Charon-Barra C, Berriolo-Riedinger A, Desmoulins I, Lorgis V, et al. Identification of biomarkers including 18FDG-PET/CT for early prediction of response to neoadjuvant chemotherapy in Triple Negative Breast Cancer. *Clin Cancer Res*. 2015 Jun 30;
44. Ulaner G a, Eaton A, Morris PG, Lilenstein J, Jhaveri K, Patil S, et al. Prognostic value of quantitative fluorodeoxyglucose measurements in newly diagnosed metastatic breast cancer. *Cancer Med*. 2013;2(5).
45. Currin E, Linden HM, Mankoff DA. Predicting breast cancer endocrine responsiveness using molecular imaging. *Curr Breast Cancer Rep*. 2011;3(4).
46. O'Flynn E a M, DeSouza NM. Functional magnetic resonance: biomarkers of response in breast cancer. *Breast Cancer Res*. 2011;13(1).
47. Coudert B, Pierga J-Y, Mouret-Reynier M-A, Kerrou K, Ferrero J-M, Petit T, et al. Use of [(18)F]-FDG PET to predict response to neoadjuvant trastuzumab and docetaxel in patients with HER2-positive breast cancer, and addition of bevacizumab to neoadjuvant trastuzumab and docetaxel in [(18)F]-FDG PET-predicted non-responders (AVATAXHER): *Lancet Oncol*. England; 2014 Dec;15(13).
48. Chevallier B, Roche H, Olivier JP, Chollet P, Hurteloup P. Inflammatory breast cancer. Pilot study of intensive induction chemotherapy (FEC-HD) results in a high histologic response rate. *Am J Clin Oncol*. 1993;16(3).
49. Ogston KN, Miller ID, Payne S, Hutcheon AW, Sarkar TK, Smith I, et al. A new histological grading system to assess response of breast cancers to primary chemotherapy: Prognostic significance and survival. *Breast*. 2003;12(5).
50. Sataloff DM, Mason BA, Prestipino AJ, Seinige UL, Lieber CP, Baloch Z. Pathologic response to induction chemotherapy in locally advanced carcinoma of the breast: a determinant of outcome. *J Am Coll Surg*. 1995;180(3).

Supplementary material

Methods: systematic search strategy

Database PubMed
 Time span from January 2000 until March 2015
 Search in Title and abstract

Category	Keywords
"Breast cancer"	breast neoplasms[mesh] OR breast neoplasm OR breast cancer OR breast tumour OR breast tumor OR breast malignan
"Imaging"	diagnostic imaging[mesh] OR imaging* OR MRI OR magnetic resonance imaging OR PET OR PET/CT OR PET-CT OR ultrasonograph* OR mammograph* OR PET/MRI OR PET-MRI OR positron emission tomograph* OR computed tomograph* OR image OR images
"Neo adjuvant therapy"	neoadjuvant therapy[mesh] OR preoperative therapy[MeSH] OR ((neoadjuvant therapy[mesh] OR neo-adjuvant OR neoadjuvant) AND (neoadjuvant therapy[mesh] OR preoperative therapy[MeSH] OR ((neoadjuvant therapy[mesh] OR neo-adjuvant OR neoadjuvant) AND (chemo OR chemotherap* OR chemo therap*))) OR ((pre-operative OR preoperative) AND (chemo OR chemotherap* OR chemo therap*))
"Outcome"	disease-free survival[mesh] OR surviv* OR survival rate[mesh] OR survival analysis[mesh] OR effective* OR cost-effective* OR treatment response* OR treatment outcome[mesh] OR complete pathologic response* OR complete pathological response* OR pathologic complete response* OR pathological complete response* OR pathologic response OR Ki67 OR Ki-67 OR MKI67
"Breast cancer subtype"	HER2 positive OR HER2/neu positive OR HER2neu positive OR HER2-neu positive OR non-luminal OR ((human epidermal growth factor receptor 2 OR receptor, erbB-2 [mesh] OR receptor, epidermal growth factor [mesh]) AND (positive)) OR (estrogen receptor-positive OR hormone receptor-positive OR estrogen receptor-positive OR oestrogen receptor-positive OR ER-positive OR hormone positive OR positive hormone receptor OR positive estrogen) OR Luminal OR triple negative OR TN OR TNBC OR ER-negative PR-negative HER2-negative OR basal-like OR basal like

Results: study characteristics

Author, year	Sample size per subtype	Age	Study design	Enrolled	Clinical stage	Monitoring technique	Monitoring interval	Neoadjuvant therapy	Response definition monitoring	PCR definition (category)	PCR rate	AUC (95% CI)	Sens, Spec, NPV, PPV, Accuracy	Setting imaging
Charehbil, 2014	HER2 – (194); ER+ (187); ER– (35)	Mean 49 years; postmenopausal (88); premenopausal (146)	Retro-spectively	July 2010 – April 2012	II and III; T1-2; T2: 128; T3/4: 92; N+: 99; N+: 123	DCE MRI 1.5 and 3.0 T	Baseline and after three cycles	TAC with (107) or without (R) (115) zoledronic acid	>30% decrease of tumour size	Miller-Payne grade 5 or ypT0is; III	17%	0.55 (0.45-0.65)	37%, 87%, 22%, 93%, 45%	DCE-MRI; 1.5 and 3.0T
Gebhart, 2013	HER2+ / HR– (43)	-	Pro-spective	Jan 2008 – May 2010	Metabolic lymph nodes (52) and distant lesions (9)	FDG-PET/CT	Baseline, week 2 and 6	(R) Lapatinib or Trastuzumab or both. All received paclitaxel	After 2 weeks ≥ 15% reduction of SUVmax; after 6 weeks ≥ 25%	Absence of invasive cancer in the breast; III	61%	-	After 2 weeks: 27%, 88%, 65%, 60%, 64% After 2 weeks: 38%, 71%, 20%, 86%, 44%	GE / Philips or Siemens PET/CT; fasted 6h before injection; 3.7 – 7.4 MBq/kg; scan at least 50 min after injection; same scanner and parameters in each institution
Groheux, 2012	TN(20)	-	Pro-spective	Enrolled within 30 months	II (9) and III (11)	FDG-PET/CT	Baseline, after two cycles	EC-D (14) or SIM (6)	≥ -42% ΔSUVmax and ≥ -50% ΔSUVmax	No evidence of residual invasive cancer in both breast tissue and lymph nodes; II	30%	ΔSUV = 0.88	≥ -42% ΔSUVmax 64%, 100%, 55%, 100%, 75%	Gemini XL PET/CT; fasted 6h before injection; scan 60 min after injection; 5MBq/kg; CT: 120 kV; 100mAs; 16 slices; 2 min per bed position
Groheux, 2012	ER+/HER2 – (64)	Mean: 52; postmenopausal (41); Premenopausal: (22)	Pro-spective	July 2007 to Oct 2011	T1(1), T2 (21), T3 (25), T4 (17);N0 (24), N1 (29), N2 (8), N3 (5)	FDG-PET/CT	Baseline, after two cycles	EC-D	≥ - 38% ΔSUVmax and ≥ -71% ΔTLG	Sataloff TA-TB; NA-NB-NC considered as responder and partial responder; IIII	6%	ΔSUVmax: 0.73; ΔTLG 0.81	ΔSUVmax: 62%, 78%; 12%; 98%; -	Gemini XL Philips; fasted 6h before; scan 60 min after injection; 5MBq/kg; 2 min per bed position; \ CT:120 kV, 100mAs;
Groheux, 2012	HER2+ (30)	-	Retro-spective	-	II (14) and III (16)	FDG-PET/CT	Baseline, after two cycles	EC-D and trastuzumab	Reduction ≥ 62% ΔSUVmax	No residual invasive disease in tumour and lymph nodes; II	53%	ΔSUVmax = 0.86	86%, 75%, 86%, 75%, 80%	Gemini XL PET/CT; fasted 6h before injection; scan 60 min after injection; 5MBq/kg; CT: 120 kV; 100 mAs; 2 min per bed position
Groheux, 2014	TN (50)	-	Pro-spective	Nov 2007 to Sept 2012	II (21) and III (29)	FDG PET/CT	Baseline, after two cycles	EC-D (20) or SIM (30)	≥ -42% ΔSUVmax and ≥ -50% ΔSUVmax	No evidence of residual invasive cancer in breast tissues and lymph nodes; II	38%	ΔSUVmax 0.80 for EC-D and 0.86 for SIM	≥ -42% ΔSUVmax: 58%; 100%; 59%; 100%; 74%	Gemini XL PET/CT; Fasted 6h before injection; scan started after 60 min after injection; 5 MBq/kg; from mid-thigh to skull with arms raised; resolution (3D): 4x4x4 mm ³ CT: 16 slices; 120kV; 100 mAs; 2 min per position

Hatt, 2013	TN(13); ER+/HER2- (26) HER2+ (12)	-	Retro- spective	July 2007 - May 2009	II (24) and III (27)	FDG PET/CT	Baseline, after two cycles	EC-D and in HER2+ EC-D plus trastuzumab	Optimal cut-off values: ΔSUVmax: -48% ΔTLG: -56% ΔMATV: -42%	Staloff scale: TA-B with NABC are considered as responder and partial responder; III	23% 0% 33%	Use of different parameters did not improve predictive value of SUVmax ΔSUVmax: 0.88 SUVpeak; 0.84 ΔSUVmean: 0.69 ΔTLG: 0.96 ΔMATV: 0.98 Use of different parameters did not improve predictive value of SUVmax	Gemini XL Philips; fasted 6h before injection; 5 MBq/kg; after 60 min mid-thigh to skull with arms raised; resolution: 4x4x4; CT: 16 slices; 120kV; 100mAs;
Humbert, 2012	TN (25)	≤50 (61) and Pro- >50 (54); mean: 51 years	Pro- spective	-	T1-2(62) T3(42); N+ (35); N+ (79)	FDG PET/CT	Baseline and just before second course NACT	FEC 100 (25); FEC 100 plus docetaxel (39); Docetaxel followed by Epirubicin and docetaxel (8); CEX (6)	No correlation between early metabolic and final pathological response	Chevallier's classification grade 1 and 2, II	36%	No correlation between early metabolic and final pathological response	C-PET Plus scanner and Gemini GXL scanner; fasted 6h before injection of F-FDG; whole body scan 60 min after injection; 2 MBq/kg (C-PET)and 5MBq/Kg (Gemini); Prone position started 80-90 min after administration
Humbert, 2014	ER+/HER2- (53) HER2+ (37)	≤50 (36) and Pro- >50 (21); postmeno- pausal (21); premeno- pausal (35)	Pro- spective	Nov 2006 – Oct 2012	I and II (26), III (28)	FDG PET/CT	Baseline and after first course NAC	TH TH +/- carboplatin (37)	ΔSUVmax ≥ 60% ΔSUVmax of -75%	No residual invasive cancer in the breast and nodes though in-situ breast residuals were allowed (ypT0/is ypN0); II	44%	AUC: 0.70 83%, 52%, (0.55- 84%, 50%, - 0.85)	Gemini GXL and TF Philips; fasted 6 hours before injection:5 MBq/ kg (GXL) 3.5 MBq/kg (TP); brain to mid-thigh after 60 min; prone position after 90 min
Koolen, 2014	ER+/HER2- (50) HER2+ (26)	Median:47 (range: 25-68)	Retro- spective	Since Sept 2008	T1 (9), T2 (66), T3 (24), T4 (8), N0 (18), N1 (61), N2 (2), N3 (26)	FDG PET/CT	Baseline, after one and three cycles and in HER2+ after three and 8 administrations	AC (53); CD(1); AC-CD(23); AC- CTC(4); PTC (26)	ΔSUVmax	Complete absence of residual tumour cells in the breast and axillary nodes; II	2% 65%	After 1 cycle, AUC: 0.61 (0.37 – 0.86) After 3 cycles, AUC: 0.87 (0.69 – 1.00) - After 3 administrations AUC 0.61 (0.33 – 0.89) After 8administrations: 0.59 (0.34-0.85)	Gemini TF Philips, Fastes 6 h before injection; 180 – 240 MBq depending on BMI; scanning after +/- 70 min; hanging breast method; 3.0 min per bed position; resolution: 2x2x2mm CT: low dose; 40mA s, 2 mm slices;
TN (31)											52%	After 1 cycle, AUC: 0.76 (0.55-0.96) After three cycles, AUC: 0.87(0.73 – 1.00)	
Koolen, 2013	ER+/HER2- (45) HER2+ (25)	Median:47 (range: 25-68)	Retro- spective	Since Sept 2008	T1 (8), T2 (59), T3 (24), T4 (7), N0 (14), N1 (57), N2(2), N3(25)	FDG PET/CT	Baseline and after first course NAC	AC (48); AC-CTC (4); AC-CD (20); CD (1); PTC (25)	Change in FDG uptake	Complete absence of residual tumour cells at microscopy, irrespective of DCIS; III	11% 68%	0.77 (0.68 – 0.87) 0.41 (0.16 – 0.67)	Gemini TF Philips, Fastes 6 h before injection; 180 – 240 MBq depending on BMI; scanning after +/- 70 min; hanging breast method; 3.0 min per bed position; resolution: 2x2x2mm CT: low dose; 40mA s, 2 mm slices
TN (25)											61%	0.85 (0.69 – 1.00)	

Loo, 2011	ER+/HER2- (103) HER2+ (38) TN (47)	Mean: 46 (range: 23-76)	Retro-spective	Between 2000 - 2008	T1 (6), T2 (97), T3 (62), T4 (23) or 3.0T N0 (28), N1 (125), N3 (11), Nk (24)	DCE MRI 1.5T or 3.0T	Baseline and after three courses or eight administrations	AC (90); AC - CD (45); CD or AD (15); Trastuzumab based (38)	Change in largest diameter	Complete absence of residual tumour cells or small number of scattered cells at microscopy; III	7%	No association between residual tumour and change in largest diameter Residual tumour after NAC associated with change in largest diameter (p<0.05) Residual tumour after NAC associated with change in largest diameter (p<0.001)	Magnetom Vision scanner 1.5T; 3.0 T Philips Achieva breast coil; gadolinium (14ml/0.1mmol/kg); 5 series at 90s interval; FOV: 310 (1.5T); 360 (3.0T)
Martoni, 2010	ER+/HER2-; (16) HER2+; (7) TN (9)	Median: 48 years (31-72)	Pro-spective	-	II (15), III (13), IV (6)	FDG PET/CT	Baseline and after second and fourth cycle	Anthracycline based and taxane based PCT	≥ -50% ΔSUVmax	Miller and Payne; 4 and 5 with NRG A and D; III	19%	-	GE medical system; Discovery LS; Fasted 6h before scanning; scan after 60-70 min after injection; 5.3 MBq/kg; 4 min per bed position; CT: 120kV 60 mA. Slices 4 a 5 mm thick
Rigter, 2013	ER+/HER2- (246)	Median 48 (range 18-68)	Retro-spective	Oct 2004 - March 2012	T1 (21), T2 (91), T3 (43), T4 (9); Na (49), Nb (40), Nc (50), Nd (98), Ne (9)	DCE MRI 1.5T or 3.0T	Baseline after three and six courses	6 x ddAC (164); 3 x ddAC - 3 x DC (82)	Difference in largest diameter	ypT0/is ypN0 /+ ypT0/is ypN0 and ypT0ypN0; III	3%	-	Magnetom Vision scanner 1.5T; 3.0 T Philips Achieva breast coil; gadolinium (14ml/0.1mmol/kg); 5 series at 90s interval; FOV: 310 (1.5T); 360 (3.0T)
Zucchini, 2013	ER+/HER2- (31) HER2+ (14) TN (15)	Median: 49 years	Pro-spective	July 2004 - March 2011	II (30) and III (23), IV (7)	FDG PET/CT	Baseline and after second PCT cycle	6 x Anthracycline taxane regimen (9); 8 x Anthracycline taxane regimen (45) 4-8 x taxane and trastuzumab (6)	≥ -50% ΔSUVmax	Miller and Payne; TRG 4 and 5 with NRG A and D; III	16%	-	GE medical system; Discovery LS; Fasted 6h before scanning; scan after 60-70 min after injection; 5.3 MBq/kg; 4 min per bed position; CT: 120kV 60 mA. Slices 4 a 5 mm thick

Abbreviations: R: Randomized; CI: Confidence Interval; NS: Not Specified; SUV: Standardized Uptake Value; pCR: pathologic complete response; AUC: Area Under Receiver Operating Curve; AC: doxorubicin and cyclophosphamide; CD: capecitabine and docetaxel; CTC: cyclophosphamide, thiotepa, carboplatin; PTC: paclitaxel, trastuzumab, carboplatin; TAC: doxorubicin followed by cyclophosphamide and docetaxel; TCAH: taxol, carboplatin, herceptin; AbCaA: abraxane, carboplatin, avastin; TCA: taxol, carboplatin; FEC: fluorouracil, epirubicin and cyclophosphamide; EC-D: epirubicin, cyclophosphamide followed by docetaxel; SIM: epirubicin and cyclophosphamide (1200 mg/m²); TH: docetaxel and trastuzumab

Results: Quadas criteria

Author	Year	1. Representative spectrum of patients	2. Selection criteria clear	3. Reference standard reliable	4. Time interval between MRI/PET and pathology	5. Whole or random sample received verification	6. Same reference standard	7. Reference standard independent of the index test	8. Description execution of MRI/PET	9. Description execution of pathology	10. Interpretation of PET/MRI blinded from reference	11. Interpretation of PET/MRI blinded from MRI/PET	12. Same clinical data available	13. Uninterpretable test results reported	14. Uninterpretable test results explained	Score (max = 14; yes = 1)
Charehill	2014	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	yes	unclear	12
Gebhart	2013	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	yes	yes	13
Gonheux	2013	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	10
Gonheux	2012	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	10
Gonheux	2013	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	10
Gonheux	2014	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	10
Hatt	2013	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	9
Humbert	2012	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	11
Humbert	2014	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	unclear	unclear	yes	yes	yes	12
Kooten	2014	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	unclear	unclear	yes	unclear	unclear	9
Kooten	2013	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	yes	yes	12
Loo	2011	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	10
Marioni	2010	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	11
Rigter	2013	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	yes	unclear	11
Zucchini	2013	yes	yes	yes	yes	yes, but not random	yes	yes	yes	yes	yes	unclear	yes	unclear	unclear	11

Results: pooled sensitivity and specificity analysis

FDG-PET/CT in ER-positive/HER2-negative

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Groheux 2013	37	1	23	3	0.62 [0.48, 0.74]	0.75 [0.19, 0.99]		
Zucchini 2013	10	0	16	5	0.38 [0.20, 0.59]	1.00 [0.48, 1.00]

MRI in ER-positive/HER2-negative

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Charehbili 2014	68	5	122	34	0.36 [0.29, 0.43]	0.87 [0.73, 0.96]		
Rigter 2013	78	4	148	16	0.35 [0.28, 0.41]	0.80 [0.56, 0.94]

FDG-PET/CT in Triple negative

Study	TP	FP	FN	TN	Cut-off value	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Groheux 2012	9	0	5	6	-42.0	0.64 [0.35, 0.87]	1.00 [0.54, 1.00]		
Groheux 2012	11	0	3	6	-50.0	0.79 [0.49, 0.95]	1.00 [0.54, 1.00]		
Groheux 2014	18	0	13	19	-42.0	0.58 [0.39, 0.75]	1.00 [0.82, 1.00]		
Groheux 2014	22	1	9	18	-50.0	0.71 [0.52, 0.86]	0.95 [0.74, 1.00]		
Martoni 2010	0	0	6	3	-50.0	0.00 [0.00, 0.46]	1.00 [0.29, 1.00]		
Zucchini 2013	0	0	11	4	-50.0	0.00 [0.00, 0.28]	1.00 [0.40, 1.00]		

50% threshold

42% threshold

FDG-PET/CT in HER2-positive

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Groheux 2013	12	4	2	12	0.86 [0.57, 0.98]	0.75 [0.48, 0.93]		
Humbert 2012	15	2	8	12	0.65 [0.43, 0.84]	0.86 [0.57, 0.98]		
Humbert 2014	27	12	5	13	0.84 [0.67, 0.95]	0.52 [0.31, 0.72]		
Zucchini 2013	2	0	8	4	0.20 [0.03, 0.56]	1.00 [0.40, 1.00]

CHAPTER 6

Exploratory cost-effectiveness analysis of response-guided neoadjuvant chemotherapy for hormone positive breast cancer patients

Anna Miquel-Cases

Valesca P Retèl

Bianca Lederer

Gunter von Minckwitz

Lotte MG Steuten

Wim H van Harten

Accepted with minor revisions

Abstract

Purpose: Guiding response to neoadjuvant chemotherapy (guided-NACT) allows for an adaptative treatment approach likely to improve breast cancer survival. In this study, our primary aim is to explore the expected cost-effectiveness of guided-NACT using as a case study the first randomized control trial that demonstrated effectiveness (GeparTrio trial).

Materials and Methods: As effectiveness was shown in hormone-receptor positive (HR+) early breast cancers (EBC), our decision model compared the health-economic outcomes of treating a cohort of such women with guided-NACT to conventional-NACT using clinical input data from the GeparTrio trial. The expected cost-effectiveness and the uncertainty around this estimate were estimated via probabilistic cost-effectiveness analysis (CEA), from a Dutch societal perspective over a 5-year time-horizon.

Results: Our exploratory CEA predicted that guided-NACT as proposed by the GeparTrio, costs additional €67, but results in 0.014 QALYs gained per patient. This scenario of guided-NACT was considered cost-effective at any willingness to pay per additional QALY. At the prevailing Dutch willingness to pay threshold (€80.000/QALY) cost-effectiveness was expected with 79% certainty.

Conclusion: This exploratory CEA indicated that guided-NACT (as proposed by the GeparTrio trial) is likely cost-effective in treating HR+ EBC women. While prospective validation of the GeparTrio findings is advisable from a clinical perspective, early CEAs can be used to prioritize further research from a broader health economic perspective, by identifying which parameters contribute most to current decision uncertainty. Furthermore, their use can be extended to explore the expected cost-effectiveness of alternative guided-NACT scenarios that combine the use of promising imaging techniques together with personalized treatments.

Introduction

Neoadjuvant (preoperative) chemotherapy (NACT) is an option in patients with breast cancer. Equally effective as adjuvant chemotherapy [1,2], this approach allows direct and early observation of treatment response [3]. Based on this response, patient's further systematic treatment can be tailored, i.e. responders continue with the same initial treatment, and non-responders can be switched to a presumably non-cross resistant regimen. This adaptive treatment approach is likely to improve breast cancer survival.

The GeparTrio trial [4] presents the first long-term survival results (overall survival; OS and disease free survival; DFS) of guided-NACT in breast cancer. In this trial, 2012 early breast cancer (EBC) women were initially treated with two cycles of docetaxel, doxorubicin, and cyclophosphamide (TAC) followed by response assessment by palpation and ultrasound. Thereafter, patients classified as early responders were randomly assigned to four or six additional TAC cycles, and patients classified as non-responders to four cycles of TAC or four cycles vinorelbine and capecitabine (NX) before surgery (Fig1). For the survival analysis the two investigational response-guided arms (8xTAC and 2xTAC/4x NX) were grouped and compared with the conventional therapy arms (6xTAC). No significant differences in OS were observed, however a longer DFS after guided-NACT was seen in the subgroup of hormone-receptor positive (HR+) patients (hazard ratio 5-years DFS = 0.56).

The interpretation of these results is that intensifying the same chemotherapy to respondents, or switching to NX in non-respondents, only works in HR+ patients. While the lack of effectiveness seen in HR-/HER2+ patients could be justified by the lack of Trastuzumab administration in this trial, in the case of HR-/HER2-, this could be consequence of treatment ineffectiveness; there is a large body of evidence suggesting that in this subgroup there may be other treatments beyond chemotherapy [5].

The results of this study need to be interpreted with caution for several reason: 1) they rely on a secondary exploratory subgroup analysis; 2) they are the first to provide such an indication for guided-NACT and need validation, especially in the context of current therapeutic decision-making (as Trastuzumab was not used); and 3) there is no clear understanding of the underlying reason for its single benefit to HR+ patients only (whether that is direct consequence of the cytotoxic effect from the regimes used, or whether that is caused from an indirect endocrine effect causing chemotherapy induced amenorrhea [6,7]). Our interpretation is that this hypothesis needs to be prospectively tested before guided-NACT as investigated in this trial is ready for routine clinical practice in HR+ breast cancer.

If this scenario of guided-NACT proves effective, cost-effectiveness will play a central role in adoption and reimbursement decision-making. Hence, a timely explorative CEA to estimate its expected cost-effectiveness is warranted. This study aims at determining the expected cost-effectiveness of guided-NACT as proposed by the GeparTrio trial using input clinical data from the trial.

Figure 1: Decision tree and Markov model. Decision nodes (■) are points at which the patient or health provider makes a choice. Chance nodes (●) are points at which more than one event is possible but is not decided by neither the patient or health provider. During the 1st model cycle, patients receive the intervention; response-guided neoadjuvant chemotherapy (NACT), starting with 2xTAC followed by 4xNX (unfavorable at monitoring) or by 6xTAC (favorable at monitoring), or conventional-NACT, with equal treatment of 6xTAC to all patients, followed by surgery. In the following 4-year cycles, the Markov model simulates the clinical evolution of the patients, TAC docetaxel, doxorubicin, and cyclophosphamide, NX vinorelbine and capecitabine

Materials and Methods

Treatment strategies compared

Two NACT interventions were compared: *Guided-NACT* (as presented in the GeparTrio trial): 2-cycles of docetaxel 75 mg/m², doxorubicin 50 mg/m², and cyclophosphamide 500 mg/m², on day 1 every 3 weeks (2xTAC), followed by monitoring with ultrasound (US) and palpation, and by either 6xTAC or 4 courses of vinorelbine 25 mg/m² on day 1 and 8 plus capecitabine 1.000 mg/m² orally twice a day on day 1 through 14, every 3 weeks (4xNX) if patients were favorable or unfavorable respondents at monitoring respectively, following published criteria [8].

In short, favorable response was defined as a “ $\geq 50\%$ reduction in the product of the two largest perpendicular diameters of the primary tumor” assessed at the end of the second cycle and before surgery. *Conventional-NACT*: Treatment with 6xTAC without monitoring. Within the same year, all patients underwent surgery (classified as either mastectomy only, or breast-conserving-surgery (BCS) with radiotherapy).

Model overview

A Markov model (Microsoft Excel 2010, Microsoft Corporation, Redmond, WA) estimated the health-economic consequences of treating 50-years old [8] HR+ EBC women with guided-NACT vs. conventional-NACT. The model with three health-states: disease free (DFS), relapse (R, including local, regional, and distant) and death (D, including breast cancer and non-breast cancer), simulated the clinical evolution of these patients over a time-horizon of 5-years (Fig 1). Patients entered the model in the DFS health-state, after completing NACT and surgery, classified as true-favorable, true-unfavorable, false-favorable and false-unfavorable respondents of NACT at monitoring (definitions in table 1). The “gold standard” for NACT response was the 5-years relapse free survival (RFS), as it provides a reasonable threshold to capture all relapses related to NACT response [9].

Table 1: Definitions of true-positive, false-positive, true-negative and false-negative patients in our study

Group of patients	Definition
True favourable	Patient that is classified as favourable at monitoring, continues receiving 6xTAC, and after 5 years of follow up is classified as favourable due to absence of relapse event
False favourable	Patient that is classified as favourable at monitoring, continues receiving 6xTAC, and after 5 years of follow up is classified as unfavourable due to presence of relapse event
True unfavourable	Patient that is unfavourable at monitoring, switches to 4xNX, and after 5 years of follow up is classified as favourable due to absence of relapse event (<i>the underlying assumption is that the patient was not responding to 2xTAC but did to 4xNX, thereby demonstrating that monitoring classified the patient properly</i>)
False unfavourable	Patient that is unfavourable at monitoring, switches to 4xNX, and after 5 years of follow up is classified as unfavourable due to presence of relapse event (<i>the underlying assumption is that the patient was responding to 2xTAC and did not to 4xNX, thereby demonstrating that monitoring classified the patient wrongly</i>)

From this DFS health-state, patients could either 1) move to the R health-state, i.e., ‘relapse’; 2) move to the D health-state, i.e., ‘non-breast cancer death’; or 3) stay in the DFS health-state, i.e., ‘no event and administration of adjuvant hormonal treatment, assumed to be an aromatase inhibitor (AI)’. During the 1st year of the DFS health state, patients could incur NACT-related toxicities, including heart failure, (febrile) neutropenia, asthenia and alopecia [8]. From the R

health-state, patients could either 1) move to the D health-state, i.e., ‘breast cancer related death’; or 2) stay in the R health-state, i.e., ‘cured relapse’. We assumed that patients could only develop one relapse.

In each annual model cycle, patients moved/stayed in one of the mutually exclusive health-states, as explained above, according to transition probabilities (tps). During each year, patients cumulated life-years (LY), quality-adjusted life-years (QALYs), and costs. The costs and health-related quality-of-life (HRQoL) associated to the health-states are presented in Table 2.

Table 2: Costs and quality-of-life associated to the Markov model health-states

Health state		Year cycle	Costs	HRQoL
DFS		1 st	NACT and surgery	NACT
		+ if NACT related toxicities	Toxicity/es treatment	Disutility from toxicity
		2 nd /5 th	AI	AI
R	event	1 st	Relapse treatment	Relapse
	cured	2 nd /5 th	DFS year 2 nd /5 th	DFS year 2 nd /5 th
D	breast cancer	1 st /5 th	Palliative treatment	none
	other causes	1 st /5 th	none	none

HRQoL health related quality of life, *DFS* disease free survival, *R* relapse, *D* death, *NACT* neoadjuvant chemotherapy, *AI* aromatase inhibitors

Clinical data

The clinical data used to derive tp in our CEA is a subset of previously published data [8]; the group of HR+ patients of the GeparTrio trial. Our definition of HR+ was somewhat different from that of the original trial, as we selected positivity of the estrogen-receptor (ER+) only, thus excluding the group of progesterone-receptor positive (PR+)/estrogen-receptor negative (ER-) patients. This was reasoned by their small proportion among all cases, 92/1295 patients (7%), and by their absence of influence in ER+ prognosis [10,11]. The total number of HR+ patients included in our analysis was of 1203.

From these patients, Kaplan-Meier (KM) curves (IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp.) of RFS (interval from finishing the NACT intervention to occurrence of first relapse) and breast cancer specific survival (BCSS; interval from relapse to occurrence of breast cancer death) were derived for the group of conventional-NACT patients on one hand (n=602), and for the combined group of false-favorable and false-unfavorable patients (of the guided-NACT arm) on the other hand (n=67). No KMs nor tps were calculated for the true-favorable and true-unfavorable (with 100% response on the switch treatment) patients (n=233), whom by definition do not relapse and thereby do not die from breast cancer. The number of of false-favorable/unfavorable and true-favorable/unfavorable were derived by using the

5-year DFS threshold to the total patients receiving response-guided NACT (n=601). The formula $S(t)=\exp\{-kt\}$ where k is the hazard rate and t is time was used to derive the tps of relapse and breast cancer death from the aforementioned KM curves. Patients who suffered from toxicities were assumed to benefit equally from NACT and the same tps were applied. Non-breast cancer deaths were accounted by using age-specific death rates from the Central Bureau of Statistics of the Netherlands [12].

Furthermore, from this dataset we derived data on medically significant NACT-related toxicities [13] and the type of surgeries performed. These were included in the model as proportions.

Quality of life

Utilities (preferences weights) related to model health-states, chemotherapy, AI and heart failure were derived from literature [14–16] based on EuroQoL-5D measures [17]. Utility scores for febrile neutropenia, asthenia and alopecia were derived by subtracting toxicity related dis-utilities in breast cancer [18] to the baseline chemotherapy utility. The same method was used to derive the utility score for neutropenia, but using non-small-cell-lung-cancer literature as a proxy [19] owing to absence of more specific data in the breast cancer literature. Utility scores for both surgery types were assumed equal [20–22]. No literature on the effect of monitoring on HRQoL was found, thus it was assumed unaltered.

Costs

Costs (€2013) included direct medical and non-medical costs (i.e., traveling costs), and costs of productivity losses (friction cost method [23]). Drug resource use (calculated for patients of 60 Kg and body-surface area of 1.6 m²), estimates on direct-non medical costs and costs of productivity losses were derived from the GeparTrio protocol and their unit costs from Dutch sources on costs and prices [24–26] or literature [27,28]. Costs of treating toxicities [29–32], of surgery [33], of radiotherapy [33] and of the model health-states [34] were also derived from literature. Costs of monitoring included one breast examination by palpation (counted as one medical visit) and a sonography [35]. We used exchange currencies [36] when needed, and the consumer price index to account for inflation [37].

Values for tps, HRQoL data and costs are presented in S1 Table.

Base-case cost-effectiveness analysis

Effects were expressed in LYs and QALYs, costs as mean cost per patient, and cost-effectiveness as the incremental cost-effectiveness ratio (ICER; difference in expected costs divided by the difference in expected QALYs for the guided-NACT vs. conventional-NACT strategy). The ICER was compared to the prevailing Dutch threshold for cost-effectiveness of severe disease (€80.000/QALY) [38]. To facilitate the adoption decision, the ICER was arranged into the net monetary benefit (NMB). If the expected NMB is >0 , guided-NACT is cost-effective and a positive adoption recommendation follows [39].

Probabilistic sensitivity analysis

Uncertainty around the ICER estimate was calculated via probabilistic sensitivity analysis (PSA) with 10.000 second order Monte-Carlo simulations of the model. For the PSA, each model parameter was entered in the model along with a distribution (S1 Table). We discounted future costs and health effects at a 4% and 1.5% yearly rate respectively, according to the Dutch guidelines on health-economics evaluations [26]. Results were reported in cost-effectiveness acceptability curves (CEAC), which reflect the probability of each alternative to be cost-effective at a range of threshold values for cost-effectiveness.

One-way sensitivity analysis

We performed a one-way sensitivity analysis (SA) to all model parameters by varying them within one standard deviation of error or, a 25% of their base case value if this information was missing, and observed its effect on the NMB.

Results

Base-case cost-effectiveness analysis

We predicted with our model that guided-NACT prevents 1.210 relapses and 102 breast cancer deaths in 10.000 treated patients over a period of 5-year. This translated into 0.011 LYs and 0.014 QALYs gained. Furthermore, we observed that while switching response to 4xNX only added €6.199, continuing with 6xTAC added €21.837. Differences came from a combination of high drug costs in the TAC regimen (highest costs per cycle: T = €1065 and pegfilgrastim = €1161), vs the NX regimen (highest costs per cycle: N = €201 and X = €160), and a higher frequency of costly adverse events. Favorable respondents (8xTAC) were the most costly patients, followed

by conventionally treated patients (6xTAC) and unfavorable respondents (2xTAC/4xNX). Overall, guided-NACT was more expensive than conventional-NACT due to having 65% of patients assigned to 8xTAC. However, as this was more effective than conventional-NACT, the resulting discounted ICER was cost-effective (€4.707/QALY, under a €80.000/QALY, corresponding with a NMB of €1.068).

Probabilistic sensitivity analysis

The CEAC showing the cost-effectiveness of guided-NACT at different willingness to pay thresholds is presented in Fig 2. This shows that guided-NACT is expected cost-effective at any willingness to pay per additional QALY. At the Dutch willingness to pay threshold of €80.000/QALY, guided-NACT was expected cost-effective with 79% certainty.

Results for the base-case CEA and the PSA are presented in Table 3.

Sensitivity analysis

In one-way SA, the NMB remained cost-effective at all parameters values tested, except at low specificity values (55%) and high sensitivity values (100%), where the NMB became negative. Furthermore, an increase in the proportion of relapses and deaths in the conventional-NACT strategy, an increase in the costs of the R health-state and a decrease in the costs of NX markedly increased cost-effectiveness (Fig 3).

Table 3: Results of the base-case cost-effectiveness analysis and the probabilistic sensitivity analysis

Base-case CEA									PSA
Strategy	Costs (€)	LY	QALY	ΔLY	ΔQALY	Δcosts	ICER (€/QALY)	INB (€)	Prob. (%)
Guided-NACT	80.937	4,717	3,324	0,011	0,014	67	4.707	1.068	79
Conventional- NACT	80.871	4,706	3,310	-	-	-	-	-	21

Figure 2: Cost-effectiveness acceptability curves. They show the probability of response-guided neoadjuvant chemotherapy (NACT) and conventional-NACT of being cost-effective at different levels of willingness-to-pay threshold (WTP). At WTP thresholds below €80.000/QALY, response-guided NACT had a higher probability of being cost-effective, ranging from 60% at €10.000/QALY to 79% at the Dutch WTP threshold for severe diseases of €80.000/QALY

Figure 3: One-way sensitivity analysis to all model parameters. We explored how varying model parameter values could affect the net monetary benefit (NMB). If this became negative, it means that response guided neoadjuvant chemotherapy became cost-ineffective. The NMB remained cost-effective at all parameters values tested, except at specificity of 55% and sensitivity of 100%, where the NMB became negative. Furthermore, an increase in the proportion of relapses and deaths in the conventional-NACT strategy, an increase in the costs of the R health-state and a decrease in the costs of NX markedly increased cost-effectiveness.

Discussion

Response-guided NACT is likely to improve breast cancer survival. The first RCT to demonstrate this was the GeparTrio trial. It showed that guiding versus not guiding NACT improved the 5-year survival of HR+ EBC with a HR of 0.56. Although this trial was limited by several reasons (that we listed in the introduction) and requires prospective validation before it can be considered for use in routine clinical practice, it provides the first example of guided-NACT in breast cancer.

The results of our study suggest that guided-NACT as proposed by the GeparTrio trial is expected to be cost-effective (compared to conventional-NACT) at any willingness to pay threshold. This means that its additional €670.000 are expected to be outweighed by the prevention of 1.210 relapses and 102 breast cancer deaths in 10.000 treated patients over a period of 5-years. At a specific Dutch threshold for cost-effectiveness of €80.000/QALY, the probability that guided-NACT was cost-effective was of 79%. We are not aware of other cost-effectiveness studies on guided-NACT. Our results can therefore not yet be compared to other estimates.

The observed higher incremental gain in terms of QALYs than LYs (0.014 and 0.011) was explained by a higher proportion of relapsed patients (with lower HRQoL) in the conventional-NACT compared to the guided-NACT strategy (2.372 vs. 1.162). These differences were evidently driven by the HR of the GeparTrio trial that suggested that guiding NACT reduced cancer-related events to half of those observed with conventional NACT. In terms of costs, we observed that the additional €670.000 of guided-NACT were consequence of having 65% of patients assigned to 8xTAC, the most costly regimen of the model. Costs were higher in the 8xTAC regimen, followed by the 6xTAC regimen and 2xTAC/4xNX regimen. This order was an aftereffect of the differential costs between Docetaxel and Capectiabine (Docetaxel is ~100 times higher than that of Capectiabine of NX regimen) combined with the frequency of costly adverse events in the TAC regimens. As 35% of patients in the guided-NACT strategy received the low costs and presumably effective 2xTAC/4xNX regimen, it seems reasonable to assume that this contributed to guided-NACT cost-effectiveness.

Our one-way SA identified monitoring performance as the main driver of cost-effectiveness, as this was the only parameter that lead to cost-ineffectiveness. The NMB became negative at low specificity values and at high sensitivity values. This was mainly consequence of an increase of patients that received the costly treatment TACx8 i.e., true-favorable patients at high sensitivities and false-favorable patients at low specificities. Optimal performance requires a trade-off between sensitivity and specificity. Given false-favorable patients are the patients that neither benefit from TACx2 nor TACx6, while receiving the most costly treatment, in this intervention specificity should be prioritized. Recent literature has shown that MRI and PET/CT are promising in this respect i.e., sensitivities and specificities of 68% and 91, and 84%-71% respectively [40,41].

Other parameters influenced the magnitude of cost-effectiveness. For example, the lower values of conventional NACT effectiveness and the lower costs of NX. These are interesting observations to explore in further cost-effectiveness studies. These can show what happens to cost-effectiveness of guided-NACT if different imaging modalities and targeted alternatives [42] are used, and these are compared to different regimens. These type of biomarker-driven guided-NACT scenarios [43–45] are expected to entail higher costs, yet their effectiveness is also expected superior [46–50]. While awaiting for evidence to emerge on them [51–53], we advocate embarking on early stage CEAs [54], as the one we have presented here. These CEAs can be used to explore via SA the effects of interactions between model parameters in cost-effectiveness. In turn, these can help identifying those scenarios that are expected to be most cost-effective for each patient subgroup, thereby guiding researchers' translational efforts on imaging and drug development.

The results of this study are specific to the guided-NACT scenario as described by the GeparTrio trial. As this is the first study that shows the effectiveness of this NACT approach using this specific chemotherapeutic regimens, it is fundamental that this evidence is further validated before any final conclusions on the cost-effectiveness of this guided-NACT scenario can be reached.

Our decision model has limitations of data availability and assumptions. Data availability was a shortcoming for two reasons: 1) when patients had to be split according to monitoring and survival outcomes, that resulted in too small sample sizes to derive reliable KM curves, and it required merging patient groups. Nonetheless, as survival modifiers like age or hormone-receptor status were homogenous in the population, we do not expect relevant survival differences if the analysis had been done separately; 2) when estimating HRQoL, as this was absent in the GeparTrio trial and had to be collected from various, sometimes suboptimal, literature sources. Our model assumptions included the inclusion of radiotherapy costs only after BCS, following recommendations by the National Institutes of Health Consensus panel on early breast cancer [55]; and the restrictive inclusion of NACT-related toxicities to frequencies $\geq 10\%$, as less frequent events were assumed to not significantly alter costs and HRQoL. Last, a limitation of the response-guided approach itself was the impossibility to distinguish in the false-favorable group, the patients truly falsely classified at monitoring from the patients irresponsive to 4xNX or NACT in general. Nonetheless, as this is a direct consequence of the use of guided-NACT, it was included as such in the model.

Conclusion

Guided-NACT (as proposed by the GeparTrio trial) is expected cost-effective in treating HR+ EBC women. While prospective validation of the GeparTrio findings is advisable from a clinical perspective, early CEAs can be used to prioritize further research from a broader health economic

perspective, by identifying which parameters contribute most to current decision uncertainty. Furthermore, their use can be extended to explore the expected cost-effectiveness of alternative guided-NACT scenarios that combine the use of promising imaging techniques together with personalized treatments.

References

1. Fisher B, Bryant J, Wolmark N, Mamounas E, Brown A, Fisher ER, et al. Effect of preoperative chemotherapy on the outcome of women with operable breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol*. 1998;16: 2672–2685
2. van der Hage JA, van de Velde CJ, Julien JP, Tubiana-Hulin M, Vandervelden C, Duchateau L. Preoperative chemotherapy in primary operable breast cancer: results from the European Organization for Research and Treatment of Cancer trial 10902. *J Clin Oncol Off J Am Soc Clin Oncol*. 2001;19: 4224–4237
3. Gralow JR, Zujewski JA, Winer E. Preoperative therapy in invasive breast cancer: reviewing the state of the science and exploring new research directions. *J Clin Oncol Off J Am Soc Clin Oncol*. 2008;26: 696–697. doi:10.1200/JCO.2007.15.9459
4. von Minckwitz G, Blohmer JU, Costa SD, Denkert C, Eidtmann H, Eiermann W, et al. Response-Guided Neoadjuvant Chemotherapy for Breast Cancer. *J Clin Oncol*. 2013;31: 3623–3630. doi:10.1200/JCO.2012.45.0940
5. Telli ML. Insight or confusion: survival after response-guided neoadjuvant chemotherapy in breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol*. 2013;31: 3613–3615. doi:10.1200/JCO.2013.51.0313
6. Yu K-D, Shao Z-M. Survival benefit from response-guided approach: a direct effect of more effective cytotoxic regimens or an indirect effect of chemotherapy-induced amenorrhea? *J Clin Oncol Off J Am Soc Clin Oncol*. 2014;32: 1282–1283. doi:10.1200/JCO.2013.53.7555
7. von Minckwitz G, Loibl S. Survival benefit from response-guided approach: a direct effect of more effective cytotoxic regimens or an indirect effect of chemotherapy-induced amenorrhea? Reply to K.-D. Yu et al. *J Clin Oncol Off J Am Soc Clin Oncol*. 2014;32: 1283–1284. doi:10.1200/JCO.2013.54.4437
8. von Minckwitz G, Blohmer JU, Costa SD, Denkert C, Eidtmann H, Eiermann W, et al. Response-guided neoadjuvant chemotherapy for breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol*. 2013;31: 3623–3630. doi:10.1200/JCO.2012.45.0940
9. Early Breast Cancer Trialists' Collaborative Group. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet*. 2005;365: 1687–1717. doi:10.1016/S0140-6736(05)66544-0
10. Hefti MM, Hu R, Knoblauch NW, Collins LC, Haibe-Kains B, Tamimi RM, et al. Estrogen receptor negative/progesterone receptor positive breast cancer is not a reproducible subtype. *Breast Cancer Res*. 2013;15: R68. doi:10.1186/bcr3462
11. Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. *The Lancet*. 2011;378: 771–784. doi:10.1016/S0140-6736(11)60993-8
12. Dutch National Center for Health Statistics [Internet]. Available: http://statline.cbs.nl/StatWeb/publication/?VW=T&DM=SLNL&PA=7052_95&D1=0-1,7,30-31,34,38,42,49,56,62-63,66,69-71,75,79&D2=0&D3=0&D4=a,10-28&HD=080509-0829&HDR=G2,G1,G3&STB=T
13. National Cancer Institute. Common Terminology Criteria for Adverse Events (CTCAE) Version 4.0. 2010.
14. Yao G, Freemantle N, Calvert MJ, Bryan S, Daubert J-C, Cleland JGF. The long-term cost-effectiveness of cardiac resynchronization therapy with or without an implantable cardioverter-defibrillator. *Eur Heart J*. 2007;28: 42–51. doi:10.1093/eurheartj/ehl382
15. Ford E, Adams J, Graves N. Development of an economic model to assess the cost-effectiveness of hawthorn extract as an adjunct treatment for heart failure in Australia. *BMJ Open*. 2012;2. doi:10.1136/bmjopen-2012-001094
16. Lidgren M, Wilking N, Jönsson B, Rehnberg C. Health related quality of life in different states of breast cancer. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil*. 2007;16: 1073-1081. doi:10.1007/s11136-007-9202-8
17. EuroQol - Home [Internet]. [cited 8 Jan 2015]. Available: <http://www.euroqol.org/>
18. Lloyd A, Nafees B, Narewska J, Dewilde S, Watkins J. Health state utilities for metastatic breast cancer. *Br J Cancer*. 2006;95: 683–690. doi:10.1038/sj.bjc.6603326
19. Nafees B, Stafford M, Gavriel S, Bhalla S, Watkins J. Health state utilities for non small cell lung cancer. *Health Qual Life Outcomes*. 2008;6: 84. doi:10.1186/1477-7525-6-84

20. Engel J, Kerr J, Schlesinger-Raab A, Sauer H, Hölzel D. Quality of life following breast-conserving therapy or mastectomy: results of a 5-year prospective study. *Breast J*. 2004;10: 223-231. doi:10.1111/j.1075-122X.2004.21323.x
21. Arndt V, Stegmaier C, Ziegler H, Brenner H. Quality of life over 5 years in women with breast cancer after breast-conserving therapy versus mastectomy: a population-based study. *J Cancer Res Clin Oncol*. 2008;134: 1311-1318. doi:10.1007/s00432-008-0418-y
22. Cortés-Flores AO, Morgan-Villela G, Del Valle CJZ-F, Jiménez-Tornero J, Juárez-Uzeta E, Urias-Valdez DP, et al. Quality of life among women treated for breast cancer: a survey of three procedures in Mexico. *Aesthetic Plast Surg*. 2014;38: 887-895. doi:10.1007/s00266-014-0384-5
23. Koopmanschap MA, Rutten FF, van Ineveld BM, van Roijen L. The friction cost method for measuring indirect costs of disease. *J Health Econ*. 1995;14: 171-189
24. Medicijnkosten [Internet]. Available: <http://medicijnkosten.nl/>
25. Farmacotherapeutisch Kompas [Internet]. [cited 6 Sep 2013]. Available: <http://www.fk.cvz.nl/>
26. L. Hakkaart - van Roijen, S.S Tan, CAM Brouwmans. 'Handling voor kosten onderzoek' 2010. College voor zorgverzekeringen, Rotterdam, 2010 - Guide for research costs - Methods and standard cost prices for economic evaluations in healthcare \ commissioned by the Health Care Insurance Board. [Internet]. [cited 3 Sep 2013]. Available: about:home
27. Retèl VP, Joore MA, Knauer M, Linn SC, Hauptmann M, Harten WH van. Cost-effectiveness of the 70-gene signature versus St. Gallen guidelines and Adjuvant Online for early breast cancer. *Eur J Cancer Oxf Engl* 1990. 2010;46: 1382-1391. doi:10.1016/j.ejca.2010.02.035
28. Frederix GW. Disease specific methods for economic evaluations of breast cancer therapies [Internet]. universiteit Utrecht. Available: about:home
29. Schilling MB. Costs and outcomes associated with hospitalized cancer patients with neutropenic complications: A retrospective study. *Exp Ther Med*. 2011; doi:10.3892/etm.2011.312
30. Wang G, Zhang Z, Ayala C, Wall HK, Fang J. Costs of heart failure-related hospitalizations in patients aged 18 to 64 years. *Am J Manag Care*. 2010;16: 769-776
31. Calhoun EA, Chang CH, Welshman EE, Fishman DA, Lurain JR, Bennett CL. Evaluating the total costs of chemotherapy-induced toxicity: results from a pilot study with ovarian cancer patients. *The oncologist*. 2001;6: 441-445
32. Lathia N, Mittmann N, DeAngelis C, Knowles S, Cheung M, Piliotis E, et al. Evaluation of direct medical costs of hospitalization for febrile neutropenia. *Cancer*. 2010;116: 742-748. doi:10.1002/cncr.24773
33. Barlow WE, Taplin SH, Yoshida CK, Buist DS, Seger D, Brown M. Cost Comparison of Mastectomy Versus Breast-Conserving Therapy for Early-Stage Breast Cancer. *JNCI J Natl Cancer Inst*. 2001;93: 447-455. doi:10.1093/jnci/93.6.447
34. Lidgren M, Wilking N, Jönsson B, Rehnberg C. Resource use and costs associated with different states of breast cancer. *Int J Technol Assess Health Care*. 2007;23: 223-231. doi:10.1017/S0266462307070328
35. Dutch Healthcare Authority - NZa.nl [Internet]. [cited 27 Feb 2014]. Available: <http://www.nza.nl/organisatie/>
36. XE currency Converter [Internet]. Available: <http://www.xe.com/>
37. OECD.Stat. OECD (2013) [Internet]. Available: <http://stats.oecd.org/index.aspx?queryid=22519#>
38. Raad voor de Volksgezondheid en Zorg. Zinnige en duurzame zorg. Zoetermeer: 2006 [Internet]. Available: http://rvz.net/uploads/docs/Advies_-_Rechtvaardige_en_duurzame_zorg.pdf
39. Stinnett AA, Mullahy J. Net health benefits: a new framework for the analysis of uncertainty in cost-effectiveness analysis. *Med Decis Mak Int J Soc Med Decis Mak*. 1998;18: S68-80
40. Cheng X, Li Y, Liu B, Xu Z, Bao L, Wang J. 18F-FDG PET/CT and PET for evaluation of pathological response to neoadjuvant chemotherapy in breast cancer: a meta-analysis. *Acta Radiol Stockh Swed* 1987. 2012;53: 615-627. doi:10.1258/ar.2012.110603
41. Wu L-M, Hu J-N, Gu H-Y, Hua J, Chen J, Xu J-R. Can diffusion-weighted MR imaging and contrast-enhanced MR imaging precisely evaluate and predict pathological response to neoadjuvant chemotherapy in patients with breast cancer? *Breast Cancer Res Treat*. 2012;135: 17-28. doi:10.1007/s10549-012-2033-5

42. Coudert B, Pierga J-Y, Mouret-Reynier M-A, Kerrou K, Ferrero J-M, Petit T, et al. Use of [(18)F]-FDG PET to predict response to neoadjuvant trastuzumab and docetaxel in patients with HER2-positive breast cancer, and addition of bevacizumab to neoadjuvant trastuzumab and docetaxel in [(18)F]-FDG PET-predicted non-responders (AVATAXHER): an open-label, randomised phase 2 trial. *Lancet Oncol.* 2014;15: 1493–1502
43. Burstein HJ, Temin S, Anderson H, Buchholz TA, Davidson NE, Gelmon KE, et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: american society of clinical oncology clinical practice guideline focused update. *J Clin Oncol Off J Am Soc Clin Oncol.* 2014;32: 2255–2269. doi:10.1200/JCO.2013.54.2258
44. Partridge AH, Rumble RB, Carey LA, Come SE, Davidson NE, Di Leo A, et al. Chemotherapy and targeted therapy for women with human epidermal growth factor receptor 2-negative (or unknown) advanced breast cancer: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol Off J Am Soc Clin Oncol.* 2014;32: 3307–3329. doi:10.1200/JCO.2014.56.7479
45. Gradishar WJ, Anderson BO, Blair SL, Burstein HJ, Cyr A, Elias AD, et al. Breast cancer version 3.2014. *J Natl Compr Cancer Netw JNCCN.* 2014;12: 542–590.
46. Loo CE, Straver ME, Rodenhuis S, Muller SH, Wesseling J, Vrancken Peeters M-JTFD, et al. Magnetic resonance imaging response monitoring of breast cancer during neoadjuvant chemotherapy: relevance of breast cancer subtype. *J Clin Oncol Off J Am Soc Clin Oncol.* 2011;29: 660–666. doi:10.1200/JCO.2010.31.1258
47. Koolen BB, Pengel KE, Wesseling J, Vogel WV, Vrancken Peeters M-JTFD, Vincent AD, et al. FDG PET/CT during neoadjuvant chemotherapy may predict response in ER-positive/HER2-negative and triple negative, but not in HER2-positive breast cancer. *Breast Edinb Scotl.* 2013;22: 691–697. doi:10.1016/j.breast.2012.12.020
48. Dogan BE, Turnbull LW. Imaging of triple-negative breast cancer. *Ann Oncol.* 2012;23: vi23–vi29. doi:10.1093/annonc/mds191
49. De Los Santos JF, Cantor A, Amos KD, Forero A, Golshan M, Horton JK, et al. Magnetic resonance imaging as a predictor of pathologic response in patients treated with neoadjuvant systemic treatment for operable breast cancer. Translational Breast Cancer Research Consortium trial 017. *Cancer.* 2013;119: 1776–1783. doi:10.1002/cncr.27995
50. von Minckwitz G, Fontanella C. Selecting the neoadjuvant treatment by molecular subtype: how to maximize the benefit? *Breast Edinb Scotl.* 2013;22 Suppl 2: S149–151. doi:10.1016/j.breast.2013.07.028
51. QuantumLeap Healthcare Collaborative. I-SPY 2 TRIAL: Neoadjuvant and Personalized Adaptive Novel Agents to Treat Breast Cancer - NCT01042379 - ClinicalTrials.gov [Internet]. [cited 19 Feb 2015]. Available: <https://clinicaltrials.gov/ct2/show/NCT01042379?term=NCT01042379&rank=1>
52. German Breast Group. A Phase III Trials Program Exploring the Integration of Bevacizumab, Everolimus (RAD001), and Lapatinib Into Current Neoadjuvant Chemotherapy Regimes for Primary Breast Cancer - NCT00567554- ClinicalTrials.gov [Internet]. [cited 19 Feb 2015]. Available: <https://clinicaltrials.gov/ct2/show/NCT00567554?term=NCT00567554&rank=1>
53. Karolinska University Hospital. A Translational Trial on Molecular Markers and Functional Imaging to Predict Response of Preoperative Treatment of Breast Cancer Early - NCT00957125 - ClinicalTrials.gov [Internet]. [cited 19 Feb 2015]. Available: <https://clinicaltrials.gov/ct2/show/NCT00957125?term=NCT00957125&rank=1>
54. Steuten LM, Ramsey SD. Improving early cycle economic evaluation of diagnostic technologies. *Expert Rev Pharmacoecon Outcomes Res.* 2014;14: 491–498. doi:10.1586/14737167.2014.914435
55. Eifel P, Axelson JA, Costa J, Crowley J, Curran WJ, Deshler A, et al. National Institutes of Health Consensus Development Conference Statement: adjuvant therapy for breast cancer, November 1-3, 2000. *J Natl Cancer Inst.* 2001;93: 979–989

Supplementary material

Supplementary Table 1

Baseline model data on proportions, survival and costs

Parameter	mean	SD	Distribution	Source
Proportions				
Responsiveness				
True favorable	0,510	0,031	Dirichlet	⁴
True unfavorable	0,137	0,059	Dirichlet	⁴
False favorable	0,267	0,043	Dirichlet	⁴
False unfavorable	0,087	0,026	Dirichlet	⁴
Surgery				
True favorable undergoing lumpectomy	0,655	0,040	beta	⁴
True unfavorable undergoing lumpectomy	0,679	0,052	beta	⁴
False favorable undergoing lumpectomy	0,568	0,072	beta	⁴
False unfavorable undergoing lumpectomy	0,360	0,094	beta	⁴
Conventional-NACT undergoing lumpectomy	0,636	0,020	beta	⁴
Toxicities (>10% incidence ^{a)})				

		TACx6	0,421	0,019	beta	52
Neutropenia		TACx8	0,483	0,019	beta	52
		TAC/NX	0,235	0,024	beta	52
Febrile neutropenia		TACx6	0,074	0,010	beta	52
		TACx8	0,103	0,012	beta	52
Asthenia		TACx6	0,118	0,012	beta	52
		TACx8	0,154	0,014	beta	52
Heart failure		TACx6	0,009	0,004	beta	52
		TACx8	0,006	0,002	beta	52
		TAC/NX	0,007	0,005	beta	52
Alopecia		TACx6	0,104	0,012	beta	52
		TACx8	0,115	0,012	beta	52
Transition probabilities						
Relapse						
False favorable/unfavorable		Tp1	0,069	0,031	beta	4
		Tp2	0,092	0,035	beta	4
		Tp3	0,156	0,044	beta	4
		Tp4	0,243	0,052	beta	4
		Tp5	0,243	0,052	beta	4

True favorable/unfavorable	Tp1, tp2, tp3, tp4 and tp5	0,000	NA	fixed	
Conventional NACT	Tp1	0,038	0,008	beta	⁴
	Tp2	0,072	0,010	beta	⁴
	Tp3	0,070	0,010	beta	⁴
	Tp4	0,059	0,010	beta	⁴
	Tp5	0,059	0,010	beta	⁴
Breast cancer death					
False favorable/unfavorable	Tp1	0,000	NA	fixed	assumption
	Tp2	0,001 ^b	0,004	beta	⁴
	Tp3	0,049	0,026	beta	⁴
	Tp4	0,055	0,028	beta	⁴
	Tp5	0,090	0,035	beta	⁴
Conventional NACT	Tp1	0,000	NA	fixed	assumption
	Tp2	0,016	0,011	beta	⁴
	Tp3	0,008	0,008	beta	⁴
	Tp4	0,030	0,015	beta	⁴
	Tp5	0,083	0,024	beta	⁴

Utilities						
TAC	0,620	0,039	beta		22	
NX	0,620	0,039	beta		22	
Anastrozole	0,774	0,049	beta		22	
Neutropenia	0,530	0,015	beta		25	
Heart failure II & IV	0,594	NA	beta		21	
Heart failure III	0,590	0,020	beta		21	
Heart failure IV	0,505	0,049	beta		21	
Febrile neutropenia	0,470	0,085	beta		24	
Asthenia	0,505	0,099	beta		24	
Alopecia	0,506	0,099	beta		24	
Relapse	0,732	0,031	beta		22	
Disease free survival	0,935	0,020	beta		22	

Costs							
Parameter	Unit costs	Unit measure	Mean resource use	Mean cost	SD ^c	Distribution	Source
Chemotherapy/Hormone therapy							
TAC	Dir. Med (total)			€2.572	€643	Gamma	53,54
	Docetaxel	108 mg	1.11	€1.065	€266	Gamma	53
	Doxorubicin	90 mg	0.89	€183	€46	Gamma	53
	Cyclophosphamide	1080 mg	0.74	€33	€8	Gamma	32
	Day care	Day	1	€279	€70	Gamma	53
	Oncologist's visit	Visit	1	€109	€27	Gamma	53
	Dexamethasone IV	5 mg	4	€14	€3	Gamma	53
	Dexamethasone OA	10 mg	5	€3	€1	Gamma	53
	Ciprofloxacin	500 mg	9	€1	€0.3	Gamma	54
	Ondasentron ^d	8 mg	6	€36	€9	Gamma	53
	Pegfilgrastim	1 mg	1	€1161	€290	Gamma	54
	Dir. Non-Med	Day	1	€3	€1	Gamma	32
	Prod. Loss	Day	3	€753	€188	Gamma	32
	Total			€3.639	€910	Gamma	32,53,54

NX	Dir. Med (total)					€791	€198	Gamma	53
	Vinorelbine		€91	36 mg	2.22	€201	€50	Gamma	53
	Capecitabine		€16	4.500 mg	9.96	€160	€40	Gamma	53
	Day care		€279	Day	1	€279	€70	Gamma	32
	Oncologist's visit		€109	Visit	1	€109	€27	Gamma	53
	Dexamethasone IV		€3	5 mg	1.60	€5	€1	Gamma	53
	Ondansetron		€6	8 mg	6	€36	€9	Gamma	53
	Dir. Non-Med		€3	Day	2	€6	€1	Gamma	32
	Prod. Loss		€251	Day	3	€753	€188	Gamma	32
	Total					€1.550	€112	Gamma	32, 53
	Dir. Med (total)					€193	€48	Gamma	53, 55
	Anastrozole		€0.05	20 mg/day	365	€18.38	€4.59	Gamma	53
Anastrozole (1 year)	Dexa scan		€174	Scan	1	€174	€44	Gamma	55
	Dir. Non-Med		€3	Day	1	€3	€1	Gamma	32
	Prod. Loss		€251	Day	1	€251	€63	Gamma	32
	Total					€447	€112	Gamma	32, 53, 55

Monitoring									
Ultrasound	Dir. Med (total)					€215	€54	Gamma	40
	Hospital costs	€163	Scan		1	€163	€41	Gamma	40
	Specialists fees	€52	Scan		1	€52	€13	Gamma	40
	Dir. Non-Med	€3	Day		1	€3	€1	Gamma	32
	Prod. Loss	€251	Day		0.125	€31	€8	Gamma	32
	Total					€ 250	€62	Gamma	32,40
Clinical examination	Dir. Med	€109	Visit		1	€109	€27	Gamma	53
	Dir. Non-Med	€3	Day		1	€3	€0.8	Gamma	32
	Prod. Loss	€251	Day		0.125	€31	€8	Gamma	32
	Total					€143	€36	Gamma	32,53
Surgery									
Mastectomy	Dir. Med	€21,451	Surgery		1	€21.451	€1.188	Gamma	57
	Dir. Non-Med	€3	Day		1	€3	€1	Gamma	32
	Prod. Loss	€251	Day		15	€3.763	€941	Gamma	32
	Total					€25.217	€6.304	Gamma	32,57

Lumpectomy and radiotherapy	Dir. Med	€21.508	Surgery and radiotherapy ^e	1	€21.508	€403	Gamma	57
	Dir. Non-Med	€3	Day	26	€78	€19	Gamma	32
	Prod. Loss	€251	Day	40	€10.036	€2.509	Gamma	32
	Total				€ 31.622	€ 7.905	Gamma	32,57
	Chemotherapy related toxicities							
Neutropenia	Dir. Med	€22.672	Episode	1	€22.672	€5.668	Gamma	58
	Dir. Non-Med	€3	Day	1	€3	€1	Gamma	32
	Prod. Loss	€251	Day	9	€2.258	€656	Gamma	32
	Total				€24.932	€6.233	Gamma	32,58
Febrile neutropenia	Dir. Med	€28.690	Episode	1	€28.699	€7.175	Gamma	58
	Dir. Non-Med	€3	Day	1	€3	€1	Gamma	32
	Prod. Loss	€251	Day	10.7	€2.685	€1.229	Gamma	32
	Total				€31.387	€7.847	Gamma	32,58

Asthenia	Dir. Med	€1.083	Episode	1	€1.083	€271	Gamma	59
	Dir. Non-Med	€3	Day	2	€6	€1	Gamma	32
	Prod. Loss	€251	Day	No reported	€976	€244	Gamma	59
	Total				€2.065	€516	Gamma	32,59
Heart failure ^f	Dir. Med	€31.528	Episode	1	€31.528	€7.882	Gamma	60
	Dir. Non-Med	€3	Day	1	€3	€1	Gamma	32
	Prod. Loss	€251	Day	6	€1.505	€376	Gamma	60
	Total				€33.036	€8.259	Gamma	32,60
Health states								
Disease free state ^g	Dir. Med (total)				€2.872	€583	Gamma	61
	In & out –patient ^g	€2.793	Episode	1	€2.793	€563	Gamma	61
	Drugs	€79	Episode	1	€79	€20	Gamma	61
	Prod. Loss. ^h	€251	Day	9.4	€2.352	€588	Gamma	61
	Total				€5.225	€1.306	Gamma	61

Local relapse									
Relapse state <									

SD standard deviation, *Dir. Med* direct medical costs, *IV* intravenous, *OA* oral administration, *TAC* docetaxel, doxorubicin, and cyclophosphamide, *NX* vinorelbine and capecitabine, *tp* transition probabilities, *NA* not applicable, *Dir. Non-Med* direct non-medical costs, *Prod.* Loss costs of productivity losses

^a Febrile neutropenia in 6x TAC was also included, although incidence was of 7,4%

^b This *tp* was zero, but to assign a distribution to it we assigned a baseline value

^c If it was missing from the data source we used 25% SD as recommended in Briggs et al ¹³

^d We selected this 5-HT3-Antagonist, but others could also be used

^e Standard radiotherapy, which consists of 25 cycles of 5 grey

^f Calculated as an average of grade III and IV toxicities

^g Source did not report travelling expenses thus were not added

^h Costs of productivity losses were calculated by using resource use of Lidgren et al ⁶¹ but with the friction method, as recommended by the Dutch guidelines

ⁱ Loss of productivity was assumed to be the same as in the metastatic state

CHAPTER 7

Cost-effectiveness and resource use of implementing MRI-guided NACT in ER-positive/HER2-negative breast cancers

**Anna Miquel-Cases
Lotte MG Steuten
Lisanne S Rigter
Wim H van Harten**

Revised submission

Abstract

Background: Response-guided neoadjuvant chemotherapy (RG-NACT) with magnetic resonance imaging (MRI) is effective in treating oestrogen receptor positive/human epidermal growth factor receptor-2 negative (ER-positive/HER2-negative) breast cancer. We estimated the expected cost-effectiveness and resources required for its implementation compared to conventional-NACT.

Methods: A Markov model compared costs, quality-adjusted-life-years (QALYs) and costs/QALY of RG-NACT vs. conventional-NACT, from a hospital perspective over a 5-year time horizon. Health services required for and health outcomes of implementation were estimated via resource modeling analysis, considering a current (4%) and a full (100%) implementation scenarios.

Results: RG-NACT was expected to be more effective and less costly than conventional NACT in both implementation scenarios, with 94% (current) and 95% (full) certainty, at a willingness to pay threshold of €20.000/QALY. Fully implementing RG-NACT in the Dutch target population of 6306 patients requires additional 5335 MRI examinations and an (absolute) increase in the number of MRI technologists, by 3.6 fte (full-time equivalent), and of breast radiologists, by 0.4 fte, while preventing 9 additional relapses, 143 cancer deaths and 0.85-fold adverse events.

Conclusion: Considering cost-effectiveness, RG-NACT is expected to dominate conventional-NACT. Furthermore, current MRI and personal capacity are likely to be sufficient for a full implementation scenario.

Introduction

Neoadjuvant (preoperative) chemotherapy (NACT) is as effective as adjuvant chemotherapy in treating breast cancer patients [1], while offering the possibility of tailoring therapy based on tumour response at monitoring [2]. Among non-invasive imaging modalities for response monitoring, contrast-enhanced magnetic resonance imaging (MRI) is generally regarded as the most accurate modality for invasive breast cancer, as it has good correlation with pathologic complete response (pCR) the most reliable surrogate endpoint of survival [3–5].

Researchers in the Netherlands Cancer Institute (NKI) have previously published criteria for monitoring NACT response with MRI [6]. This research confirmed MRI's prediction for pCR in the triple negative breast cancer subtype [7], but not in oestrogen receptor-positive (ER+) and epidermal growth factor receptor 2- negative (HER2-) tumours. This was not an unexpected finding, given the known low rates of pCR in ER-positive/HER2-negative tumors [8, 9] make it an unsuitable measure of tumour response in these tumours. Hence, to investigate their benefit from response-guided NACT (RG-NACT), a subsequent study from this group used serial MRI response monitoring as a readout of response [10]. In this study, unresponsive tumours to the first chemotherapy regimen were switched to a second, presumably, 'non-cross-resistant' regimen. Upon study completion, the tumour size reduction caused by the non-cross-resistant regimen was similar to that in initially responding tumours after the first regimen. Furthermore, relapse frequency in both groups was similar. These observations suggested that ER-positive/HER2-negative tumours do benefit from RG-NACT with MRI, despite not reaching pCR. This results are in line with those from the German Breast Group [11], which also showed survival advantage from RG-NACT in ER+ patients.

Compared to traditional NACT, RG-NACT has thus shown to positively influence ER-positive/HER2-negative patients' survival, yet comes at additional monitoring costs. Its onset costs may however be offset by a reduction in the subsequent medical costs. This can be explored via probabilistic cost-effectiveness analysis (CEA), which quantifies the probability and extent to which RG-NACT is expected to be cost-effective compared to conventional NACT as based on current evidence. Such information is of interest for health-care regulators who, under the pressure of limited resources, are increasingly using cost-effectiveness as a criterion in decision-making [12].

The ultimate goal of decision-makers is, however, the implementation of cost-effective health-care interventions into routine clinical practice. This can often be jeopardized by the lack of attention given to resource demands [13]. Implementation as described in a CEA may not always be feasible, as this assumes that all physical resources (i.e., doctors, scanners, drugs) required by the new strategy are immediately available, regardless of actual supply constraints

(or likely demand). Ignoring these constraints may result in negative consequences, from low levels of implementation through to the technology not being implemented at all [13]. Resource modelling is a method that quantitatively captures the resource implications of implementing a new technology. While this approach has scarcely been used in health-care decision-making, it can be of great help to health services planners who are challenged by implementation issues normally not addressed in CEAs.

Our aim is thus to estimate the expected cost-effectiveness and resource requirements of implementing RG-NACT with MRI for the treatment of ER-positive/HER2-negative breast cancers using The Netherlands as a case study population. This information can act as reference for health-care regulators and health services decision-makers worldwide, on the health and economic value of RG-NACT and the resources required for its implementation

Methods

This study followed the Consolidated Health Economic Evaluation Reporting Standards (CHEERS) checklist and did not require ethical approval.

Treatment strategies

Two strategies were considered for the treatment of ER-positive/HER2-negative breast cancer women; RG-NACT and conventional-NACT (Figure 1). RG-NACT followed our single-institution neoadjuvant chemotherapy program [10]: treatment with NACT 1 (AC, doxorubicin 60 mg m⁻² and cyclophosphamide 600 mg m⁻² on day 1, every 14 days, with PEG-filgrastim on day 2) for three courses (3x) followed by MRI scanning and subsequent classification into 'favourable' or 'unfavourable' responders to NACT, defined by previously published criteria [6]). Favourable patients continue with additional 3xNACT 1, and unfavourable patients switch to 3xNACT 2 (DC, docetaxel 75 mg m⁻² on day 1, every 21 days and capecitabine 2x1000 mg m⁻² on days 1–14). Conventional-NACT represented current practice: treatment with 6xAC. Following NACT, all patients underwent surgery, radiation therapy when indicated, and at least 5-years of endocrine treatment according to protocol.

Implementation scenarios

We performed the cost-effectiveness and resource modelling analysis for two implementation scenarios in the Netherlands, i.e. current implementation and full implementation. These scenarios were adopted in a hypothetical cohort of 6306 patients, reflecting the Dutch target population

of stage II/III ER-positive/HER2-negative breast cancers. These are patients with the same baseline characteristics as those of our neoadjuvant chemotherapy program, and thus, were RG-NACT seems beneficial [10]. The current implementation scenario is defined as the number of stage II/III ER-positive/HER2-negative breast cancer patients currently treated with RG-NACT divided by all stage II/III ER-positive/HER2-negative breast cancer patients. The full implementation scenario considers the use of RG-NACT in the entire stage II/III ER-positive/HER2-negative breast cancer population. Although this is not entirely likely, there is always a percentage of non-compliant providers, we decided to present the maximum possible resource use of RG-NACT. The number of patients currently treated with RG-NACT was calculated as the number of scans performed in the Netherlands (assuming 1 scan/patient) [14] minus the number of scans performed for other disease areas than oncology [15], other cancers than breast [16], other applications than guiding response to therapy [17], other stages than II/III [18], and other receptor expressions than ER-positive/HER2-negative [19]. The entire stage II/III ER-positive/HER2-negative breast cancer population was estimated by multiplying the 2013 breast cancer prevalence in the Netherlands (The Netherlands Cancer Registry) by the proportion of patients with stage II/III ER-positive/HER2-negative breast cancer (calculations presented in Table 1).

Table 1: Current implementation scenario calculation.

Formula to derive current implementation of response-guided NACT in the Netherlands:

Number of stage II-III, ER+/HER2-breast cancer currently treated with response-guided NACT		257	4%
Number of eligible stage II-III, ER+/HER2-breast cancer		6.306	
	#	Source	
Calculations of the number of stage II-III, ER+/HER2-breast cancer currently treated with response-guided NACT			
MRI scans performed in the Netherlands	843.765	[14]	
in oncology	202.503	[15]	
for response-guided NACT	2.430	[17]	
in stage II-III, ER+/HER2- breast cancer	257	[16, 18, 19]	
Calculations of number of eligible stage II-III, ER+/HER2-breast cancer			
Incidence of breast cancer patients in the Netherlands	14.326	[63]	
With stage II-III, ER+/HER2-breast cancer	6.306	[16, 19]	

Model overview

We developed a Markov model to estimate mean differences in clinical effects and costs of treatment with RG-NACT vs. conventional-NACT from a Dutch hospital perspective. For each treatment strategy, the model simulated the transitions of a hypothetical cohort of stage II/III ER-positive/HER2-negative breast cancer patients over three health-states: disease free (DFS), relapse

(R, including local, regional, and distant) and death (D, including breast cancer and non-breast cancer), during a 5-year time horizon (Figure 1). The model was programmed in Microsoft Excel (Redmond, Washington: Microsoft, 2007. Computer Software).

Figure 1: Decision analytic model to compare the health-economic outcomes of treating ER-positive/HER2-negative stage II-III breast cancer patients with response-guided NACT vs. conventional-NACT. Decision nodes (■); patient or health provider makes a choice. Chance nodes (●); more than one event is possible but is not decided by neither the patient or health provider. Abbreviations: NACT=neoadjuvant chemotherapy; RFS=relapse free survival; DFS=disease free survival; R=relapse; D=death; AC= cyclophosphamide, doxorubicine; DC= docetaxel, capecitabine.

Upon completion of the NACT intervention, patients in each cohort entered the model in the DFS state (Figure 1). Patients treated under the RG-NACT strategy entered the DFS model state classified as true-favourable, true-unfavourable, false-favourable and false-unfavourable respondents of NACT at monitoring by using the 5-year RFS (relapse free survival) as the “gold standard” for NACT response. This was considered a sensible assumption to capture all relapses related to NACT response [21]. Definitions for true-favourable, true-unfavourable, false-favourable and false-unfavourable respondents are presented in supplementary 1.

In year 1 of the DFS health-state, patients were attributed the costs and health related quality-of-life (HRQoL) of the NACT intervention, except when there was an incidental MRI finding or when they suffered from chemotherapy-related toxicities (Terminology for Adverse Events grades 3 and 4 [22]); vomiting, neutropenia, hand-foot-syndrome (HFS), desquamation and congestive heart failure (CHF) [23, 24]). In these situations, there was NACT interruption and temporary changes in costs and HRQoL, except for HFS and desquamation. For these toxicities there is no other curative treatment than time, thereby, they were exempt of costs. From the DFS health-state,

patients could either move to the R health-state, i.e., 'relapse event'; move to the D health-state, i.e., 'non-breast cancer death event'; or stay in the DFS health-state, i.e., 'no event'. From the R health-state, patients could either move to the D health-state, i.e., 'breast cancer or non-breast cancer related death event'; or stay in the R health-state, i.e., 'cured relapse'. We assumed that patients could only develop one relapse. In the 5th-year of the model, patients could incur long-term NACT-related toxicities, including myelodysplastic syndrome (MDS) and acute myeloid leukaemia (AML) [25].

Model input parameters

Input model parameters are presented in table 2.

Clinical

The proportions of favourable and unfavourable patients at monitoring and after 5-years of NACT were retrieved from an updated version of the individual patient data from Rigter *et al* [10]. The transition probabilities (tp) simulating a relapse and a breast cancer death event were derived from Kaplan-Meier (KM) curves. The first from a KM of RFS (interval from finishing the NACT intervention to occurrence of first relapse) and the second, from a KM of breast cancer specific survival (BCSS; interval from relapse to occurrence of breast cancer death). The KMs were either constructed uniquely with raw data of Rigter *et al* [10], or by using additional assumptions, which we explain in detail below. Calculations were performed in SPSS (IBM Corp. Released 2013. IBM SPSS Statistics for Windows, Version 22.0).

RG-NACT: The tps for the group of false-unfavourable and false-favourable patients were derived by using KMs and the formula $tp(t_u) = 1 - \exp\{H(t - u) - H(t)\}$ [26], where u is the length of the Markov cycle (1 year) and H is the cumulative hazard. Data for the KM of RFS came from 25 relapsed patients from Rigter *et al* [10], and that of BCSS, from literature [27]. The tps of relapse and breast cancer death for the true-favourable and true-unfavourable patients were assumed to be zero at all times, as these patients do not relapse nor die from breast cancer (see supplementary 1). **Conventional-NACT:** tps were derived from KM curves, with data from the complete dataset of Rigter *et al* [10] for the RFS curve and data from literature [27] for the BCSS curve. The formula to derive tps was: $tp(tu) = 1 - \exp\{1/\tau(H(t - u) - H(t))\}$ [26], where τ is the treatment effect or hazard ratio (HR) of RG-NACT vs. conventional-NACT. This formula allowed calculating the tps from a "hypothetical" control arm, which was inexistent in the Rigter *et al* [10] study. The used HRs were 0.5 for the RFS curve, and 0.64 for the BCSS curve. While the first was assumed, the second was derived from literature and set equal to the reported HR of OS in a similar population of ER-positive breast cancers where RG-NACT vs. conventional-NACT was being compared [11].

As these assumptions could affect our cost-effectiveness results, we performed a one-way and two-way sensitivity analysis (SA) to the HRs (range 0.1 - 1.5).

The tps of non-BC related deaths (i.e., transition from any state to D) were accounted for by using Dutch life tables [28]. The occurrence of vomiting, neutropenia, HFS and desquamation under 3xAC and 3xDC, were derived from literature [24]. When a patient received both 3xAC and 3xDC the probability of vomiting and neutropenia was represented as the combined probability of two independent events ($P(A \text{ and } B) = P(A) * P(B)$). The probability of occurrence of CHF due to the administration of anthracyclines was accounted for in the 1st-year of the model and was dose-dependent: 0.2% with 3xAC and 1.7% with 6xAC [23]. Also the probability of incidental findings at MRI was accounted for in that year [29]. The frequency of MDS and AML events was based on cumulative doses of anthracycline and cyclophosphamide [25]. Patients whose NACT was interrupted to treat toxicities were still assumed to benefit from NACT and the same relapse rate was applied.

Costs

Intervention costs comprise of chemotherapy, monitoring, chemotherapy-related toxicities and costs of confirming incidental findings. To calculate drug dosages we assumed patients of 60Kg and body-surface area of 1.6m². Drug use was derived from study protocol, and costed by using literature [30, 31] and Dutch sources on costs and prices (Dutch National Health Care Institute; Dutch Healthcare Authority; Dutch Health Care Insurance Board). Chemotherapy costs included day care and one visit to the oncologist per cycle. Costs of monitoring consisted of one MRI scan [35] and one medical visit of 1h (accounting for waiting time) [31]. Costs of treating toxicities were taken from literature [36–38]. Costs of confirming incidental findings were estimated as an average of “standard diagnostic imaging” (i.e., Ultrasound, x-Ray and bone scintigraphy) using prices from the NZA as a proxy [32]. Health state costs, i.e., follow up costs for the DFS health state and detection plus treatment costs for the R health state, were derived from literature [39]. All results were reported in 2013 Euros, using exchange currencies [40] and the consumer price index to account for inflation [41].

Health-Related Quality of life

Utilities were derived from published literature. The DFS utility was 0.78 except in the 1st-year cycle when patients either accrued the utility of the NACT regimen without toxicities i.e., 0.62 [42], the utility of the NACT regimen with toxicities i.e., 0.62 minus the utility decrements [43–45]) or the utility of anxiety in patients where incidental findings at MRI occurred i.e., 0.68 [46]. These utilities lasted for the whole cycle. The R utility was calculated as an average of the utility of local

and distant relapse [42]. All utility weights were obtained from sources using the EuroQoL EQ-5D questionnaires, except anxiety, which was derived from a Quality of Well-Being index [46]. There is no literature to suggest an effect of monitoring on HRQoL, thus this was assumed unaltered.

Scenarios and resource modelling

Additional parameters to simulate the scenarios and to perform the resource modelling exercise were added in the model. These include a parameter reflecting the RG-NACT uptake, and parameters illustrating the proportion of i) patients with MRI contraindications (impaired renal function due to the risk of developing Nephrogenic Systemic Fibrosis (NSF) [47], presence of ferrous body parts like pacemakers (mean of values reported in [48–50], and claustrophobia [51]), ii) patients with NSF [52], iii) patients with malignant incidental findings (Rinaldi et al, 2011) and iv) MRI technologists with acute transition symptoms (ATS) [53].

Cost-effectiveness analysis

The 5-year cumulative outcomes (health benefits and costs) were simulated for a cohort of 6306 individuals. The cost-effectiveness outcome measure was the incremental cost-effectiveness ratio (ICER), which is the difference in expected costs (per patient) divided by the difference in expected effects expressed as (quality-adjusted) life-years ((QA)LYs) of treating one hypothetical cohort with RG-NACT vs. treating an identical cohort with conventional-NACT. For the current implementation scenario, we compared the expected costs and QALYs of a cohort as treated with conventional-NACT, to the costs and QALYs of a cohort partially treated with RG-NACT, as dictated by the implementation rate and MRI contraindications. Patients where RG-NACT was not implemented or MRI was contraindicated were modelled as receivers of conventional-NACT. The full implementation scenario was modelled in the same way, except that the RG-NACT strategy was now applied to all patients in the cohort, except those with MRI contraindications receiving conventional-NACT.

Table 2: Input model parameters

Parameter	mean	SE	Parameters ^a	Distribution	Source
Clinical data					
Monitoring performance					
True favourable	0,53	0,04	0,53/0,04	Dirichlet	[10]
True unfavourable	0,24	0,05	0,24/0,05	Dirichlet	[10]
False favourable	0,17	0,07	0,17/0,07	Dirichlet	[10]
False unfavourable	0,07	0,09	0,07/0,09	Dirichlet	[10]
Chemotherapy related toxicities					
Vomiting					
3xAC	0,05	0,02	5/98	beta	[24]
3xDC	0,24	0,04	24/77	beta	[24]
HFS					
3xDC	0,22	0,04	23/80	beta	[24]
Neutropenia					
3xAC	0,85	0,04	86/15	beta	[24]
3xDC	0,72	0,04	74/29	beta	[24]
Desquamation					
3xDC	0,05	0,02	5/98	beta	[24]
CHF					
3xAC	0,002	0,20	1/359	beta	[23]
6xAC	0,02	0,60	11/349	beta	[23]
AML/MDS					
3xAC	0,003	0,001	12/4471	beta	[25]
6xAC	0,005	0,001	12/2372	beta	[25]
Transition probabilities					
Relapse					
RG-NACT;					
False favourable/unfavourable					
Tp1	0,14	0,06	4/24	beta	[10]
Tp2	0,29	0,08	8/20	beta	[10]
Tp3	0,47	0,09	13/15	beta	[10]
Tp4	0,44	0,09	12/16	beta	[10]
Tp5	0,40	0,09	11/17	beta	[10]
RG-NACT;					
True favourable/unfavourable					
Tp1 ¹²⁻⁵	0,00	NA	-	fixed	assumption
HR RFS					
(RG-NACT vs. conventional-NACT)					
Conventional-NACT					
Normal truncated	0,50	0,20	0,50/0,20	Normal truncated	assumption
Conventional-NACT					
Tp1	0,03	-	-	-	[10]
Tp2	0,06	-	-	-	[10]
Tp3	0,08	-	-	-	[10]
Tp4	0,05	-	-	-	[10]
Tp5	0,04	-	-	-	[10]

Breast cancer specific death						
False favourable/unfavourable						
	Tp1	0,00	NA	-	fixed	assumption
	Tp2	0,04	0,02	5/109	beta	[27]
	Tp3	0,12	0,03	14/100	beta	[27]
	Tp4	0,06	0,02	7/107	beta	[27]
	Tp5	0,19	0,04	22/92	beta	[27]
(RG-NACT vs. conventional-NACT)		0,64	0,13	0,64/0,13	normal	[11]
HR BCSS		0,00	NA	-	fixed	assumption
Conventional-NACT		0,06	-	-	-	[27]
	Tp2	0,19	-	-	-	[27]
	Tp3	0,09	-	-	-	[27]
	Tp4	0,28	-	-	-	[27]
Utilities						
Chemotherapy		0,62	0,04	94/58	beta	[42]
Neutropenia		0,53	0,01	557/488	beta	[43]
Anxiety		0,68	0,06	40/19	beta	[46]
Vomiting		0,52	0,08	17/16	beta	[44]
HFS		0,50	0,10	12/12	beta	[44]
Desquamation		0,59	0,01	1041/721	beta	[43]
CHF (average grade III/IV)		0,55	-	-	beta	[45]
CHF grade III		0,59	0,02	360/250	beta	[45]
CHF grade IV		0,51	0,05	52/50	beta	[45]
MDS/MLA		0,26	0,01	500/1423	beta	[57]
DFS		0,80	0,03	196/49	beta	[42]
R (average loco-regional and metastatic)		0,73	-	-	beta	[42]
Loco-regional relapse		0,68	0,03	226/104	beta	[42]
Metastatic relapse		0,78	0,04	104/30	beta	[42]

Scenarios and resource modelling

Incidental findings						
All	0,18	0,01	270/1265	beta	[29]	
Malign	0,20	0,02	55/270	beta	[29]	
MRI contraindications						
Impaired renal function	0,07	0.1 ^b	0.45/5.54	beta	[52]	
Gadolinium allergy	0.0003	0.01 ^c	0.08/29	-	[47]	
Body ferrous parts	0.58	0.1	0.26/4.21	beta	[48]	
Claustrophobia	0.02	0.1	0.02/0.94	beta	[51]	
Uptake	0.04		20-100%	fixed	assumption	
MRI technologists with ATS	0.26		-	fixed	[53]	
Costs						
Parameter	Unit costs	Unit measure	Mean resource use	Mean cost	SE ^d	Distribution Source
Chemotherapy						
6xAC						
Doxorubicin	€204	90 mg	5,3	€1306	€326	Gamma [31]
Cyclophosphamide	€45	1080 mg	6,4	€239	€60	Gamma [31]
Peg-filgrastim	€849	1 mg	6	€5096	€1274	Gamma [58]
Pharmacy preparation	€45	Per course	6	€267	67	Gamma NKI
Day care	€286	Day	6	€1718	€430	Gamma [30]
Oncologist's visit	€109	Visit	6	€653	€163	Gamma [31]
Total				€9279		
3xAC/ 3xDC						
Doxorubicin	€204	90 mg	3,2	€653	€163	Gamma [31]
Cyclophosphamide	€45	1080 mg	2,7	€120	€30	Gamma [31]
Peg-filgrastim	€849	1 mg	3	€2548	€637	Gamma [58]
Docetaxel	€959	108 mg	3,3	€3195	€799	Gamma [31]
Capecitabine	€27	4500 mg	29,9	€821	€205	Gamma [31]
Pharmacy preparation	€45	Per course		€267	€67	Gamma NKI
Day care	€286	Day	6	€1718	€430	Gamma [30]
Oncologist's visit	€109	Visit	6	€653	€163	Gamma [31]
Total				€9974		

Monitoring						
MRI scan	€163	Scan	1	€163	€41	Gamma [35]
Hospital costs	€52	Scan	1	€52	€13	Gamma [35]
Specialists fees				€215		
Total	€149	Episode	1	€149	€37	Gamma [35]
Confirm incidental findings						
Chemotherapy related toxicities						
Neutropenia	€14397	Episode	1	€14397	€425	Gamma [38]
Vomiting	€92	Episode	1	€92	€23	Gamma [59]
HF	€18225	Episode	1	€18225	€4556	Gamma [36]
MDS/MLA	€112946	Episode	1	€112946	€28236	Gamma [60, 61]
Health states						
DFS	€2793	Episode	1	€2793	€563	Gamma [42]
In & out -patient	€79	Episode	1	€79	€20	Gamma [42]
Total				€2872		
R						
Local relapse						
In & out -patient	€12497	Episode	1	€12497	€1692	Gamma [62]
Drugs	€2336	Episode	1	€2336	€584	Gamma [62]
Distant metastasis						
In & out -patient	€11645	Episode	1	€11645	€1346	Gamma [62]
Drugs	€5772	Episode	1	€5772	€1443	Gamma [62]
Total	€8296	Episode	1	€16125	€2074	Gamma [62]
BC death						

Abbreviations: SE= standard error; AC= cyclophosphamide, doxorubicine; DC= docetaxel, capecitabine; HFS= hand-food-syndrome; CFH= congestive heart failure; AML/ADM= acute myeloid leukaemia /myelodysplastic syndrome; MRI= magnetic resonance imaging; tp= transition probability; HR= hazard ratio; RG-NACT= response guided neoadjuvant chemotherapy; NACT= neoadjuvant chemotherapy; DFS= disease free survival; R= relapse; RFS= relapse free survival; BCSS=breast cancer specific survival; BC= breast cancer; AT5= acute transition symptoms

^a Dirichlet distribution: mean/SE, Beta distribution: α/β , Normal distribution: mean/SE

^b We assumed a SE=0.1

^c We assumed a SE=0.01

^d We assumed SE=0.25 when this was not available from literature

We performed a probabilistic sensitivity analysis (PSA) after assigning a distribution to each model parameter (Table 2). The uncertainty surrounding the model results was presented as cost-effectiveness acceptability curves (CEAC), which reflect the probability of each alternative to be cost-effective across a range of threshold values for cost-effectiveness. We discounted future costs and health effects at a 4% and 1.5% yearly rate respectively, according to the Dutch guidelines on health-economics evaluations [54]. A strategy was considered cost-effective if the ICER did not exceed the willingness-to-pay threshold of €20.000/QALY.

Resource modelling analysis

We estimated the health services required and the health outcomes experienced in each strategy. Health services required included: number of 1) MRI scans performed, 2) patients scanned per MRI, 3) Full-time equivalent (FTE) MRI technologists, 4) FTE breast radiologists and 5) confirmation of incidental findings. Health outcomes included: number of 1) relapses prevented, 2) breast cancer deaths prevented, 3) excluded patients due to contraindications, 4) patients with adverse events (including NSF, CHF, and AML/ADS), 5) patients with anxiety due to incidental findings, 6) patients with malignant incidental findings, and 7) fte MRI technologists with ATS. These outcomes were analysed deterministically for the current and full implementation scenarios and expressed for the 6306 ER-positive/HER2-negative breast cancer women. A detailed description of the calculations and sources for each outcome is presented in supplementary 2.

Volumes of health services needed were also calculated at the hospital level, which required determining the number of hospitals expected to offer RG-NACT under each scenario. For current implementation, we assumed RG-NACT to be used in the 16 hospitals of the largest Dutch hospital network currently involved in the RG-NACT trial NCT01057069 (Clinical Trials.gov). Although this trial excludes ER+ patients, we expected involved hospitals to have endorsed RG-NACT in other subtypes with single institution studies, as is the case in the NKI. For the full implementation, we considered all 113 hospitals (locations) with MRI that deliver cancer treatment (i.e., university, general and specialized hospitals), as identified from the database published by the National Public Health Atlas [55]. The presence and quantity of MRI scans per hospital was either taken from that hospital's website or based on literature [53], indicating 3 MRIs per academic hospital and 1 per general hospital.

All assumptions made were confirmed by an experienced MRI technologist in a general hospital. One-way SAs on one key-assumptions was done: 'the time required by a breast radiologist for MRI scan interpretation' (range 6.8-15 minutes).

Results

Cost-effectiveness analysis

At current implementation (4%) RG-NACT was expected to result in 0.005 QALYs gains and savings of €13 per patient. Under full implementation, RG-NACT is expected to generate 0.12 additional QALYs and savings of €328 per patient (Table 3). In both scenarios, RG-NACT is expected to dominate (be more effective and less costly) than conventional-NACT. The results of the PSAs show that at a willingness to pay threshold of €20.000/QALY, RG-NACT is expected to be the optimal strategy under the current and full implementation scenarios, with 94% and 95% certainty respectively (Figure 2).

SAs of RFS and BCSS hazard ratios (baseline values of 0.5 and 0.64 respectively), invariably showed the RG-NACT strategy to be cost-effective (Table 3). Even when LYs were slightly higher in the conventional-NACT arm (i.e., with HRs of >1), the better quality of life provided by the DC treatment of the RG-NACT strategy (lower and better tolerated adverse events) maintained the incremental QALYs for the RG-NACT strategy.

Figure 2: Cost effectiveness acceptability curves. At a willingness to pay threshold of €20.000/QALY, RG-NACT is expected to be the optimal strategy with 94% and 95% certainty under the current and full implementation scenarios respectively.

Table 3: Resource modeling and cost-effectiveness results for the current and full implementation scenarios of response-guided NACT in the Netherlands.

Cost-effectiveness analysis									
Current implementation (4%)					Full implementation (100%)				
	Costs (€)	QALYs	Δ costs (€)	Δ QALYs	ICER	Costs (€)	QALYs	Δ costs (€)	Δ QALYs
RG-NACT	28013	3.46	-13	0.005	Dominant ^a	27698	3.58	-328	0.12
Conventional-NACT	28026	3.45	-	-	-	28026	3.45	-	-
One-way and two-way sensitivity analysis									
ICER					ICER				
HR RFS	HR OS				HR RFS / BCSS				
0.1	€-12857/QALY (cost-effective)				0.1 / 0.1				
1	€2398/QALY (cost-effective)				€-10692/QALY (cost-effective)				
1.5	€-9367/QALY (cost-effective)				1.5 / 1.5				
Resource modelling analysis									
expressed in relation to the Dutch population of ER-positive/HER2-negative breast cancer women (n=6306)									
Current implementation (16 hospitals, 31 MRIs)					Full implementation (113 hospitals, 148 MRIs)				
Transition from current to full implementation									
Health services required at the country level									
No of MRIs scans performed	218				5335				
No of patients scanned per MRI	7				36				
Fte MRI technologists	0.2				3.8				
Fte breast radiologists	0.02				0.4				
	0.04 ^b (↑121%)				0.95 ^b (↑121%)				
No of confirmations of incidental findings (using standard imaging)	38				939				
Health services required at the hospital level									
No of MRIs scans performed per hospital	14				47				
No of patients scanned per MRI per hospital	7				36				
Fte MRI technologists per hospital	0.01				0.03				
Fte breast radiologists per hospital	0.001				0.004				
	0.002 ^b (↑121%)				0.001 ^b (↑121%)				

Health outcomes gained at the country level				
No of relapses prevented	0.4	9		+9
No of breast cancer deaths prevented	6	149		+143
Health outcomes lost at the country level				
No of excluded patients due to contraindications				
No of patients with NFS	40	971		+931
Fte MRI technologists with acute transient symptom	0.07	2		+2
No of patients with CHF	0.04	0.9		+1
No of patients with long term AML/ADS	106	83		-23
No of patients with anxiety due to incidental findings	23	21		-2
No of patients with malignant incidental findings	38	939		+901
	8	192		+184

Abbreviations: No= number; Fte= Full-time equivalent; MRI= magnetic resonance imaging; NSF= nephrogenic systemic fibrosis; ATS= acute transient symptom; CHF= chronic heart failure; AML/ADS= myelodysplastic syndrome/acute myeloid leukaemia.

^a RG-NACT is more effective and less costly than conventional NACT

^b if radiologists spent 15 minutes to interpret 1 MRI scan

* When possible, figures were rounded to the nearest whole number.

Resource modelling analysis

Under the current implementation scenario we calculated that over 5-years, the RG-NACT strategy requires 218 MRI scans to be performed in the target population of 6306 women, after 40 exclusions due to contraindications. With 31 MRI scans currently used for this purpose (estimated number of MRI scans in the multicentre NCT01057069 trial), 7 patients were scanned/MRI, requiring a total of 0.2 fte MRI technologists and 0.02 fte breast radiologists. At the hospital level covering a population of 6306 breast cancers, 14 MRI scans would be required for the prevalent population over a 5-year timeframe. Assuming an average capacity of 2 MRI scans/hospital (estimated weighted average of MRI scans/hospital within the multicentre NCT01057069 trial), this would translate to 7 patients scanned/MRI, demanding 0.01 fte MRI technologists and 0.001 fte breast radiologists per hospital. In terms of health outcomes, the current implementation scenario was expected to prevent 0.4 relapses and 6 breast cancer deaths, while yielding 0.07 patients with NSF. Besides, 106 patients would have a CHF, 23 patients would suffer from AML/ADS and 38 incidental findings were expected, of which 8 would be malignant. Of the required 0.2 fte MRI technologists, 0.04 fte would suffer from ATS (Table 3).

Under the full implementation scenario, we calculated that 5335 MRI scans would be needed over a 5-year period for the 6306 pertinent breast cancer population, after excluding 971 patients for contraindications. With 148 MRI scans available (estimated number of MRI scans in the estimated 113 hospitals), this would require 36 patients to be scanned/MRI for which 3.8 fte MRI technologists and 0.4 fte radiologists are needed. At the hospital level, 47 MRI scans are expected to be performed for the prevalent population of 6306 within 5-years. Assuming the mean MRI scans/hospital is 1.3 (estimated weighted average of MRIs/hospital within the estimated 113 hospitals), 36 patients would be scanned per MRI, requiring 0.03 fte MRI technologists and 0.004 fte breast radiologists per hospital. In terms of health outcomes, the full implementation scenario was expected to prevent 9 relapses and 149 breast cancer deaths, but to bring about 2 patients with NSF, 83 patients with CHF, and 21 patients with AML/ADS. Furthermore, there are 939 incidental findings expected, of which 192 would be malignant, and 0.9 fte MRI technologists are projected to get ATS (Table 3).

The transition from current (4%) to full (100%) implementation is expected to increase the number of examinations by 5117 (2347%) countrywide or by 33 (247%) per hospital, consequently demanding an increase of scan utilization (for an additional 29 patients), an increase in the number MRI technologists by 3.6 fte countrywide or by 0.02 fte per hospital, and a marginal increase in breast radiologists by 0.4 fte countrywide or by 0.003 fte per hospital. In terms of health outcomes, full implementation would diminish the number of breast cancer related deaths and relapses by 25-fold (from 6 to 149) and 23-fold (from 0.4 to 9) respectively, and the number

of CHF and AML/MDS by ~0.8-fold (from 106 to 83) and ~0.9-fold (from 23 to 21) respectively. However, these would come at the cost of a ~25-fold increase on health losses (additional 2 patients with NSF, 1 fte MRI technologist with ATS, 901 patients with anxiety due to presence of incidental findings, and 184 patients with confirmed malignant findings).

The results of the one-way SA on the radiologists' working pattern assumption showed that increasing the time required for MRI scan interpretation to 15 minutes, increased the 'fte breast radiologists' required by 121% (Table 3).

As increasing RG-NACT uptake from 4% to 100% is not realistic in a short time-frame, we explored post-hoc resource requirements and health outcomes across a range of implementation rates via one-way SA including 20%, 40%, 60% and 80% uptake. This showed that increasing implementation rates markedly increases the number of patients with MRI contraindications, the number confirmatory scans, and the number of patients with anxiety while awaiting for those (Figure 3). Simultaneously, the number of cancer deaths, and the number of patients with CHF and AML/ADS decreased consistently (by ~1.5, ~0.98 and ~0.95 -fold per 20% rate increase).

Figure 3: Influence of implementation rates on resource modelling outcomes, a) on health services required and b) on health outcomes. Abbreviations: No= number; Fte= full-time equivalent; MRI= magnetic resonance imaging; ATS= acute transition syndrome; CHF= chronic heart failure; AML/ADM= acute myeloid leukaemia / myelodysplastic syndrome; NFS= nephrogenic systemic fibrosis.

Discussion

The aim of our study was to estimate the cost-effectiveness and resource requirements of implementing RG-NACT with MRI for ER-positive/HER2-negative breast cancer patients using The Netherlands as a case study population. As RG-NACT is an emerging treatment approach and its implementation is at its onset, we performed these analyses under a current implementation scenario of 4% uptake, and under a full implementation scenario, to anticipate the outcomes of a potential wider roll-out.

At the current 4% uptake RG-NACT is expected to be less expensive and achieve more QALYs than conventional-NACT. With higher implementation rates, more patients will be treated with this cost-saving and effective strategy, rendering RG-NACT ever more dominant. At full implementation, 0.12 additional QALYs and savings of €328 per patient are expected. This is achieved despite 15% (971 out of the 6303 patients) being treated with conventional-NACT due to MRI contraindications. In both scenarios, decision uncertainty surrounding the ICERs is low (~5%).

The main drivers of advantageous survival in the RG-NACT are the HRs used to derive the hypothetical survival of the conventional-NACT strategy. Either of the HRs used (for RFS and BCSS) was below 1, thus implying less breast cancer related events in the RG-NACT strategy compared to the conventional-NACT strategy. These values were based on best available data from the GeparTrio trial [11], but this evidence is still preliminary. One- and two-way SA of these HR values demonstrated that even when survival was (slightly) higher in the conventional-NACT strategy, the better quality-of-life derived from DC treatment in the RG-NACT strategy maintained the cost-effectiveness of RG-NACT.

The cost savings of RG-NACT hinge on a satisfactory diagnostic performance of MRI. Under current diagnostic performance, 79% of patients would not yield any event in the RG-NACT strategy, compared to 76% in conventional-NACT. Although the prevention of these events came at the costs of 30% of patients receiving a more expensive treatment than conventional-NACT (>€695), as treating one relapse is even more expensive (€16125), RG-NACT was still cost saving.

The resource modelling analysis showed that increasing RG-NACT uptake rates from 4% to 100% is expected to increase the number of examinations by 5117 (2347%), consequently demanding a 5-fold increase in scans utilization, a 19-fold increase in the number MRI technologists and a 20-fold increase in the number of breast radiologists. Thereby, adapting current practice to meet these resources requires paying special attention to the availability and utilization of MRIs, as well as availability of technical personnel. For instance, fully implementing RG-NACT in the Netherlands,

were 5701 MRI examinations were performed in 2013 (considering 843765 MRI examinations [14] performed in 148 MRIs), would only require 4.5 days of additional MRI scanning per year to current MRI utilisation (given our model assumptions). Furthermore, personnel technologists and radiologists is not expected to be a limiting implementation factor either, as availability is estimated to be of 1700 MRI technologists countrywide [53] and 10 breast radiologists per hospital [56].

In terms of health outcomes gained, full implementation would diminish the number of breast cancer related deaths and relapses by 25- and 23-fold respectively, and the number of severe and costly adverse events as CHF and AML/MDS by ~ 0.8 - and ~ 0.9 -fold respectively. However, these would come at the cost of a parallel ~ 25 -fold increase in patients with NSF, MRI contraindications, MRI technologists with ATS and incidental findings causing anxiety and other diseases.

Our post-hoc analysis on resource requirements at various RG-NACT implementation rates allow identifying those that seem feasible given current resources. Considering current MRI machines and personnel capacity, RG-NACT implementation seems feasible at any rate. However, it would be interesting to further investigate whether there is sufficient capacity to handle an increase of incidental findings (requiring further diagnostic examinations), as well the cost-consequences of treating those that are diagnosed as malignant.

Our study has some limitations. A limitation of the response-guided approach itself was the impossibility to distinguish in the false-unfavourable group, patients truly falsely classified at monitoring from patients irresponsive to 3xDC or NACT in general. Yet, as this is inherent to guided-NACT, it was included as such in the model. Furthermore, we did not consider adjuvant treatment in our model, as the administration of this was similar between arms. Moreover, we considered AC, instead of a 3rd generation regimen containing taxanes as standard treatment because it was considered the best comparator for the used RG-NACT regimens. As costs of those are different, we performed a post-hoc one-way SA and found that RG-NACT would become more dominant due to increased cost savings.

While the typical CEA assumes perfect implementation of the strategy under investigation, we showed the impact of implementation rates on incremental health gains and cost-savings of RG-NACT in the Dutch population of ER-positive/HER2-negative breast cancers. Furthermore, we showed that fully implementing RG-NACT generates a ~ 24 -fold increase in health benefits, but requires MRI and personnel capacity to be increased by 5- and ~ 20 -fold. In the Netherlands, both capacities are likely to be sufficient for a full implementation scenario.

Acknowledgements

The authors gratefully acknowledge Prof. dr. Sjoerd Rodenhuis for his clinical insights, and Mirjam Franken and Prof. dr. Ruud Pijnapple for assessing the resource modeling assumptions. This project is funded by the Center for Translational Molecular Medicine (CTMM project Breast CARE, grant no.03O-104).

References

1. Mauri D, Pavlidis N, Ioannidis JPA: Neoadjuvant versus adjuvant systemic treatment in breast cancer: a meta-analysis. *J Natl Cancer Inst* 2005, 97:188–194.
2. Gralow JR, Zujewski JA, Winer E: Preoperative therapy in invasive breast cancer: reviewing the state of the science and exploring new research directions. *J Clin Oncol Off J Am Soc Clin Oncol* 2008, 26:696–697.
3. Yeh E, Slanetz P, Kopans DB, Rafferty E, Georgian-Smith D, Moy L, Halpern E, Moore R, Kuter I, Taghian A: Prospective comparison of mammography, sonography, and MRI in patients undergoing neoadjuvant chemotherapy for palpable breast cancer. *AJR Am J Roentgenol* 2005, 184:868–877.
4. Shin HJ, Kim HH, Ahn JH, Kim S-B, Jung KH, Gong G, Son BH, Ahn SH: Comparison of mammography, sonography, MRI and clinical examination in patients with locally advanced or inflammatory breast cancer who underwent neoadjuvant chemotherapy. *Br J Radiol* 2011, 84:612–620.
5. Marinovich ML, Houssami N, Macaskill P, Sardanelli F, Irwig L, Mamounas EP, von Minckwitz G, Brennan ME, Ciatto S: Meta-analysis of magnetic resonance imaging in detecting residual breast cancer after neoadjuvant therapy. *J Natl Cancer Inst* 2013, 105:321–333.
6. Loo CE, Teertstra HJ, Rodenhuis S, van de Vijver MJ, Hannemann J, Muller SH, Peeters M-JV, Gilhuijs KGA: Dynamic contrast-enhanced MRI for prediction of breast cancer response to neoadjuvant chemotherapy: initial results. *AJR Am J Roentgenol* 2008, 191:1331–1338.
7. Loo CE, Straver ME, Rodenhuis S, Muller SH, Wesseling J, Vrancken Peeters M-JTFD, Gilhuijs KGA: Magnetic resonance imaging response monitoring of breast cancer during neoadjuvant chemotherapy: relevance of breast cancer subtype. *J Clin Oncol Off J Am Soc Clin Oncol* 2011, 29:660–666.
8. von Minckwitz G, Untch M, Blohmer J-U, Costa SD, Eidtmann H, Fasching PA, Gerber B, Eiermann W, Hilfrich J, Huober J, Jackisch C, Kaufmann M, Konecny GE, Denkert C, Nekljudova V, Mehta K, Loibl S: Definition and impact of pathologic complete response on prognosis after neoadjuvant chemotherapy in various intrinsic breast cancer subtypes. *J Clin Oncol Off J Am Soc Clin Oncol* 2012, 30:1796–1804.
9. Untch M, Konecny GE, Paepke S, von Minckwitz G: Current and future role of neoadjuvant therapy for breast cancer. *Breast Edinb Scotl* 2014.
10. Rigter LS, Loo CE, Linn SC, Sonke GS, van Werkhoven E, Lips EH, Warnars HA, Doll PK, Bruining A, Mandjes IA, Vrancken Peeters MJ, Wesseling J, Gilhuijs KG, Rodenhuis S: Neoadjuvant chemotherapy adaptation and serial MRI response monitoring in ER-positive HER2-negative breast cancer. *Br J Cancer* 2013, 109:2965–2972.
11. von Minckwitz G, Blohmer JU, Costa SD, Denkert C, Eidtmann H, Eiermann W, Gerber B, Hanusch C, Hilfrich J, Huober J, Jackisch C, Kaufmann M, Kummel S, Paepke S, Schneeweiss A, Untch M, Zahm DM, Mehta K, Loibl S: Response-Guided Neoadjuvant Chemotherapy for Breast Cancer. *J Clin Oncol* 2013, 31:3623–3630.
12. Dakin H, Devlin N, Feng Y, Rice N, O'Neill P, Parkin D: The influence of cost-effectiveness and other factors on NICE decisions. *Health Econ* 2014, 24.
13. Thokala P, Dixon S, Jahn B: Resource Modelling: The Missing Piece of the HTA Jigsaw? *PharmacoEconomics* 2014.
14. OECD: *Health at a Glance 2013, OECD Indicators*. 2013.
15. Pacemakers and MRI Trends [http://www.medtronic.com/mrisurescan-us/mri_pacemakers_trends.html]
16. Coburn N, Przybysz R, Barbera L, Hodgson D, Sharir S, Laupacis A, Law C: CT, MRI and ultrasound scanning rates: Evaluation of cancer diagnosis, staging and surveillance in ontario. *J Surg Oncol* 2008, 98:490–499.
17. Wernli KJ, DeMartini WB, Ichikawa L, Lehman CD, Onega T, Kerlikowske K, Henderson LM, Geller BM, Hofmann M, Yankaskas BC: Patterns of Breast Magnetic Resonance Imaging Use in Community Practice. *JAMA Intern Med* 2014, 174:125.
18. Ries L, Eisner M, Kosary C: SEER cancer statistics review, 1973-1999. *Natl Cancer Inst* 2002.
19. Parise CA, Bauer KR, Brown MM, Caggiano V: Breast cancer subtypes as defined by the estrogen receptor (ER), progesterone receptor (PR), and the human epidermal growth factor receptor 2 (HER2) among women with invasive breast cancer in California, 1999-2004. *Breast J* 2009, 15:593–602.
20. The Netherlands Cancer Registry [<http://www.cijfersoverkanker.nl/>]

21. Early Breast Cancer Trialists' Collaborative Group: Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet* 2005, 365:1687–1717.
22. Common Terminology Criteria for Adverse Events (CTCAE) v4.0
23. Swain SM, Whaley FS, Ewer MS: Congestive heart failure in patients treated with doxorubicin: a retrospective analysis of three trials. *Cancer* 2003, 97:2869–2879.
24. Lee KS, Ro J, Nam B-H, Lee ES, Kwon Y, Kwon HS, Chung KW, Kang HS, Kim EA, Kim SW, Shin KH, Kim S-K: A randomized phase-III trial of docetaxel/capecitabine versus doxorubicin/cyclophosphamide as primary chemotherapy for patients with stage II/III breast cancer. *Breast Cancer Res Treat* 2008, 109:481–489.
25. Smith, DeCillis A, Anderson S, National Surgical Adjuvant Breast and Bowel Project Experience: Acute myeloid leukemia and myelodysplastic syndrome after doxorubicin-cyclophosphamide adjuvant therapy for operable breast cancer: the National Surgical Adjuvant Breast and Bowel Project Experience. *J Clin Oncol Off J Am Soc Clin Oncol* 2003, 21:1195–1204.
26. Briggs et al: *Decision Modelling for Health Economic Evaluation*. Oxford: Oxford university press; 2006.
27. Tsuji W, Teramukai S, Ueno M, Toi M, Inamoto T: Prognostic factors for survival after first recurrence in breast cancer: a retrospective analysis of 252 recurrent cases at a single institution. *Breast Cancer* 2014, 21:86–95.
28. Dutch National Center for Health Statistics [http://statline.cbs.nl/StatWeb/publication/?VW=T&DM=SLNL&PA=7052_95&D1=0-1,7,30-31,34,38,42,49,56,62-63,66,69-71,75,79&D2=0&D3=0&D4=a,!0-28&HD=080509-0829&HDR=G2,G1,G3&STB=T]
29. Rinaldi P, Costantini M, Belli P, Giuliani M, Bufi E, Fubelli R, Distefano D, Romani M, Bonomo L: Extra-mammary findings in breast MRI. *Eur Radiol* 2011, 21:2268–2276.
30. Retèl VP, Joore MA, Knauer M, Linn SC, Hauptmann M, van Harten WH: Cost-effectiveness of the 70-gene signature versus St. Gallen guidelines and Adjuvant Online for early breast cancer. *Eur J Cancer Oxf Engl* 1990 2010, 46:1382–1391.
31. Frederix, G: Disease specific methods for economic evaluations of breast cancer therapies. University Utrecht; 2013.
32. DBC product-finder for tariffs [<http://www.nza.nl/organisatie/>]
33. Farmacotherapeutisch Kompas [<http://www.fk.cvz.nl/>]
34. Dutch Health Care Insurance Board: Guide for research costs - Methods and standard cost prices for economic evaluations in healthcare.
35. CVZ [cvz.nl]
36. Wang G, Zhang Z, Ayala C, Wall HK, Fang J: Costs of heart failure-related hospitalizations in patients aged 18 to 64 years. *Am J Manag Care* 2010, 16:769–776.
37. Tina Shih Y-C, Xu Y, Elting LS: Costs of uncontrolled chemotherapy-induced nausea and vomiting among working-age cancer patients receiving highly or moderately emetogenic chemotherapy. *Cancer* 2007, 110:678–685.
38. Schilling MB, Parks C, Deeter RG: Costs and outcomes associated with hospitalized cancer patients with neutropenic complications: A retrospective study. *Exp Ther Med* 2011, 2:859–866.
39. Lidgren M, Wilking N, Jönsson B, Rehnberg C: Resource use and costs associated with different states of breast cancer. *Int J Technol Assess Health Care* 2007, 23:223–231.
40. XE currency Converter [<http://www.xe.com/>]
41. OECD (2013) [<http://stats.oecd.org/index.aspx?queryid=22519#>]
42. Lidgren M, Wilking N, Jönsson B, Rehnberg C: Health related quality of life in different states of breast cancer. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil* 2007, 16:1073–1081.
43. Nafees B, Stafford M, Gavriel S, Bhalla S, Watkins J: Health state utilities for non small cell lung cancer. *Health Qual Life Outcomes* 2008, 6:84.
44. Lloyd A, Nafees B, Narewska J, Dewilde S, Watkins J: Health state utilities for metastatic breast cancer. *Br J Cancer* 2006, 95:683–690.
45. Ford E, Adams J, Graves N: Development of an economic model to assess the cost-effectiveness of hawthorn extract as an adjunct treatment for heart failure in Australia. *BMJ Open* 2012, 2.

46. Coco A: The Cost-Effectiveness of Expanded Testing for Primary HIV Infection. *Ann Fam Med* 2005, 3:391–399.
47. Wertman R, Altun E, Martin DR, Mitchell DG, Leyendecker JR, O'Malley RB, Parsons DJ, Fuller ER, Semelka RC: Risk of nephrogenic systemic fibrosis: evaluation of gadolinium chelate contrast agents at four American universities. *Radiology* 2008, 248:799–806.
48. Singer OC, Sitzer M, Mesnil de Rochemont R du, Neumann-Haefelin T: Practical limitations of acute stroke MRI due to patient-related problems. *Neurology* 2004, 62:1848–1849.
49. Hand PJ, Wardlaw JM, Rowat AM, Haisma JA, Lindley RI, Dennis MS: Magnetic resonance brain imaging in patients with acute stroke: feasibility and patient related difficulties. *J Neurol Neurosurg Psychiatry* 2005, 76:1525–1527.
50. Sølling C, Ashkanian M, Hjort N, Gyldested C, Andersen G, Østergaard L: Feasibility and logistics of MRI before thrombolytic treatment. *Acta Neurol Scand* 2009, 120:143–149.
51. Dewey M: Claustrophobia preventing MR imaging of the breast. *Radiology* 2010, 256:328; author reply 328–329.
52. Coresh J, Astor BC, Greene T, Eknoyan G, Levey AS: Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis Off J Natl Kidney Found* 2003, 41:1–12.
53. Schaap K, Christopher-De Vries Y, Slottje P, Kromhout H: Inventory of MRI applications and workers exposed to MRI-related electromagnetic fields in the Netherlands. *Eur J Radiol* 2013, 82:2279–2285.
54. 'Handliding voor kosten onderzoek' 2010. College voor zorgverzekeringen, Rotterdam, 2010 - Guide for research costs - Methods and standard cost prices for economic evaluations in healthcare \ commissioned by the Health Care Insurance Board. [about:home]
55. National Public Health Atlas [<http://www.zorgatlas.nl/zorg/ziekenhuiszorg/>]
56. Radiological Society of the Netherlands [<https://www.radiologen.nl/12/1402/kerngegevens/general-information-radiological-society-of-the-netherlands.html>]

Supplementary material

Definitions of true-favourable, false-favourable, true-unfavourable and false-unfavourable used in our study.

Group of patients	Definition
True favourable	Patient that is classified as favourable at monitoring (criteria (Loo <i>et al</i> , 2011)), continues receiving NACT 1, and after 5 years of follow up is classified as favourable due to absence of relapse event
False favourable	Patient that is classified as favourable at monitoring (criteria (Loo <i>et al</i> , 2011)), continues receiving NACT 1, and after 5 years of follow up is classified as unfavourable due to presence of relapse event
True unfavourable	Patient that is unfavourable at monitoring (criteria (Loo <i>et al</i> , 2011)), switches to NACT 2, and after 5 years of follow up is classified as favourable due to absence of relapse event <i>(the underlying assumption is that the patient was not responding to NACT1 but did to NACT 2, thereby demonstrating that monitoring classified the patient properly)</i>
False unfavourable	Patient that is unfavourable at monitoring (criteria (Loo <i>et al</i> , 2011)), switches to NACT 2, and after 5 years of follow up is classified as unfavourable due to presence of relapse event <i>(the underlying assumption is that the patient was responding to NACT1 and did not to NACT 2, thereby demonstrating that monitoring classified the patient wrongly)*</i>

* Although we are aware that in the 'False favourable' group there could be patients irresponsive to both NACT 1 and 2, as the design of the RG-NACT does not allow distinguishing them, we had to make such an assumption.

Resource modeling outcomes, sources and calculations

Health services required at the country level		Current implementation (16 hospitals, 31 MRIs)	Full implementation (113 hospitals, 148 MRIs)	Source
Health services required at the country level				
No of MRIs scans performed		Calculations in table 2	No of stage II-III, ER-positive/HER2-negative breast cancers in the Netherlands	See table 2
No of patients scanned per MRI		'No of MRI scans performed'/31 MRIs ¹	'No of MRI scans performed'/148 MRIs ¹	See footnote 1
Fte MRI technologists required		Yearly hours required of MRI technologist to perform the 'No of MRIs scans performed' / Fully workable hours of an MRI technologist a year ²	idem	See footnote 2
Fte breast radiologists required		Yearly hours required of breast radiologist to perform the 'No of MRIs scans performed' / Fully workable hours of a breast radiologist a year ³	idem	See footnote 3
No of confirmations of incidental findings (using standard imaging)		Derived from the Markov model	idem	-
Health services required at the hospital level				
No of MRIs scans performed per hospital		'No of MRI scans performed' / 16 hospitals ⁴	'No of MRI scans performed' / 113 hospitals ⁵	See footnote 4 and 5
No of patients scanned per MRI per hospital		'No of MRI scans performed per hospital' / mean MRIs per hospital ¹	'No of MRI scans performed per hospital' / mean MRIs per hospital ¹	See footnote 1
Fte MRI technologists required per hospital		Yearly hours required of MRI technologist to perform the 'No of MRI scans performed per hospital' / Fully workable hours of an MRI technologist a year ²	idem	See footnote 2
Fte breast radiologists required per hospital		Yearly hours required of breast radiologist to perform the 'No of MRI scans performed per hospital' / Fully workable hours of a breast radiologist a year ³	idem	See footnote 3
Health outcomes gained at the country level				
No of relapses prevented		Derived from the Markov model	idem	-
No of breast cancer deaths prevented		Derived from the Markov model	idem	-

Health outcomes lost at the country level

No of excluded patients due to contraindications	Derived from the Markov model	idem	-
No of patients with NSF	'No of MRI scans performed' * <i>p</i> of NSF	idem	[52]
Fte MRI technologists with ATS	'Fte MRI technologists required' * <i>p</i> of ATS	idem	[53]
No of patients with CHF	Derived from the Markov model	idem	-
No of patients with long term AML/ADS	Derived from the Markov model	idem	-
No of patients with anxiety due to incidental findings	Derived from the Markov model	idem	-
No of patients with malignant incidental findings	'No of confirmations of incidental findings' * <i>p</i> malignant incidental findings ⁶	idem	[29]

Abbreviations: No= number; Fte= Full-time equivalent; MRI= magnetic resonance imaging; RG-NACT= response guided neoadjuvant chemotherapy; *p*= probability; NSF= nephrogenic systemic fibrosis; ATS= acute transient symptom; CHF= chronic heart failure; DSF=disease free survival; R=relapse; AML/ADS= myelodysplastic syndrome/acute myeloid leukaemia.

Note that when a calculation refers to another outcome of the table this is always the outcome within the same column i.e., within the same implementation rate.

Idem means calculated equal as the left cell, but adapted to the full implementation scenario figures.

¹ We search for this information in each hospital website. When this information was not available or unclear, we made use of literature [53] where the most frequent quantity of MRIs per type of hospital is presented (three for academic hospitals and one for general hospitals).

² Hours required of MRI technologists for the 'No of MRIs scans performed (per hospital)' in a year are calculated by assuming that a full scanning procedure requires 1 hour of MRI technologist. Employees were assumed to work 52 weeks/year, 5 days/week i.e., 260 days/year. Of these, 40 days would be vacation and sick days, resulting thus in 220 workable days/year. Assuming workers are employed for 8h/day this results in 1760 working hours/year. Yet workers need some time off during their working days i.e., breaks, assumed to be 20%. Thereby, a fully workable year is of 1408 hours.

³ Hours required of breast radiologist for the 'No of MRIs scans performed (per hospital)' in a year are calculated by assuming a mean of 6.8 minutes needed for a breast radiologist to interpret one MRI scan [57]. The workable hours a year of a breast radiologist were calculated exactly as explained in footnote 2.

⁴ Assuming its use in the biggest Dutch hospital network involved in RG-NACT (see 'resource modeling analysis' section).

⁵ Assuming its use in all Dutch hospitals (locations) with MRI expected to deliver cancer treatment (i.e., university, general and specialized hospitals) (see 'resource modeling analysis' section).

⁶ After confirming by ultrasound.

PART IV

IMAGING TECHNIQUES: SCREENING FOR DISTANT METASTASIS

CHAPTER 8

¹⁸F-FDG PET/CT for distant metastasis screening in stage II/III breast cancer patients: A cost-effectiveness analysis from a British, US and Dutch perspective

Anna Miquel-Cases*

Suzana C Teixeira*

Valesca P Retèl

Lotte MG Steuten

Renato A Valdés Olmos

Emiel JT Rutgers#

Wim H van Harten#

* First shared authorship, # Last shared authorship

Submitted for publication

Abstract

Purpose: ^{18}F -FDG PET/CT (PET/CT) is more accurate than conventional imaging (CI) in detecting distant metastasis (DM) in primary stage II/III breast cancer patients. As PET/CT comes at high costs, we estimated its added value from a perspective of the United Kingdom (UK), the United States (US) and the Netherlands (NL).

Patients and methods: A Markov model compared costs, life years (LYs), quality-adjusted LYs (QALYs), and cost-effectiveness (incremental net monetary benefit, iNMB) of DM screening with PET/CT vs. CI (according to European and US standards) from a hospital perspective over a 5-year time horizon in four breast cancer subtypes (classified by ER and HER2 status). Imaging performance, systemic, and local treatment data stemmed from the Netherlands Cancer Institute. Epidemiological, survival and utility data were derived from recent literature. Costs (2013) derived from national tariffs (UK/NL)/Centers for Medicaid and Medicare Services (US). One-way sensitivity analysis identified the ceiling PET/CT costs to achieve cost-effectiveness per country.

Results: PET/CT was more sensitive (92% vs. 13%) and specific (98% vs. 94%) than CI. Gains in LYs (0.007 ± 0.0001) and QALYs (0.002 ± 0.0001) were similar across subtypes. Largest cost savings were in ER-positive/HER2-negative patients (incremental costs NL/ UK/ US = €447/ €1100/ -€1461) and least in ER-positive/HER2-positive (€1739/ €4382/ €2662). PET/CT was expected cost-effective with high certainty in HER2-negative patients of the US (iNMB range = €1089-€1571, probability of cost-effectiveness range = 83-97%). Ceiling PET/CT costs for ER-positive/HER2-negative and ER-negative/HER2-positive patients were \$1000(US)/ €600(NL)/ £500(UK). For the remaining subtypes, this was conditional to additional cost-reductions in Trastuzumab (US), or Trastuzumab plus Paclitaxel (NL/UK).

Conclusions: PET/CT adds value if it reduces costly palliative treatment. So far, this is only achieved with in the HER2-negative subtypes of the US. Reductions in PET/CT and palliative treatment costs are warranted to attain cost-effectiveness in the NL and UK.

Introduction

Preoperative systemic treatment (PST) is becoming treatment of first choice in breast cancer, as it facilitates breast conservation and has positive influence on survival [1]. Breast cancer patients receiving PST require prior distant metastases (DM) screening. Currently, this is performed by bone scan, plus liver sonography and chest X-ray [2,3] in Europe, and by bone scan, plus liver sonography and CT thorax/abdomen in the US. Recently, positron emission tomography with integrated low-dose computed tomography (PET/CT) using fluorine-18 fluoro-deoxy-glucose (^{18}F -FDG) has shown to be of additional value to detect DM [4–8]. In a series of 167 patients recruited in a comprehensive cancer center (Netherlands Cancer Institute–Antoni van Leeuwenhoek Hospital; NKI) PET/CT sensitivity was found to be of 100% compared to that of 57.9% for conventional imaging (CI)[6]. These findings lead to new recommendations in the ‘Dutch guidelines for breast cancer diagnostics and treatment’ stating that “ ^{18}F -FDG-PET/CT can replace conventional staging methods for DM screening and is therefore advised for stage III breast cancer. Furthermore, it can be considered in stage II primary breast cancer”.

PET/CT is also able to better detect metastatic lesions in an earlier stage than CI. If these lesions are limited in number (max 3 or 5), so-called “oligometastatic lesions”[9], the patient can be treated with curative intent [10–12]. The clinical adoption of PET/CT is thus expected to improve survival outcomes in breast cancer patients. However, PET/CT comes at significant additional cost. Its actual implementation will depend on the extent to which these costs are justified by the incremental health benefits achieved, as well as by the potential cost savings attained in other parts of the patient pathway.

To estimate the added value of implementing PET/CT for DM screening in stage II/III breast cancer, we conducted a model-based cost-effectiveness analysis (CEA) using patient data from the NKI. As PET/CT is potentially applicable in a variety of countries, we conducted this analysis from a perspective of the Netherlands (NL), the United Kingdom (UK) and the United States (US). Furthermore, we explored the ceiling PET/CT costs to achieve cost-effectiveness in each country.

Patients and methods

We developed a Markov model to compare health economic consequences of DM screening by ‘full body ^{18}F -FDG PET/CT’ or by ‘CI’ in four cohorts of stage II-III breast cancer (ER-negative/HER2-positive, ER-positive/HER2-positive, ER-negative/HER2-negative, and ER-positive/HER2-negative) scheduled for PST. CI was modelled according to European and US standards. For technical details of PET/CT and CI see supplementary material. The CEA was performed from a hospital perspective

of the NL, the UK and the US (annual discount rates per country were of 4% for costs and 1.5% for effects [13]; 3.5% for both[14]; 3% for both respectively)[15] over a 5-years' time horizon.

Imaging performance, systemic and local treatments, and patient baseline characteristics (stage II/III breast cancer, post-menopausal status, 50 years old) were derived from patients treated at the NKI from 2007 to 2013. Epidemiological, survival and utility data were derived from recent literature or expert assumptions. Costs (2013) were obtained from national tariffs (UK and NL), and the Centres for Medicaid and Medicare Services (US).

Markov model

The Markov model has eight mutually exclusive health-states reflecting the natural history of the disease (Figure 1). Patients entered the model classified as true-positive (TP), false-positive (FP), true-negative (TN) or false-negative (FN) with respect to the presence of DM at imaging, based on the PET/CT or the CI strategy. DM lesions were grouped into single lung, single bone, single liver or multiple. Patients were classified as positive following a tumour-positive biopsy, or if no biopsy was taken, by confirmation on another imaging modality. Patients were classified as negative based on disease free survival at 6 months after the PET/CT was made. Specific definitions for TP, FP, TN and FN are shown in table 1.

Transition of a patient from one health-state to another was defined in yearly cycles for a time horizon of 5-years. A description of the course that patients followed in the model as well as the assigned health-state costs and utilities are presented in the supplementary material.

Figure 1: Decision tree and Markov model of distant metastasis screening with PET/CT vs CI in four subtypes of stage II/III breast cancer patients. Two strategies are presented: DM screening with PET/CT vs. DM screening with CI (chest X-ray, liver sonography plus bone scan (UK/NL) and CT-thorax-abdomen plus bone scan (US)). In the first year of the model, simulated by the decision tree, all patients incur the costs of DM screening and primary breast cancer treatment. Furthermore, in the case of true- and false- positive patients, they also incur the additional cost of biopsy, plus DM treatment (true positives) and imaging (false positives), and in the case of false- negative patients, additional costs of biopsy plus imaging and DM treatment. The quality-of-life of patients in this first year will mainly be determined by the presence or absence of DM. The last square of the tree represent the health-state of Markov model were patients enter in the 1st year, either stable or DM health-state. The Markov model simulates the disease progression of the patients, were costs and quality of life are accumulated at the time horizon of 5-years. Abbreviations: DM= distant metastases; Tx=treatment; L=local, PBC= primary breast cancer treatment.

Table 1: Definitions, survival, costs and quality of life associated assumptions regarding true-positive, false-positive, true-negative and false-negative patients.

	Definition	Survival	Costs	Quality of life
TP	Imaging reveals metastasis and is confirmed by biopsy or additional imaging	++ (early detection DM)	+++ (biopsy and DM _{Tx})	++ (Presence DM and Palliative _{Tx})
FP	Imaging reveals metastasis but the presence of metastatic disease is not confirmed by biopsy or additional imaging	+++ (no DM)	++ (biopsy and confirmation scans)	+++ (PBC _{Tx})
TN	Imaging reveals no metastasis and this is confirmed by "6 months follow-up"	+++ (no DM)	+ (none)	+++ (PBC _{Tx})
FN*	Imaging reveals no metastasis but metastatic disease is present at "6 months follow-up"	+ (late detection of DM)	++++ (biopsy, confirmation scans and DM _{Tx})	+ (painful DM and Palliative _{Tx})

Abbreviations: TP= true-positive; FP= false positive; TN= true negative; FN= false negative; DM= distant metastasis; PBC= primary breast cancer treatment; Tx=treatment

*As all patients in our database were scanned by CI and PET/CT, when calculating the performance of CI the following had to be assumed: patients that were negative under the conventional strategy but that were treated as positive at the discretion of the physician after PET/CT discovered DM were included in the false negative (FN) group. These patients were assigned the same costs, utilities and transition probabilities as the remaining FNs.

Model input data

Clinical database

We retrospectively collected data from 545 stage II/III breast cancer patients who underwent CI and PET/CT to detect distant dissemination before start of PST, in the NKI from 2007 to 2013. From this database, we derived imaging performance (PET/CT and CI) and data on primary breast cancer treatment (PST, breast surgery, adjuvant chemotherapy and breast radiotherapy). Performance data was obtained from 413 patients (supplementary table 2). Data on primary breast cancer treatment came from 157 patients treated in the year 2013 (supplementary table 3). As this was the most recent data in our database, it was expected to most adequately represent current treatment.

Pre-treatment core biopsies of the primary tumor were classified according to the conventional criteria of the World Health Organization [14] to determine breast cancer subtypes. After pathology assessment, but prior to PST initiation, patients were scanned with CI and PET/CT.

The reports of PET/CT and CI were discussed in a multi-disciplinary meeting where the nuclear physician and radiologist gave their advice and discussed whether further investigations were desirable.

The treatment for patients with DM was assumed, as only nine patients in our dataset developed a metastasis. A patient with a single metastasis received local treatment consisting of surgery for metastases in liver and lung, and radiotherapy for lesions in the bone. Furthermore, patients with bone DM were treated with Zometa (bisphosphonate). Multi organ metastasis were assumed to always include a bone lesion, and were treated with one line of systemic treatment, (according to Dutch guidelines)[17]. If DM lesions were detected prior to start of treatment, patients received Anastrozole plus Zometa for 5-years (ER+/HER2-), Trastuzumab plus Paclitaxel until death (ER+/HER2+, ER-/HER2+) or Paclitaxel monotherapy until death (ER-/HER2-). If multi DM lesions were detected during treatment, regimens were Capecitabine (ER+/HER-, ER+/HER2+, ER-/HER2-) and Trastuzumab plus Paclitaxel (ER-/HER2+). Systemic treatment dosages are presented in supplementary table 1.

Data derived from literature

Epidemiological data (i.e., common types and sites of metastasis per subtype, and frequencies of chemotherapy-related toxicities) and survival data (i.e., per site of metastasis) were derived from recent literature. Epidemiology data came from studies with similar subtype and DM sites classification as our model. Frequencies on the types of DM (multi or single) were derived from a Finish cohort study on 2.032 invasive operable breast cancer [18] with similar frequencies as our database (22% multiple vs. 78% single). Frequencies on the DM sites (lung, liver, bone and multiple) came from a cohort of 531 U.S citizens with distant metastatic disease from breast cancer[19]. Both type of frequencies were reported similar in other recent literature [20–23]. Short-term chemotherapy-related adverse-events included vomiting, neutropenia, hand-food-syndrome, thrombocytopenia, mucositis and cardio-toxicities (symptomatic, class II-IV from the NYHA[24]). These were included in the model if prevalence $\geq 10\%$ and classified as related to anthracyclines, taxanes, anthracyclines plus taxanes, anthracyclines plus Trastuzumab and paclitaxel plus Trastuzumab (supplementary table 4).

Data on breast cancer mortality came from a Norwegian study on the survival of 304 metastasized breast cancers [20]. Survival was assigned based on first site of metastasis: bone (bone DM), visceral (liver and lung DM) or 'bone plus visceral' (multi organ DM). Survival rates in years 4 and 5 were assumed equal for patients with 'visceral' and 'bone plus visceral' lesions. This was decided upon the low patient numbers in these years, generating unexpectedly different survival rates between these groups. In FN patients, the probability of breast cancer death was simulated

higher than in FP, as metastases are detected with a delay and there is a lower likelihood of cure. The applied factor was estimated from our database, where a 1.8 higher probability of breast cancer death was observed in FNs vs FPs. This was corroborated by an experienced surgical oncologist. The probability of dying from a non-breast cancer related event was derived from the Dutch cancer registry [25].

Costs of diagnostic imaging, biopsy (assumed to be ultrasound guided core biopsy), surgery (breast or metastatic site), radiotherapy and follow-up were derived from Dutch reference tariffs [26], NHS reference costs [27] and the centres for Medicare & Medicaid services (CMS)[28] and literature [29–42]. Costs of systemic treatments and of the treatment of adverse events were derived from Dutch published literature [43–47], except vomiting where we used data from Canada [48] due to the lack of a Dutch estimator, NHS reference costs [27,41,49–56] and average selling prices from CMS [28] and literature [57–63] for the US. All costs were inflated to 2013 values using the Consumer Price Index [64] and transformed to Euros [65].

Utility estimates were obtained from the review of Peasgood et al [66] or from the CEA registry [67]. When multiple utilities were identified, we prioritized those reflecting the patient's perspective using the EQ-5D profile. Biopsy was assumed 100% accurate and that induces no QALY decrement.

Supplementary table 4 summarizes all model parameters and its sources.

Model outcomes

Outcomes were the 5-years' incremental effects (FN and FP prevented, TP and TN gained, and life years (LY) and quality-adjusted-life-years (QALYs) gained), incremental costs (2013, reported in country-specific currencies and euros) and incremental net monetary benefit ratio (iNMB)[68] of DM screening with PET/CT minus DM screening with CI. If $iNMB > 0$ PET/CT was considered cost-effective.

Cost effectiveness analysis

A probabilistic sensitivity analysis (PSA) with 10.000 Monte Carlo simulations was undertaken for each breast cancer subtype, using the costs of each country (NL, UK and US). Each model parameter was assigned a probability distribution: Dirichlet for performance, beta for effectiveness and utilities, and gamma for costs parameters (supplementary table 4). By randomly drawing a value for each input parameter from the assigned distribution, the PSA quantifies the joint decision uncertainty in model outcomes. This is summarized in cost-effectiveness acceptability curves (CEACs)[69]. They represent the probability that PET/CT is cost-effective given a certain threshold of willingness to pay for an additional QALY. The iNMB (i.e., cost-effectiveness) was

determined using the prevailing threshold for cost-effectiveness in each country ($\lambda = \text{€}80.000/\text{QALY}$ in the Netherlands[13], $\text{£}30.000/\text{QALY}$ in the UK⁷¹ and $\text{\$}50.000/\text{QALY}$ in the US). CEACs were presented per country and subtype.

One-way sensitivity analysis

One-way sensitivity analysis (SA) was conducted to all model parameters to determine to which parameters each model was most sensitive. This was performed from a US and NL perspective; we did not use the UK perspective because this was expected to behave similar to the NL model. Furthermore, we determined the upper margin of PET/CT costs that warrant the PET/CT strategy cost-effective per country.

Results

Sensitivity and specificity were 13% and 92% for CI, and 94% and 98% for PET/CT respectively. The PET/CT strategy prevented FNs and FPs by 0.89 and 0.65 times respectively, while increasing TN and TP by 1.04 and 8.3 times respectively. Subtypes with higher probability to develop bone DM (ER-positive/HER2-positive and ER-positive/HER2-negative) had higher LYs, as these lead to longer short-term survival as compared to visceral DMs. Subtypes with high frequency of multiple DMs (ER-negative/HER2-negative and ER-positive/HER2-positive) had lower utility weights resulting in lower QALYs. This led to 0.007 ± 0.0001 LYs and 0.002 ± 0.0001 QALYs gained, depending on tumour subtype.

An increase in costs by the PET/CT strategy was consistently seen in the UK (range $\text{€}1100/\text{€}4382$) and in the NL ($\text{€}447/\text{€}1739$), but not in the US ($\text{€}-1461/\text{€}2662$). In the UK and the NL, largest cost savings were seen in ER-positive/HER2-negative ($\text{€}1100/\text{€}447$), followed by ER-negative/HER2-positive ($\text{€}1319/\text{€}582$), ER-negative/HER2-negative ($\text{€}2629/\text{€}1050$), and ER-positive/HER2-positive ($\text{€}4382/\text{€}1739$). In the US, largest savings were in ER-positive/HER2-negative ($\text{€}-1461$), followed by ER-negative/HER2-negative ($\text{€}-991$), ER-negative/HER2-positive ($\text{€}133$) and ER-positive/HER2-positive ($\text{€}2662$).

The iNMBs were highest in the US (range $\text{€}-2517/\text{€}1571$), compared to the NL ($\text{€}-259/\text{€}1560$) and the UK ($\text{€}-4289/\text{€}1003$), and following the opposite order of incremental costs. In the US, PET/CT became cost-effective in the subtypes that had cost savings. The probability that PET/CT was cost-effective was low in the UK (range 0/22%) and the NL (4/31%), dependent on subtype. In the US, this was high for the ER-positive/HER2-negative (97%) and ER-negative/HER2-negative subtypes (83%), but below 50% for the remaining subtypes. Cost-effectiveness results are summarized in table 2 and CEACs are presented in Figure 2.

Table 2: Results from the cost-effectiveness analysis

	The US				The Netherlands				The UK			
	Δ Costs	Δ LYs	Δ QALYs	iNMB (pCE)	Δ Costs	Δ LYs	Δ QALYs	iNMB (pCE)	Δ Costs	Δ LYs	Δ QALYs	iNMB (pCE)
ER-positive/ HER2-negative	-€1461 -\$1606	0,007	0,002	€1571 \$1727 (97%)	€447	0,007	0,002	-€259 (25%)	€1100 £780	0,007	0,002	-€1003 -£712 (3%)
ER-negative/ HER2-positive	€133 \$146	0,007	0,002	-€18 -\$20 (48%)	€582	0,007	0,002	-€384 (31%)	€1319 £936	0,007	0,002	-€1215 -£862 (22%)
ER-negative/ HER2-negative	-€991 -\$1090	0,007	0,002	€1089 \$1197 (83%)	€1050	0,007	0,002	-€883 (10%)	€2629 £1864	0,007	0,002	-€2542 -£1803 (5%)
ER-positive/ HER2-positive	€2662 \$2822	0,007	0,002	-€2517 -\$2766 (11%)	€1739	0,007	0,002	-€1560 (4%)	€4382 £3107	0,007	0,002	-€4289 -£3042 (0%)

Abbreviations: LY= life years; QALY= quality adjusted life year; iNMB= incremental monetary benefit; pCE: probability of cost-effectiveness. 1 pound = 1.41 euros; 1 dollar= 0.91 euros

Figure 2: Cost-effectiveness acceptability curves per subtype and country (10,000 simulations). In each figure, the bottom curves represent the probability that the PET-CT strategy is more cost-effective than conventional imaging (CI) (INMB > 0), at a specific willingness to pay threshold, different per country (marked with a vertical line).

Results from one-way SA to all model parameters showed that DM screening costs, palliative treatment costs and imaging performance drove cost-effectiveness. These are presented in the supplementary material. The upper margin of PET/CT costs to warrant the PET/CT strategy cost-effective were \$1000 (US), €600 (NL) and £500 (UK) in ER-positive/HER2-negative and ER-negative/HER2-positive patients (table 3). Even at these cost levels, PET/CT did not become cost-effective for ER-positive/HER2-positive and ER-negative/HER2-negative patients of the NL and the UK, and ER-positive/HER2-positive patients of the US. To achieve cost-effectiveness in these groups the costs of Trastuzumab and Paclitaxel would have to be lowered (potential scenarios for the treatment costs are presented in supplementary table 5).

Table 3: Upper margin of cost of PET/CT to reach cost-effectiveness per subtype and country

	ER-positive/ HER2-negative	ER-negative/ HER2-positive	ER-negative/ HER2-negative	ER-positive/ HER2-positive
US	<\$2900	<\$1000	<\$2300	Conditional on cost reduction in palliative regimen costs
NL	<€700	<€600	Conditional on cost reduction in palliative regimen costs	Conditional on cost reduction in palliative regimen costs
UK	<£600	<£500	Conditional on cost reduction in palliative regimen costs	Conditional on cost reduction in palliative regimen costs

Palliative treatment in ER-positive/HER2-positive is Trastuzumab plus Paclitaxel, and ER-negative/HER2-negative, Paclitaxel monotherapy. Treatment costs are yearly costs given with metastatic dosages, as detailed in supplementary table 1.

Discussion

Our study reveals that PET/CT outperforms CI in detecting DMs in stage II-III breast cancer patients. However, this comes at additional costs of imaging and palliative treatment. So far these are only outweighed by health benefits in the US. Cost-effectiveness in the UK and the NL could be achieved by lowering the costs of PET/CT as well as the costs of specific treatments given as palliative treatment.

The 8.3-time increase in early and 0.89-time decrease in late detection of DMs with the PET/CT strategy resulted in LYs and QALY gains in all subtypes and countries analysed (equal between countries, and similar between subtypes). The observed health gains were however modest, as can be expected for the limited survival of metastatic patients (0.007 LYs and 0.002 QALYs).

Incremental costs were mainly driven by the costs of DM screening; as these are incurred in the total breast cancer population under study. This trend was noticed in the incremental costs per country. The country with the highest incremental DM screening costs (the UK) had the highest

overall incremental cost per patient. A secondary driver of incremental costs were palliative treatment costs. Their influence was visible when costs were extremely high, as is the case for the systemic treatment of HER2-positive subtypes treated in the US; PET/CT became cost-ineffective despite having the lowest increase in DM screening cost.

The main driver of incremental cost differences between subtypes was palliative treatment costs, received by TP and FN patients. As by using PET/CT the number of TPs increased (by 8.3 times) and the number of FPs decreased (by 1.04 times), patients who needed the most costly TP treatment (Trastuzumab plus Paclitaxel) but a proportionally cheaper FP treatment (capecitabine), i.e., ER-positive/HER2-positive patients, had the highest incremental costs in all countries. In the other end of the spectrum, ER-positive/HER2-negative patients, who had the cheapest TP treatment (Anastrozole plus Zometa) and a proportionally expensive treatment for FPs (capecitabine), had the least incremental costs.

As health gains were similar across countries and subtypes, but costs differed, the latter drove the cost-effectiveness results. Our model revealed that only in the subtypes with cost savings, as is the case of HER2-negative subtypes treated in the US, cost-effectiveness was achieved with high probabilities. In the remaining of cases probability of cost-effectiveness remained below 50% (Figure 2).

The main driver of cost-effectiveness was imaging performance, followed by DM screening costs or palliative treatment costs, depending on subtype. These are the aspects one should focus in order to determine courses of action to warrant the PET/CT strategy more cost-effective. However, as PET/CT performance is already superior to that of CI our suggestion would be to concentrate on the other two drivers. While for ER-positive/HER2-negative and ER-negative/HER2-positive patients determining an upper margin cost for PET/CT is sufficient, in the remaining subtypes this should go along with additional cost-reductions in Trastuzumab (US), or Trastuzumab plus Paclitaxel (NL/UK). Costs reductions in palliative treatment costs could be achieved by increasing the detection of “oligometastatic” metastasis, as these patients can be treated with curative intent.

The cost-effectiveness of DM screening with PET/CT in breast cancer has previously been reported from a Dutch perspective. Unfortunately, this study reported incremental costs per saved biopsy [71], and can therefore not be compared to our cost/QALY estimates.

One of our study limitations is that biopsy performance was assumed perfect, yet false-negative rates reported in literature (0-9% [72]) make this a fairly feasible assumption. Moreover, the factor applied to lower FNs survival, warrants further research, as despite being derived from our

clinical database and confirmed by an experienced surgeon, it is uncertain and a key driver of cost-effectiveness. Yet at the time of study, this was the best available source. Although it is not well known whether 6-months of follow-up is sufficient to capture missed DM at screening, this time frame was chosen in accordance with previously reported results of our institute [6]. Finally, we assumed primary breast cancer treatment in all countries to be equal of that of the NKI, as we expect treatment guidelines to be similar.

Our study demonstrates that PET/CT adds value in detecting DM in breast cancer if it detects TP patients treated with low-priced palliative treatment and prevents FNs with low-prognosis i.e., if it reduces costly palliative treatment. So far, this is only achieved in the HER2-negative subtypes treated in the US. To achieve cost-effectiveness in the NL and the UK, reductions in PET/CT and palliative treatment costs are warranted. A way forward to decrease palliative treatment costs is by increasing the detection of 'oligometastatic lesions' treated with local procedures and curative intent.

Acknowledgements

We would like to thank prof. dr. Rodenhuis for his help on the systemic treatment assumptions used in our model.

References

- [1] Mieog JSD, van der Hage JA, van de Velde CJH. Preoperative chemotherapy for women with operable breast cancer. *Cochrane Database Syst Rev* 2007;CD005002. doi:10.1002/14651858.CD005002.pub2.
- [2] Fuster D, Duch J, Paredes P, Velasco M, Muñoz M, Santamaría G, et al. Preoperative staging of large primary breast cancer with [18F]fluorodeoxyglucose positron emission tomography/computed tomography compared with conventional imaging procedures. *J Clin Oncol Off J Am Soc Clin Oncol* 2008;26:4746–51. doi:10.1200/JCO.2008.17.1496.
- [3] Bovelli D, Plataniotis G, Roila F, On behalf of the ESMO Guidelines Working Group. Cardiotoxicity of chemotherapeutic agents and radiotherapy-related heart disease: ESMO Clinical Practice Guidelines. *Ann Oncol* 2010;21:v277–82. doi:10.1093/annonc/mdq200.
- [4] Fuster D, Duch J, Paredes P, Velasco M, Muñoz M, Santamaría G, et al. Preoperative staging of large primary breast cancer with [18F]fluorodeoxyglucose positron emission tomography/computed tomography compared with conventional imaging procedures. *J Clin Oncol Off J Am Soc Clin Oncol* 2008;26:4746–51. doi:10.1200/JCO.2008.17.1496.
- [5] Riegger C, Herrmann J, Nagarajah J, Hecktor J, Kuemmel S, Otterbach F, et al. Whole-body FDG PET/CT is more accurate than conventional imaging for staging primary breast cancer patients. *Eur J Nucl Med Mol Imaging* 2012;39:852–63. doi:10.1007/s00259-012-2077-0.
- [6] Koolen BB, Vrancken Peeters M-JTFD, Aukema TS, Vogel W V, Oldenburg HSA, van der Hage JA, et al. 18F-FDG PET/CT as a staging procedure in primary stage II and III breast cancer: comparison with conventional imaging techniques. *Breast Cancer Res Treat* 2012;131:117–26. doi:10.1007/s10549-011-1767-9.
- [7] Groheux D, Giacchetti S, Espié M, Vercellino L, Hamy A-S, Delord M, et al. The yield of 18F-FDG PET/CT in patients with clinical stage IIA, IIB, or IIIA breast cancer: a prospective study. *J Nucl Med Off Publ Soc Nucl Med* 2011;52:1526–34. doi:10.2967/jnumed.111.093864.
- [8] Groheux D, Giacchetti S, Delord M, Hindié E, Vercellino L, Cuvier C, et al. 18F-FDG PET/CT in staging patients with locally advanced or inflammatory breast cancer: comparison to conventional staging. *J Nucl Med Off Publ Soc Nucl Med* 2013;54:5–11. doi:10.2967/jnumed.112.106864.
- [9] Di Lascio S, Pagani O. Oligometastatic breast cancer: A shift from palliative to potentially curative treatment? *Breast Care* 2014;9:7–14. doi:10.1159/000358750.
- [10] Singh D, Chen Y, Hare MZ, Usuki KY, Zhang H, Lundquist T, et al. Local control rates with five-fraction stereotactic body radiotherapy for oligometastatic cancer to the lung. *J Thorac Dis* 2014;6:369–74. doi:10.3978/j.issn.2072-1439.2013.12.03.
- [11] Niibe Y, Kuranami M, Matsunaga K, Takaya M, Kakita S, Hara T, et al. Value of high-dose radiation therapy for isolated osseous metastasis in breast cancer in terms of oligo-recurrence. *Anticancer Res* 2008;28:3929–31.
- [12] Fumagalli I, Bibault J-E, Dewas S, Kramar A, Mirabel X, Prevost B, et al. A single-institution study of stereotactic body radiotherapy for patients with unresectable visceral pulmonary or hepatic oligometastases. *Radiat Oncol* 2012;7:164. doi:10.1186/1748-717X-7-164.
- [13] Dutch National Health Care Institute. CVZ n.d. cvz.nl.
- [14] National Institute for Health and Clinical Excellence. Guide to the Methods of Technology Appraisal. London: 2008.
- [15] Academy of managed care pharmacy (AMCP): Format submission of Clinical and Economic evidence of Pharmaceuticals in Support of Formulary Consideration. 2012.
- [16] Tavassoli F, Weltgesundheitsorganisation, International Agency for Research on Cancer, editors. Pathology and genetics of tumours of the Breast and Female Genital Organs. Reprint. Lyon: IARC Press; 2003.
- [17] Integraal Kankercentrum Nederland (IKNL). Breast Cancer Guideline, NABON 2012. n.d.
- [18] Sihto H, Lundin J, Lundin M, Lehtimäki T, Ristimäki A, Holli K, et al. Breast cancer biological subtypes and protein expression predict for the preferential distant metastasis sites: a nationwide cohort study. *Breast Cancer Res BCR* 2011;13:R87. doi:10.1186/bcr2944.

- [19] Soni A, Ren Z, Hameed O, Chanda D, Morgan CJ, Siegal GP, et al. Breast cancer subtypes predispose the site of distant metastases. *Am J Clin Pathol* 2015;143:471–8. doi:10.1309/AJCPYO5FSV3UPEXS.
- [20] Kast K, Link T, Friedrich K, Petzold A, Niedostatek A, Schoffer O, et al. Impact of breast cancer subtypes and patterns of metastasis on outcome. *Breast Cancer Res Treat* 2015;150:621–9. doi:10.1007/s10549-015-3341-3.
- [21] Lobbezoo DJA, van Kampen RJW, Voogd AC, Dercksen MW, van den Berkmortel F, Smilde TJ, et al. Prognosis of metastatic breast cancer subtypes: the hormone receptor/HER2-positive subtype is associated with the most favorable outcome. *Breast Cancer Res Treat* 2013;141:507–14. doi:10.1007/s10549-013-2711-y.
- [22] Alqaisi A, Chen L, Romond E, Chambers M, Stevens M, Pasley G, et al. Impact of estrogen receptor (ER) and human epidermal growth factor receptor-2 (HER2) co-expression on breast cancer disease characteristics: implications for tumor biology and research. *Breast Cancer Res Treat* 2014;148:437–44. doi:10.1007/s10549-014-3145-x.
- [23] Kennecke H, Yerushalmi R, Woods R, Cheang MCU, Voduc D, Speers CH, et al. Metastatic Behavior of Breast Cancer Subtypes. *J Clin Oncol* 2010;28:3271–7. doi:10.1200/JCO.2009.25.9820.
- [24] William E. Chavey, Caroline S. Blaum, Barry E. Bleske, Richard van Harrison, Sean Kesterson, John M. Nicklas. Guideline for the Management of Heart Failure Caused by Systolic Dysfunction: Part II. Treatment. *Am Fam Physician* 2001;104:5–55.
- [25] iKNL (Integraal Kankercentrum Nederland). Nederlandse Kankerregistratie n.d. <http://www.cijfersoverkanker.nl/over-de-registratie-12.html>.
- [26] Dutch Healthcare Authority. DBC product-finder for tariffs n.d. <http://www.nza.nl/organisatie/>.
- [27] NHS reference costs. 2013.
- [28] Centers for Medicare & Medicaid Services. n.d.
- [29] Melnikow J, Tancredi DJ, Yang Z, Ritley D, Jiang Y, Slee C, et al. Program-Specific Cost-Effectiveness Analysis: Breast Cancer Screening Policies for a Safety-Net Program. *Value Health* 2013;16:932–41. doi:10.1016/j.jval.2013.06.013.
- [30] Barlow WE, Taplin SH, Yoshida CK, Buist DS, Seger D, Brown M. Cost comparison of mastectomy versus breast-conserving therapy for early-stage breast cancer. *J Natl Cancer Inst* 2001;93:447–55.
- [31] Landman MP, Feurer ID, Pinson CW, Moore DE. Which is more cost-effective under the MELD system: primary liver transplantation, or salvage transplantation after hepatic resection or after loco-regional therapy for hepatocellular carcinoma within Milan criteria? *HPB* 2011;13:783–91. doi:10.1111/j.1477-2574.2011.00355.x.
- [32] Puri V, Crabtree TD, Kymes S, Gregory M, Bell J, Bradley JD, et al. A comparison of surgical intervention and stereotactic body radiation therapy for stage I lung cancer in high-risk patients: A decision analysis. *J Thorac Cardiovasc Surg* 2012;143:428–36. doi:10.1016/j.jtcvs.2011.10.078.
- [33] Reed SD, Li Y, Anstrom KJ, Schulman KA. Cost effectiveness of ixabepilone plus capecitabine for metastatic breast cancer progressing after anthracycline and taxane treatment. *J Clin Oncol Off J Am Soc Clin Oncol* 2009;27:2185–91. doi:10.1200/JCO.2008.19.6352.
- [34] Alvarado MD, Mohan AJ, Esserman LJ, Park CC, Harrison BL, Howe RJ, et al. Cost-Effectiveness Analysis of Intraoperative Radiation Therapy for Early-Stage Breast Cancer. *Ann Surg Oncol* 2013;20:2873–80. doi:10.1245/s10434-013-2997-3.
- [35] Konski A. Radiotherapy is a cost-effective palliative treatment for patients with bone metastasis from prostate cancer. *Int J Radiat Oncol* 2004;60:1373–8. doi:10.1016/j.ijrobp.2004.05.053.
- [36] Kimman ML, Dirksen CD, Voogd AC, Falger P, Gijzen BCM, Thuring M, et al. Economic evaluation of four follow-up strategies after curative treatment for breast cancer: Results of an RCT. *Eur J Cancer* 2011;47:1175–85. doi:10.1016/j.ejca.2010.12.017.
- [37] Auguste P, Barton P, Hyde C, Roberts T. An economic evaluation of positron emission tomography (PET) and positron emission tomography/computed tomography (PET/CT) for the diagnosis of breast cancer recurrence. *Health Technol Assess* 2011;15. doi:10.3310/hta15180.
- [38] Hummel S, Simpson E, Hemingway P, Stevenson M, Rees A. Intensity-modulated radiotherapy for the treatment of prostate cancer: a systematic review and economic evaluation. *Health Technol Assess* 2010;14. doi:10.3310/hta14470.

- [39] Meng Y, Ward S, Cooper K, Harnan S, Wyld L. Cost-effectiveness of MRI and PET imaging for the evaluation of axillary lymph node metastases in early stage breast cancer. *Eur J Surg Oncol EJSO* 2011;37:40–6. doi:10.1016/j.ejso.2010.10.001.
- [40] Hamada S, Hinotsu S, Ishiguro H, Toi M, Kawakami K. Cross-National Comparison of Medical Costs Shared by Payers and Patients: A Study of Postmenopausal Women with Early-Stage Breast Cancer Based on Assumption Case Scenarios and Reimbursement Fees. *Breast Care* 2013;8:282–8. doi:10.1159/000354249.
- [41] Loveman E, Jones J, Hartwell D, Bird A, Harris P, Welch K, et al. The clinical effectiveness and cost-effectiveness of topotecan for small cell lung cancer: a systematic review and economic evaluation. *Health Technol Assess* 2010;14. doi:10.3310/hta14190.
- [42] Hall PS, Hulme C, McCabe C, Oluboyede Y, Round J, Cameron DA. Updated Cost-Effectiveness Analysis of Trastuzumab for Early Breast Cancer: A UK Perspective Considering Duration of Benefit, Long-Term Toxicity and Pattern of Recurrence. *Pharmacoeconomics* 2011;29:415–32. doi:10.2165/11588340-000000000-00000.
- [43] Dutch National Health Care Institute. Medicijnkosten n.d. <http://medicijnkosten.nl/>.
- [44] Elting LS, Cooksley C, Chambers M, Cantor SB, Manzullo E, Rubenstein EB. The burdens of cancer therapy. Clinical and economic outcomes of chemotherapy-induced mucositis. *Cancer* 2003;98:1531–9. doi:10.1002/cncr.11671.
- [45] Wang G, Zhang Z, Ayala C, Wall HK, Fang J. Costs of heart failure-related hospitalizations in patients aged 18 to 64 years. *Am J Manag Care* 2010;16:769–76.
- [46] Bouwmans C, Janssen J, Huijgens P, Uyl-de Groot C. Costs of haematological adverse events in chronic myeloid leukaemia patients: a retrospective cost analysis of the treatment of anaemia, neutropenia and thrombocytopenia in patients with chronic myeloid leukaemia. *J Med Econ* 2009;12:164–9. doi:10.3111/13696990903149479.
- [47] Frederix, G. Disease specific methods for economic evaluations of breast cancer therapies. University Utrecht, 2013.
- [48] Lachaine J, Yelle L, Kaizer L, Dufour A, Hopkins S, Deuson R. Chemotherapy-induced emesis: quality of life and economic impact in the context of current practice in Canada. *Support Cancer Ther* 2005;2:181–7. doi:10.3816/SCT.2005.n.011.
- [49] Ward S, Simpson E, Davis S, Hind D, Rees A, Wilkinson A. Taxanes for the adjuvant treatment of early breast cancer: systematic review and economic evaluation. *Health Technol Assess Winch Engl* 2007;11:1–144.
- [50] Brown T, Pilkington G, Bagust A, Boland A, Oyee J, Tudur-Smith C, et al. Clinical effectiveness and cost-effectiveness of first-line chemotherapy for adult patients with locally advanced or metastatic non-small cell lung cancer: a systematic review and economic evaluation. *Health Technol Assess* 2013;17. doi:10.3310/hta17310.
- [51] Ward S, Scope A, Rafia R, Pandor A, Harnan S, Evans P, et al. Gene expression profiling and expanded immunohistochemistry tests to guide the use of adjuvant chemotherapy in breast cancer management: a systematic review and cost-effectiveness analysis. *Health Technol Assess* 2013;17. doi:10.3310/hta17440.
- [52] Collins R, Fenwick E, Trowman R, Perard R, Norman G, Light K, et al. A systematic review and economic model of the clinical effectiveness and cost-effectiveness of docetaxel in combination with prednisone or prednisolone for the treatment of hormone-refractory metastatic prostate cancer. *Health Technol Assess Winch Engl* 2007;11:iii – iv, xv – xviii, 1–179.
- [53] National Collaborating Centre for Cancer (UK). Ovarian Cancer: The Recognition and Initial Management of Ovarian Cancer. Cardiff (UK): National Collaborating Centre for Cancer (UK); 2011.
- [54] Horizon Scanning Centre, NHS. Denosumab (Xgeva) for recurrent or unresectable giant cell tumour of the bone – first or second line. 2012.
- [55] Trastuzumab (Herceptin) for HER2 positive early breast cancer – two year regimen. Horizon Scanning Centre, NHS; 2010.
- [56] Picot J, Cooper K, Bryant J, Clegg A. The clinical effectiveness and cost-effectiveness of bortezomib and thalidomide in combination regimens with an alkylating agent and a corticosteroid for the first-line treatment of multiple myeloma: a systematic review and economic evaluation. *Health Technol Assess* 2011;15. doi:10.3310/hta15410.
- [57] Elting LS, Cantor SB, Martin CG, Hamblin L, Kurtin D, Rivera E, et al. Cost of chemotherapy-induced thrombocytopenia among patients with lymphoma or solid tumors. *Cancer* 2003;97:1541–50. doi:10.1002/cncr.11195.

- [58] Elting LS, Cooksley CD, Chambers MS, Garden AS. Risk, outcomes, and costs of radiation-induced oral mucositis among patients with head-and-neck malignancies. *Int J Radiat Oncol Biol Phys* 2007;68:1110–20. doi:10.1016/j.ijrobp.2007.01.053.
- [59] Carlson J, Reyes C, Oestreicher N, Lubeck D, Ramsey S, Veenstra D. Comparative clinical and economic outcomes of treatments for refractory non-small cell lung cancer (NSCLC). *Lung Cancer* 2008;61:405–15. doi:10.1016/j.lungcan.2007.12.023.
- [60] Stokes ME, Muehlenbein CE, Marciniak MD, Faries DE, Motabar S, Gillespie TW, et al. Neutropenia-related costs in patients treated with first-line chemotherapy for advanced non-small cell lung cancer. *J Manag Care Pharm JMCP* 2009;15:669–82.
- [61] Braunschweig F, Cowie MR, Auricchio A. What are the costs of heart failure? *Europace* 2011;13:ii13–7. doi:10.1093/europace/eur081.
- [62] Sheng Duh M, Reynolds Weiner J, Lefebvre P, Neary M, Skarin AT. Costs associated with intravenous chemotherapy administration in patients with small cell lung cancer: a retrospective claims database analysis. *Curr Med Res Opin* 2008;24:967–74. doi:10.1185/030079908X280464.
- [63] Delea TE, Amdahl J, Diaz J, Nakhaipour HR, Hackshaw MD. Cost-effectiveness of pazopanib versus sunitinib for renal cancer in the United States. *J Manag Care Spec Pharm* 2015;21:46–54, 54a – b.
- [64] OECD.Stat. OECD (2013) n.d. <http://stats.oecd.org/index.aspx?queryid=22519#>.
- [65] XE currency Converter n.d. <http://www.xe.com/>.
- [66] Peasgood T, Ward SE, Brazier J. Health-state utility values in breast cancer. *Expert Rev Pharmacoecon Outcomes Res* 2010;10:553–66. doi:10.1586/erp.10.65.
- [67] CEA Registry Website > Home - Blog n.d. <https://research.tufts-nemc.org/cear4/> (accessed May 5, 2015).
- [68] Briggs et al. *Decision modelling for Health Economic Evaluation*. Oxford: Oxford university press; 2006.
- [69] Fenwick E, Marshall DA, Levy AR, Nichol G. Using and interpreting cost-effectiveness acceptability curves: an example using data from a trial of management strategies for atrial fibrillation. *BMC Health Serv Res* 2006;6:52. doi:10.1186/1472-6963-6-52.
- [70] Devlin N, Parkin D. Does NICE have a cost-effectiveness threshold and what other factors influence its decisions? A binary choice analysis. *Health Econ* 2004;13:437–52. doi:10.1002/hec.864.
- [71] Koleva-Kolarova RG, Greuter MJW, van Kruchten M, Vermeulen KM, Feenstra T, Buskens E, et al. The value of PET/CT with FES or FDG tracers in metastatic breast cancer: a computer simulation study in ER-positive patients. *Br J Cancer* 2015;112:1617–25. doi:10.1038/bjc.2015.138.
- [72] Schueller G, Schueller-Weidekamm C, Helbich TH. Accuracy of ultrasound-guided, large-core needle breast biopsy. *Eur Radiol* 2008;18:1761–73. doi:10.1007/s00330-008-0955-4.
- [73] Soni A, Ren Z, Hameed O, Chanda D, Morgan CJ, Siegal GP, et al. Breast cancer subtypes predispose the site of distant metastases. *Am J Clin Pathol* 2015;143:471–8. doi:10.1309/AJCPY05FSV3UPEXS.
- [74] Auguste P, Barton P, Hyde C, Roberts TE. An economic evaluation of positron emission tomography (PET) and positron emission tomography/computed tomography (PET/CT) for the diagnosis of breast cancer recurrence. *Health Technol Assess Winch Engl* 2011;15:iii – iv, 1–54. doi:10.3310/hta15180.
- [75] Center for Medicare and Medicaid Services. Physician fee schedule and acute inpatient prospective payment system. n.d.
- [76] n.d.
- [77] Uyl-de Groot CA, Senft A, de Bree R, Leemans CR, Hoekstra OS. Chest CT and whole-body 18F-FDG PET are cost-effective in screening for distant metastases in head and neck cancer patients. *J Nucl Med Off Publ Soc Nucl Med* 2010;51:176–82. doi:10.2967/jnumed.109.067371.
- [78] Melnikow J, Tancredi DJ, Yang Z, Ritley D, Jiang Y, Slee C, et al. Program-Specific Cost-Effectiveness Analysis: Breast Cancer Screening Policies for a Safety-Net Program. *Value Health* 2013;16:932–41. doi:10.1016/j.jval.2013.06.013.
- [79] Bouwmans C, Janssen J, Huijgens P, Uyl-de Groot C. Costs of haematological adverse events in chronic myeloid leukaemia patients: a retrospective cost analysis of the treatment of anaemia, neutropenia and thrombocytopenia in patients with chronic myeloid leukaemia. *J Med Econ* 2009;12:164–9. doi:10.3111/13696990903149479.

- [80] Loveman E, Jones J, Hartwell D, Bird A, Harris P, Welch K, et al. The clinical effectiveness and cost-effectiveness of topotecan for small cell lung cancer: a systematic review and economic evaluation. *Health Technol Assess Winch Engl* 2010;14:1–204. doi:10.3310/hta14190.
- [81] Elting LS, Cantor SB, Martin CG, Hamblin L, Kurtin D, Rivera E, et al. Cost of chemotherapy-induced thrombocytopenia among patients with lymphoma or solid tumors. *Cancer* 2003;97:1541–50. doi:10.1002/cncr.11195.
- [82] Lachaine J, Yelle L, Kaizer L, Dufour A, Hopkins S, Deuson R. Chemotherapy-induced emesis: quality of life and economic impact in the context of current practice in Canada. *Support Cancer Ther* 2005;2:181–7. doi:10.3816/SCT.2005.n.011.
- [83] Ward S, Simpson E, Davis S, Hind D, Rees A, Wilkinson A. Taxanes for the adjuvant treatment of early breast cancer: systematic review and economic evaluation. *Health Technol Assess Winch Engl* 2007;11:1–144.
- [84] Carlson J, Reyes C, Oestreicher N, Lubeck D, Ramsey S, Veenstra D. Comparative clinical and economic outcomes of treatments for refractory non-small cell lung cancer (NSCLC). *Lung Cancer* 2008;61:405–15. doi:10.1016/j.lungcan.2007.12.023.
- [85] Annemans L, Giaccone G, Vergnenègre A. The cost-effectiveness of paclitaxel (Taxol) + cisplatin is similar to that of teniposide + cisplatin in advanced non-small cell lung cancer: a multicountry analysis. *Anticancer Drugs* 1999;10:605–15.
- [86] Stokes ME, Muehlenbein CE, Marciniak MD, Faries DE, Motabar S, Gillespie TW, et al. Neutropenia-related costs in patients treated with first-line chemotherapy for advanced non-small cell lung cancer. *J Manag Care Pharm JMCP* 2009;15:669–82.
- [87] Brown T, Pilkington G, Bagust A, Boland A, Oyee J, Tudur-Smith C, et al. Clinical effectiveness and cost-effectiveness of first-line chemotherapy for adult patients with locally advanced or metastatic non-small cell lung cancer: a systematic review and economic evaluation. *Health Technol Assess Winch Engl* 2013;17:1–278. doi:10.3310/hta17310.
- [88] Elting LS, Cooksley CD, Chambers MS, Garden AS. Risk, Outcomes, and Costs of Radiation-Induced Oral Mucositis Among Patients With Head-and-Neck Malignancies. *Int J Radiat Oncol* 2007;68:1110–20. doi:10.1016/j.ijrobp.2007.01.053.
- [89] Stevanovic J, Denee L, Koenders JM, Postma MJ. Incidence Description and Costs of Acute Heart Failure in the Netherlands. *Value Health* 2014;17:A328. doi:10.1016/j.jval.2014.08.597.
- [90] Braunschweig F, Cowie MR, Auricchio A. What are the costs of heart failure? *Europace* 2011;13:ii13–7. doi:10.1093/europace/eur081.
- [91] van der Linden N, Buter J, Pescott CP, Lalisang RI, de Boer JP, de Graeff A, et al. Treatments and costs for recurrent and/or metastatic squamous cell carcinoma of the head and neck in the Netherlands. *Eur Arch Oto-Rhino-Laryngol Off J Eur Fed Oto-Rhino-Laryngol Soc EUFOS Affil Ger Soc Oto-Rhino-Laryngol - Head Neck Surg* 2015. doi:10.1007/s00405-015-3495-y.
- [92] Hamada S, Hinotsu S, Ishiguro H, Toi M, Kawakami K. Cross-National Comparison of Medical Costs Shared by Payers and Patients: A Study of Postmenopausal Women with Early-Stage Breast Cancer Based on Assumption Case Scenarios and Reimbursement Fees. *Breast Care* 2013;8:282–8. doi:10.1159/000354249.
- [93] Reed SD, Li Y, Anstrom KJ, Schulman KA. Cost effectiveness of ixabepilone plus capecitabine for metastatic breast cancer progressing after anthracycline and taxane treatment. *J Clin Oncol Off J Am Soc Clin Oncol* 2009;27:2185–91. doi:10.1200/JCO.2008.19.6352.
- [94] Barlow WE, Taplin SH, Yoshida CK, Buist DS, Seger D, Brown M. Cost comparison of mastectomy versus breast-conserving therapy for early-stage breast cancer. *J Natl Cancer Inst* 2001;93:447–55.
- [95] Botteman M, Barghout V, Stephens J, Hay J, Brandman J, Aapro M. Cost effectiveness of bisphosphonates in the management of breast cancer patients with bone metastases. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2006;17:1072–82. doi:10.1093/annonc/mdl093.
- [96] Konski A. Radiotherapy is a cost-effective palliative treatment for patients with bone metastasis from prostate cancer. *Int J Radiat Oncol* 2004;60:1373–8. doi:10.1016/j.ijrobp.2004.05.053.
- [97] Puri V, Crabtree TD, Kymes S, Gregory M, Bell J, Bradley JD, et al. A comparison of surgical intervention and stereotactic body radiation therapy for stage I lung cancer in high-risk patients: a decision analysis. *J Thorac Cardiovasc Surg* 2012;143:428–36. doi:10.1016/j.jtcvs.2011.10.078.

- [98] Landman MP, Feurer ID, Pinson CW, Moore DE. Which is more cost-effective under the MELD system: primary liver transplantation, or salvage transplantation after hepatic resection or after loco-regional therapy for hepatocellular carcinoma within Milan criteria?: HPB. HPB 2011;13:783–91. doi:10.1111/j.1477-2574.2011.00355.x.
- [99] Kimman ML, Dirksen CD, Voogd AC, Falger P, Gijzen BCM, Thuring M, et al. Economic evaluation of four follow-up strategies after curative treatment for breast cancer: results of an RCT. Eur J Cancer Oxf Engl 1990 2011;47:1175–85. doi:10.1016/j.ejca.2010.12.017.
- [100] Alvarado MD, Mohan AJ, Esserman LJ, Park CC, Harrison BL, Howe RJ, et al. Cost-Effectiveness Analysis of Intraoperative Radiation Therapy for Early-Stage Breast Cancer. Ann Surg Oncol 2013;20:2873–80. doi:10.1245/s10434-013-2997-3.
- [101] Riley GF, Potosky AL, Lubitz JD, Kessler LG. Medicare payments from diagnosis to death for elderly cancer patients by stage at diagnosis. Med Care 1995;33:828–41.
- [102] Frederix, G. Disease specific methods for economic evaluations of breast cancer therapies. University Utrecht, 2013.
- [103] Dutch Health Care Insurance Board. Guide for research costs - Methods and standard cost prices for economic evaluations in healthcare n.d.
- [104] Ward S, Scope A, Rafia R, Pandor A, Harnan S, Evans P, et al. Gene expression profiling and expanded immunohistochemistry tests to guide the use of adjuvant chemotherapy in breast cancer management: a systematic review and cost-effectiveness analysis. Health Technol Assess Winch Engl 2013;17:1–302. doi:10.3310/hta17440.
- [105] Centers for Medicare & Medicaid Services. Average sales pricing (ASP) Drug Pricing. Medicare Part B drug average sales price. 2014.
- [106] Sheng Duh M, Reynolds Weiner J, Lefebvre P, Neary M, Skarin AT. Costs associated with intravenous chemotherapy administration in patients with small cell lung cancer: a retrospective claims database analysis*. Curr Med Res Opin 2008;24:967–74. doi:10.1185/030079908X280464.
- [107] Collins R, Fenwick E, Trowman R, Perard R, Norman G, Light K, et al. A systematic review and economic model of the clinical effectiveness and cost-effectiveness of docetaxel in combination with prednisone or prednisolone for the treatment of hormone-refractory metastatic prostate cancer. Health Technol Assess Winch Engl 2007;11:iii – iv, xv – xviii, 1–179.
- [108] Jones L, Hawkins N, Westwood M, Wright K, Richardson G, Riemsma R. Systematic review of the clinical effectiveness and cost-effectiveness of capecitabine (Xeloda) for locally advanced and/or metastatic breast cancer. Health Technol Assess Winch Engl 2004;8:iii, xiii – xvi, 1–143.
- [109] Ovarian Cancer: The Recognition and Initial Management of Ovarian Cancer. Cardiff (UK): National Collaborating Centre for Cancer (UK); 2011.
- [110] Horizon Scanning Centre, NHS. Trastuzumab (Herceptin) for HER2 positive early breast cancer – two year regimen. 2010.
- [111] Dutch National Health Care Institute. Medicijnkosten n.d.
- [112] Picot J, Cooper K, Bryant J, Clegg AJ. The clinical effectiveness and cost-effectiveness of bortezomib and thalidomide in combination regimens with an alkylating agent and a corticosteroid for the first-line treatment of multiple myeloma: a systematic review and economic evaluation. Health Technol Assess Winch Engl 2011;15:1–204. doi:10.3310/hta15410.
- [113] Guest JF, Clegg JP, Davie AM, McCloskey E. Costs and consequences of using pamidronate compared with zoledronic acid in the management of breast cancer patients in the UK. Curr Med Res Opin 2005;21:805–15. doi:10.1185/030079905X40472.
- [114] Lee KS, Ro J, Nam B-H, Lee ES, Kwon Y, Kwon HS, et al. A randomized phase-III trial of docetaxel/capecitabine versus doxorubicin/cyclophosphamide as primary chemotherapy for patients with stage II/III breast cancer. Breast Cancer Res Treat 2008;109:481–9. doi:10.1007/s10549-007-9672-y.
- [115] Sparano JA, Wang M, Martino S, Jones V, Perez EA, Saphner T, et al. Weekly Paclitaxel in the Adjuvant Treatment of Breast Cancer. N Engl J Med 2008;358:1663–71. doi:10.1056/NEJMoa0707056.
- [116] Sonke GS, Mandjes IA, Holtkamp MJ, Schot M, van Werkhoven E, Wesseling J, et al. Paclitaxel, carboplatin, and trastuzumab in a neo-adjuvant regimen for HER2-positive breast cancer. Breast J 2013;19:419–26. doi:10.1111/tbj.12124.

- [117] Suter T., Cook-Bruns N, Barton C. Cardiotoxicity associated with trastuzumab (Herceptin) therapy in the treatment of metastatic breast cancer. *The Breast* 2004;13:173–83. doi:10.1016/j.breast.2003.09.002.
- [118] Lidgren M, Wilking N, Jönsson B, Rehnberg C. Health related quality of life in different states of breast cancer. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil* 2007;16:1073–81. doi:10.1007/s11136-007-9202-8.
- [119] Milne RJ, Heaton-Brown KH, Hansen P, Thomas D, Harvey V, Cubitt A. Quality-of-life valuations of advanced breast cancer by New Zealand women. *PharmacoEconomics* 2006;24:281–92.
- [120] Färkkilä N, Torvinen S, Roine RP, Sintonen H, Hänninen J, Taari K, et al. Health-related quality of life among breast, prostate, and colorectal cancer patients with end-stage disease. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil* 2014;23:1387–94. doi:10.1007/s11136-013-0562-y.
- [121] Prescott RJ, Kunkler IH, Williams LJ, King CC, Jack W, van der Pol M, et al. A randomised controlled trial of postoperative radiotherapy following breast-conserving surgery in a minimum-risk older population. The PRIME trial. *Health Technol Assess Winch Engl* 2007;11:1–149, iii – iv.
- [122] Stacie Hudgens. 1046P - Impact of treatment with eribulin (ERI) or capecitabine (CAP) for metastatic breast cancer (MBC) on EQ-5D utility derived from EORTC QLQ-C30, *Annals of Oncology* (2014) 25 (suppl_4): iv357-iv360.; n.d. doi:10.1093/annonc/mdl341.
- [123] Lloyd A, Nafees B, Narewska J, Dewilde S, Watkins J. Health state utilities for metastatic breast cancer. *Br J Cancer* 2006;95:683–90. doi:10.1038/sj.bjc.6603326.
- [124] Tachi T, Teramachi H, Tanaka K, Asano S, Osawa T, Kawashima A, et al. The Impact of Outpatient Chemotherapy-Related Adverse Events on the Quality of Life of Breast Cancer Patients. *PLOS ONE* 2015;10:e0124169. doi:10.1371/journal.pone.0124169.
- [125] Dyer MT, Goldsmith KA, Sharples LS, Buxton MJ. A review of health utilities using the EQ-5D in studies of cardiovascular disease. *Health Qual Life Outcomes* 2010;8:13. doi:10.1186/1477-7525-8-13.
- [126] Sanz MA, Aledort L, Mathias SD, Wang X, Isitt JJ. Analysis of EQ-5D scores from two phase 3 clinical trials of romiplostim in the treatment of immune thrombocytopenia (ITP). *Value Health J Int Soc Pharmacoeconomics Outcomes Res* 2011;14:90–6. doi:10.1016/j.jval.2010.10.017.
- [127] Goldhirsch A, Gelber RD, Piccart-Gebhart MJ, De Azambuja E, Procter M, Suter TM, et al. 2 years versus 1 year of adjuvant trastuzumab for HER2-positive breast cancer (HERA): An open-label, randomised controlled trial. *The Lancet* 2013;382:1021–8. doi:10.1016/S0140-6736(13)61094-6.
- [128] Kelly C, Alken. Benefit risk assessment and update on the use of docetaxel in the management of breast cancer. *Cancer Manag Res* 2013;357. doi:10.2147/CMAR.S49321.
- [129] Arimidex, Tamoxifen, Alone or in Combination Trialists' Group, Buzdar A, Howell A, Cuzick J, Wale C, Distler W, et al. Comprehensive side-effect profile of anastrozole and tamoxifen as adjuvant treatment for early-stage breast cancer: long-term safety analysis of the ATAC trial. *Lancet Oncol* 2006;7:633–43. doi:10.1016/S1470-2045(06)70767-7.
- [130] Smith IE. Efficacy and safety of Herceptin in women with metastatic breast cancer: results from pivotal clinical studies. *Anticancer Drugs* 2001;12 Suppl 4:S3–10.
- [131] National Cancer Institute. Common Terminology Criteria for Adverse Events (CTCAE) Version 4.0 2010.
- [132] Tan-Chiu E, Yothers G, Romond E, Geyer CE, Ewer M, Keefe D, et al. Assessment of cardiac dysfunction in a randomized trial comparing doxorubicin and cyclophosphamide followed by paclitaxel, with or without trastuzumab as adjuvant therapy in node-positive, human epidermal growth factor receptor 2-overexpressing breast cancer: NSABP B-31. *J Clin Oncol Off J Am Soc Clin Oncol* 2005;23:7811–9. doi:10.1200/JCO.2005.02.4091.

Supplementary material

Table 1: Dosages per systemic regimen

ddAC*	2 cycles of 2-weekly 600mg/m ² cyclophosphamide (C) and 60 mg/m ² doxorubicin (A)
CD*	2 cycles of 3-weekly 75 mg/m ² docetaxel (D) and two-daily 1000 mg/m ² capecitabine (C) during 14 days
PTC*	3-cycles Weekly AUC=3 carboplatin (C), 70 mg/m ² paclitaxel (P) and Trastuzumab (T), with first dose of 4mg/kg and subsequent of 2 mg/kg
FE ⁷⁵ C-T*	3-cycles In one day: 5-FU 500 mg/kg, epirubicine 90 mg/m ² , cyclofosfamide 500 mg/m ² , and on the first day of the first cycle pertuzumab 420 mg
Tamoxifen*	20 mg oral once daily for 2,5 years followed by Anastrozole
Anastrozole*	1mg/daily oral for 5-years following Tamoxifen
Zometa**	4 mg intravenously every 3-4 weeks for 9 months then 4 mg every 12 weeks. Total 5 years
Paclitaxel**	80 mg/m ² intravenously every 3 weeks (in combination with Trastuzumab). Patient will be treated until death.
Trastuzumab**	Once every 3 weeks: first day of first cycle 8 mg/kg intravenously and 6 mg/kg the other cycles (in combination with Paclitaxel). Patient will be treated until death.
Capecitabine**	1000-1250 mg/m ² intravenously every 12 hours. After 14 days, 7 days rest. Patient will be treated until death.

* Neo-adjuvant and adjuvant setting. ** Palliative setting

Table 2: Patient characteristics of the group used to derive imaging performance (n=545).

Total (N)	545
Mean age in years (range)	51
DM found at screening	
Total	9
Bone only	5
Lung only	1
Liver only	1
Multiple*	2

*More than 3 lesions and thus not considered oligometastasis or curable.

Table 3: Patient characteristics of the group used to derive primary breast cancer treatment (n=157).

	ER-positive/ HER2-negative	ER-positive/ HER2-positive	ER-negative/ HER2-positive	ER-negative/ HER2-negative	Total
	n (%)	n (%)	n (%)	n (%)	n (%)
	94 (60)	15 (10)	18 (11)	30 (19)	157 (100)
Pre-operative systemic treatment (PST)					
Initial PST ¹					
No PST	9 (10)	0 (0)	0 (0)	0 (0)	10 (6)
ddAC	81 ² (86)	2 (13)	0 (0)	29 (97)	111 (71)
CD	4 ² (4)	0 (0)	0 (0)	0 (0)	4 (30)
PTC	0 (0)	13 ⁷ (87)	18 ⁷ (100)	1 (3)	32 (20)
FE ⁷⁵ C-T	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Other	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	94 (100)	15 (100)	18 (100)	30 (100)	157 (100)
Second PST					
No PST	70 (74)	12 (80)	18 (100)	21 (70)	121 (77)
ddAC	1 (1)	0 (0)	0 (0)	3 ⁹ (10)	4 (3)
CD	7 ⁴ (7)	0 (0)	0 (0)	2 (7)	9 (6)
PTC	1 (1)	2 (13)	0 (0)	0 (0)	3 (2)
FE ⁷⁵ C-T	0 (0)	1 (7)	0 (0)	1 (3)	2 (1)
Other	15 ⁵ (16)	0 (0)	0 (0)	3 ⁵ (10)	18 (11)
	94 (100)	15 (100)	18 (100)	30 (100)	157 (100)
Adjuvant treatment					
Tamoxifen					
yes	84 (89)	15 (100)	0 (0)	0 (0)	99 (63)
no	10 (11)	0 (0)	18 (100)	30 (100)	58 (37)
	94 (100)	15 (100)	18 (100)	30 (100)	157 (100)
Aromatase inhibitors (AI)					
yes	25 (28)	12 (80)	0 (0)	0 (0)	37 (24)
no	69 (72)	3 (20)	18 (100)	30 (100)	120 (76)
	94 (100)	15 (100)	18 (100)	30 (100)	157 (100)
Chemotherapy					
No chemo	73 (78)	13 (87)	18 (100)	24 (80)	128 (82)
ddAC	1 ² (1)	0 (0)	0 (0)	0 (0)	1 (1)
CD	1 (1)	0 (0)	0 (0)	0 (0)	1 (1)
PTC	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
FE ⁷⁵ C-T	0 (0)	2 (13) ⁸	0 (0)	0 (0)	2 (1)
Other	19 ⁵ (19)	0 (0)	0 (0)	6 (20)	23 (15)
	94 (100)	15 (100)	18 (100)	30 ^{5,10} (100)	157 (100)
Trastuzumab					
yes	0 (0)	11 (73)	14 (78)	0 (0)	25 (16)
no	94 (100)	4 (27)	4 (22)	30 (100)	132 (84)
	94 (100)	15 (100)	18 (100)	30 (100)	157 (100)
Combinations of systemic treatment (Initial PST/ Second PST / Adjuvant)					
ddAC/ AI/ Px ¹¹	16 (17)				
ddAC/ Px / AI ¹²	15 (16)				
ddAC / --/ AI	40 (43)				
ddAC/ DC/ AI	5 (5)				
ddAC/ ddAC/ Px	1 (1)				
ddAC/ DC/ AI & DC	1 (1)				
ddAC/ PTC/ AI	1 (1)				

CD/ --/ AI	3 (3)				
CD/ CD/ AI	1 (1)				
--/ --/ AI	9 (10)				
PTC/ AI/ Tras.		7 (47)			
PTC/ FECT/ AI & Tras.		2 (13)			
PTC/ AI		3 (20)			
PTC/ FECT/ AI		1 (7)			
ddAC/ PTC/ AI & Tras.		2 (13)			
PTC/ herc			14 (78)		
PTC			4 (22)		
ddAC/ ddAC				3 (10)	
ddAC/ Px				1 (3)	
ddAC/-- / Px				2 (7)	
ddAC				19 (63)	
ddAC/ Px / Px				2 (7)	
ddAC/ DC/ Px				1 (3)	
ddAC/ DC				1 (3)	
PTC/ FECT/ Px				1 (3)	
	94 (100)	15 (100)	18 (100)	30 (100)	157 (100)
Breast radiotherapy					
yes	86 (91)	12 (80)	14 (78)	23 (77)	135 (86)
no	8 (9)	3 (20)	4 (22)	7 (23)	22 (14)
	94 (100)	15 (100)	18 (100)	30 (100)	157 (100)
Breast surgery					
WLE	54 (56)	8 (53)	7 (39)	16 (53)	85 (53)
Ablatio	42 (42)	7 (47)	11 (61)	14 (47)	74 (47)
	9 ⁶ (100)	15 (100)	18 (100)	30 (100)	157 (100)

Abbreviations: Px= Paclitaxel; FECT= FE⁷⁵C-T; Tras= Trastuzumab; ddAC= dose-dense cyclophosphamide and doxorubicin; DC=docetaxel and capecitabine; PTC= Paclitaxel, trastuzumab and carboplatin; FEC⁷⁵-T= Fluorouracil, Epirubicine, and cyclophosphamide; AI= aromatase inhibitor; Px= paclitaxel; WLE= wide local excision.

¹ Patients receiving PST were enrolled in a response- adaptive trial, where a treatment switch could occur after a specific number of cycles.

² Three patients received TAC (docetaxel, doxorubicin, cyclophosphamide) instead of ddAC, yet they were included in this group.

³ For one patient the number of CD cycles were not specified, yet they were assumed to follow the CD regimen in table 1.

⁴ Only D in 2nd and 3rd course, yet we assumed it follow the CD regimen in table 1.

⁵ Many patients in this group received 9 cycles of paclitaxel, thus this was assumed the most common treatment of "other". Patients that received <9 cycles, were assumed to have 9.

⁶ Two patients had both types of surgery.

⁷ One patient received PTC plus pertuzumab, yet this was not taken into account.

⁸ Two patients received 3 cycles, yet they were assumed to follow dosage of the FE⁷⁵C-T regimen as specified in table 1.

⁹ Two patients received high dose alkylating chemotherapy as part of a trial, yet we assumed they received ddAC as in table 1.

¹⁰ Four patients received paclitaxel accompanied by carboplatin, yet we assumed they received 9 cycles of paclitaxel.

¹¹ As other usually involved Paclitaxel, this was assumed.

¹² As in our model hormonal treatment was assumed AI, in the "combined systemic treatments" this was always termed as AI, regardless of the actual treatment received.

Table 4: Model input parameters

Variables different per subtypes												
	ER-positive/HER2-negative				ER-negative/HER2-positive				ER-negative/HER2-negative			
	Mean Value	Lower limit	Upper Limit		Mean Value	Lower limit	Upper Limit		Mean Value	Lower limit	Upper Limit	
Metastasis distributions												
Bone metastasis	0.586	0.284	0.961		0.286	0.096	0.977		0.370	0.144	0.956	
Liver metastasis	0.218	0.043	0.958		0.390	0.177	0.950		0.205	0.019	0.988	
Lung metastasis	0.195	0.012	0.993		0.325	0.123	0.960		0.425	0.160	0.953	
Single metastasis	0.333	0.197	0.505		0.333	0.101	0.644		0.467	0.200	0.751	
									0.433	0.160	0.743	
Variables different per countries												
	The Netherlands				The United Kingdom				The United States			
	Mean Value	Lower limit	Upper Limit	Source ^a	Mean Value	Lower limit	Upper Limit	Source ^a	Mean Value	Lower limit	Upper Limit	Source ^a
Costs of imaging, biopsy, chemotherapy-related toxicities, cancer treatment and health states (€ for NL, £ for UK, and \$ for US)												
Full body PET/CT	1163	380	2577	NL-NZA reference cost[13]	1458	374	3352	NHS reference costs 2008[74]	1077	379	2345	CPT/HCPCS reference fees [75,76]
Chest X-ray	77	28	173	NL-NZA reference cost[13]	112	33	247	NHS reference costs 2008[74]	77	25	210	CPT/HCPCS reference fees [75,76]
Bone scintigraphy	282	89	739	NL-NZA reference cost[13]	193	58	452	NHS reference costs 2008[74]	162	42	350	CPT/HCPCS reference fees [75,76]
Liver sonography	88	26	230	NL-NZA reference cost[13]	66	18	144	NHS reference costs 2008[74]	197	64	430	CPT/HCPCS reference fees [75,76]
MRI (for bone metastases) ^b	274	78	579	NL-NZA reference cost[13]	274	86	608	NHS reference costs 2008[74]	969	290	2122	CPT/HCPCS reference fees [75,76]
CT (thorax)	192	48	422	[77]	159	43	355	NHS reference costs 2008[74]	545	172	1289	CPT/HCPCS reference fees [75,76]

CT (full body)	NA	NA	NA	NA	NA	NA	NA	NA	NA	574	163	1408	CPT/HCPCS reference fees [75,76]
Biopsy (US guided)	199	69	516	NA	NA	NL-NZA reference cost[13]	166	49	382	NHS reference costs 2008[74]	538	158	1134 [78]
Thrombocytopenia	3422	6	26442	69	516	[79]	1427	359	3373	NHS reference costs 2008[80]	1037	781	1366 [81]
Vomiting	92	30	210	30	210	[82] c	467	130	1029	NHS reference costs 2005[83]	122	44	256 [84]
Neutropenia	972	309	2471	309	2471	[85]	538	154	1144	NHS reference costs 2005[83]	3099	2515	3798 [86]
Febrile neutropenia	4486	1296	9797	1296	9797	[85]	6260	1709	13462	NHS reference costs 2009[87]	12148	9436	15317 [86]
Mucositis	3723	1284	8568	1284	8568	[81]	932	252	2170	NHS reference costs 2009[87]	3600	1096	7973 [88]
Cardio toxicities (symptomatic)	4632	1300	11086	1300	11086	[89]	1970	476	4557	NHS reference costs 2013/14	8500	2580	18712 [90]
Breast radiotherapy	8840	2397	18661	2397	18661	[91]	10748	3212	24087	NHS reference costs 2012[92]	13383	4612	28780 [93]
Breast surgery	6633	2159	14042	2159	14042	NKI-NZA reference cost	4023	1368	8679	NHS reference costs 2012[92]	12454	10551	14868 [94]
Bone radiotherapy	1750	568	4273	568	4273	NKI-NZA reference cost	958	303	2127	NHS reference costs 2006[95]	1799	482	4169 [96]
Lung surgery (metas)	10891	3477	23997	3477	23997	NKI-NZA reference cost	9782	2679	22613	NHS reference costs 2013/14	15034	4420	35137 [97]
Liver surgery (metas)	10393	3027	22398	3027	22398	NKI-NZA reference cost	10102	3521	23456	NHS reference costs 2009[80]	3500	4272	119371 [98]
Follow up (stable)	2447	1535	3581	1535	3581	[99]	245	67	550	NHS reference costs 2008[42]	1833	497	4111 [100]

Breast cancer death	16350	4713	36202	NK(-NZA reference cost	14730	5234	31026	NHS reference costs 2013/14	19436	6291	49066	[101]
Neo(adjuvant) systemic treatments												
3xAC	2047	619	4475	[102,103]	2241	713	5088	NHS reference costs 2005[83,104], 2009[104]	1881	662	4707	[105,106]
3xDC	5601	1800	12697	[102,103]	5522	1772	12046	NHS reference costs 2003[107,108], 2009[104]	11735	3018	25027	[105],CPT/ HCPCS reference fees [75,76]
8xPTC	9073	2773	20236	[102,103]	11317	3634	24808	NHS reference costs 2007[83], 2008, 2009[104], 2010[109], 2012[110]	16476	5591	42329	[105,106], CPT/HCPCS reference fees [75,76]
FE ⁷⁵ C-T	5463	1503	11836	[102,103, 111]	6879	1850	16408	NHS reference costs 2007[83], 2008, 2009[104], 2010[109], 2012[110]	11314	3399	28297	[105,106], CPT/HCPCS reference fees [75,76]
AI	466	582	349	[102,111]	324	405	243	NHS reference costs 2013/14, 2009[104]	212	265	159	[105], CPT/ HCPCS reference fees [75,76]
9xPx	5448	1736	13403	[102,103]	8662	2303	20124	NHS reference costs 2007[83], 2009[104]	5374	1573	11644	CPT/HCPCS reference fees [75,76]
AD	1947	611	4659	[102,103]	2221	692	4609	NHS reference costs 2007[83], 2009[104], 2011[112]	1842	601	4270	[105,106], CPT/HCPCS reference fees [75,76]

Trastuzumab (1 year)	17812	4842	40856	[102,103]	28291	9382	67969	NHS reference costs 2008, 2009[104], 2012[110]	70479	23664	161676	[105,106] CPT/HCPCS reference fees [75,76]
Metastatic systemic treatments												
Trastuzumab + Paclitaxel (1 year)	21647	16235	27058	[102,103]	37877	28408	47364	NHS reference costs 2007[49], 2008, 2009[51], 2012[55]	70344	52758	87929	[57,63] CPT/HCPCS reference fees[75,76]
Anastrozole + Zometa (year 1)	2440	1830	3050	[102,111]	2618	1964	3273	NHS reference costs 2003[113] 2009[51], 2013/14	2350	1762	2937	[105] CPT/HCPCS reference fees[76]
Anastrozole + Zometa (>year 1)	1232	924	1540	[102,111]	1322	992	1653	NHS reference costs 2003[113] 2009[51], 2013/14	1178	883	1472	[105] CPT/HCPCS reference fees[76]
Paclitaxel	10148	7611	12685	[102,103]	17134	12851	21418	NHS reference costs 2007[49], 2009[51]	9942	7457	12428	[57,63] CPT/HCPCS reference fees [75,76]
Capecitabine	3637	2728	4546	[102]	3969	2977	4961	NHS reference costs 2003[108], 2009[51]	29544	22158	36930	[105] CPT/HCPCS reference fees[76]
Zometa (1 year)	2421	1816	3026	[102,111]	2592	1944	3240	NHS reference costs 2003[113] 2009[51]	2344	1758	2930	[105] CPT/HCPCS reference fees[76]
Zometa (>1 year)	1270	952	1587	[102,111]	1296	972	1620	NHS reference costs 2003[113] 2009[51]	1172	879	1465	[105] CPT/HCPCS reference fees[76]

Variables that are equal for all models						
	Mean value	Lower limit	Upper Limit	Source/ Observations		
Imaging performance						
Sensitivity PET/CT	92%	74%	100%	NKI		
Specificity PET/CT	98%	95%	100%	NKI		
Sensitivity CI	13%	0%	31%	NKI		
Specificity CI	94%	89%	97%	NKI		
Transition probabilities of breast cancer death						
Bone metastasis						
Year 1	0.240	0.092	0.417	[20]		
Year 2	0.132	0.033	0.277	[20]		
Year 3	0.227	0.074	0.394	[20]		
Year 4	0.333	0.171	0.557	[20]		
Year 5	0.382	0.198	0.586	[20]		
Visceral metastasis						
Year 1	0.410	0.236	0.615	[20]		
Year 2	0.424	0.251	0.622	[20]		
Year 3	0.265	0.120	0.643	[20]		
Year 4	0.320	0.170	0.529	[20]		
Year 5	0.294	0.156	0.529	[20]		
Bone plus visceral metastasis						
Year 1	0.480	0.303	0.693	[20]		
Year 2	0.308	0.117	0.514	[20]		
Year 3	0.361	0.159	0.564	[20]		
Year 4	0.320	0.167	0.504	Assumed as visceral		
Year 5	0.294	0.136	0.491	Assumed as visceral		
Chemotherapy-related toxicities ^d						
Vomiting						
Anthracyclines (plus trastuzumab ^e)	0.240	0.102	0.394	Assumed as anthracyclines alone[14]		

Neutropenia				
Taxanes	0.720	0.556	0.856	[114]
Anthracyclines (plus trastuzumab)	0.850	0.711	0.960	[114]
Anthracyclines plus taxanes	0.460	0.406	0.511	[115]
PTC	0.720	0.480	0.913	[116]
Febrile neutropenia (anthracyclines plus taxanes)	0.160	0.124	0.205	[115]
Hand-foot-syndrome (taxanes)	0.220	0.095	0.381	[114]
Mucositis (taxanes)	0.100	0.025	0.265	[114]
Thrombocytopenia (PTC)	0.360	0.374	0.844	[116]
Cardio toxicity ^a (anthracyclines plus trastuzumab)	0.280	0.160	0.428	[117]
Utilities ^a				
Metastasis	0.685	0.656	0.845	[118]
Bone metastasis	0.310	0.270	0.350	[119]
Stable disease	0.690	0.630	0.753	[118]
Terminal disease	0.447	0.285	0.604	[120]
Radiotherapy	0.780	0.740	0.810	[121]
Surgery ^h	0.855	0.341	1	[66]
Vomiting	0.640	0.640	0.805	[122]
Febrile neutropenia	0.540	0.145	0.956	[123]
Mucositis	0.530	0.130	0.998	[124]
Cardio toxicity (symptomatic)	0.545	0.200	0.985	[125]
Thrombocytopenia	0.770	0.687	0.913	[126]
Hormonal treatment	0.648	0.458	0.923	[118]

Abbreviations: US=ultrasound; NA= not applicable; ddAC= dose-dense cyclophosphamide and doxorubicin; DC=docetaxel and capecitabine; PTC= Paclitaxel, trastuzumab and carboplatin; FEC⁷⁵-T= Fluorouracil, Epirubicine, and cyclophosphamide; AI= aromatase inhibitor; Px= paclitaxel.

Neither weekly Paclitaxel, single Trastuzumab or Anastrozole were documented have any serious side effects $\geq 10\%$. [127–129]

^a If no bibliographic reference is added to the source it means we derived it directly from the reference source.

^b Calculate as the average of upper body, lower body and spine scan.

^c No Dutch source was found.

^d In our dataset, ddAC was given with PEG-filgrastim, which results in a similar toxicity profile standard AC regimen, defined as anthracyclines in the table.

^e Assumed equal as AC, as adding T does not really affect vomiting¹⁰¹. In fact, cardio-toxicity is the only 'combined' side effect, thus the remaining side effects of AC + T are assumed those of AC.

^f A review on Trastuzumab by Suter et al¹⁰⁸ only identifies the combination of AC + T as having ≥ 10% incidence of cardio-toxicity.

^g Presented utility weights of adverse events grade III/IV (common NCTCN criteria¹⁰²). Values are from EQ-5D questionnaires (UK or Europe), except for febrile neutropenia, derived from conventional gamble.

^h SD assumed of 0,1.

Table 5: Upper margin of cost of PET/CT and palliative treatment to attain cost-effectiveness

	ER-negative/ HER2-negative	ER-positive/ HER2-positive	Suggestion to reach cost-effectiveness in all subtypes
US	-	Only if palliative regimen <€28.000 & PET/CT costs \$1000	Lower PET/CT costs to \$1000, but also lower palliative treatment costs in ER- positive/ HER2-positive
NL	Only if palliative regimen <€3.000 & PET/CT costs €600	Only if palliative regimen <€3.000 & PET/CT costs €600	Lower PET/CT costs to €600, but also lower palliative treatment costs in ER- positive/HER2-positive and in ER-negative/ HER2-negative
UK	Only if palliative regimen <£3.000 & PET/CT costs £500	Only if palliative regimen <£3.000 & PET/CT costs £500	Lower PET/CT costs to £500, but also lower palliative treatment costs in ER- positive/HER2-positive and in ER-negative/ HER2-negative

Palliative treatment in ER-positive/HER2-positive is Trastuzumab plus Paclitaxel, and ER-negative/HER2-negative, Paclitaxel monotherapy. Treatment costs are yearly costs given with metastatic dosages, as detailed in supplementary table 1.

Figure 1: One way sensitivity analysis of the NL

Figure 2: One way sensitivity analysis of the US

Technical details of the imaging modalities

Whole body ^{18}F -FDG PET/CT was performed with the scanner Gemini TF, Philips, Cleveland, Ohio, USA. CI comprised of bone scintigraphy (Symbia dual head gamma camera, Siemens, Erlangen, Germany) based on whole-body scanning anterior and posterior simultaneously, 2.5 h after administration of 555 MBq of $^{99\text{m}}\text{Tc}$ hydroxymethane diphosphonate), ultrasound of the liver (Hitachi Ultrasound (Hitachi Medical Corporation, model EZU- MT27-S1, Tokyo, Japan) and chest radiograph (posterior–anterior and lateral view; Buckydiagnost CS, Philips, Hamburg, Germany). Patients were prepared for the whole-body PET/CT scan with a fasting period of 6 h. Before intravenous injection 180–240 MBq ^{18}F -FDG 10 mg diazepam was orally administered and blood glucose levels had to be <10 mmol/l. After a resting period of approximately 60 min the PET/CT acquisition was made in supine position from the base of the skull to the upper half of the femora (1.30 min per bed position).

Description of the Markov model

During the 5-years' time horizon, patients who entered the model with presence of DM or developed a DM due to a false result at screening, could: i) remain stable (simulated by remaining in the same state); ii) die from a non-breast cancer event (simulated by a transition to the non-breast cancer death state); or iii) die from breast cancer (simulated by a transition to the terminal state and ultimately to the breast cancer death state). Patients who did not develop DM could remain stable or die from a non-breast cancer event.

In the 1st-year cycle the costs of primary breast cancer treatment (PST, breast surgery, breast radiotherapy, adjuvant chemotherapy and chemotherapy-related adverse events, except cardio-toxicities which were included in year 2) were attributed to all patients. Additionally, positive patients at baseline were attributed costs of biopsy, plus local DM treatment (single DM) or palliative treatment (multiple DM) to TPs, and plus confirmation scans to FPs. Confirmation scans for FP patients under the PET/CT strategy consisted of bone MRI, liver sonography, and CT lung, and full-body PET/CT, under the CI strategy. While TN patients did not incur additional costs, FN patients incurred costs of confirmation scans, biopsy, and additional systemic and local DM treatment. FNs confirmation scans for the PET/CT strategy consisted of the modality of CI intended for the region of interest, and for the conventional strategy the full body PET/CT.

Stable patients, without prior detection of DM or after local treatment of single liver or lung DM, were assigned the costs of follow-up (mammogram plus a specialist visit). Patients who remained stable after being detected with single bone DM received bisphosphonates, and patients who remained stable after being detected with multiple DM received palliative treatment. Details on treatments used in the model for DM patients are detailed in the "model input data section" and its posology details in supplementary table 1. The costs of a cardio-toxic adverse events were added in the 2nd-year cycle, as the cardio-toxic pick of incidence is 1-year after treatment initiation[132]. Additional costs of palliative treatment were assigned to patients who died from a breast-cancer event, while patients dying from other causes than breast cancer had no additional costs.

During the 1st-year cycle, utilities were also attributed based on the TP, FP, TN and FN classification. Thus, TPs were assigned the utility of DM; FPs and TNs, the weighted average utility of all primary breast cancer treatments undergone during that year (using time as a weighting factor); and FNs, the utility of bone DM, representing the quality-of-life of painful metastases. Patients who remained stable in the following cycles were assigned the utility of the adjuvant treatment received, or in its absence, of stable disease. Utility for cardio-toxic adverse events was assigned in the 2nd-year cycle. Patients who died from a breast-cancer event were assigned the utility of palliative treatment.

Results of the one way sensitivity analysis

The one-way sensitivity analysis to all model parameters revealed cost-effectiveness in the US is driven by either the prevention of FPs palliative treatment costs (in ER-positive/HER2-negative and ER-negative/HER2-negative), the decrease in PET/CT costs together with an increase in CI costs (ER-negative/HER2-positive) or the decrease in TPs palliative treatment costs (ER-positive/HER2-positive), in the NL by either a decrease in PET/CT costs (ER-positive/HER2-negative and ER-negative/HER2-positive), or a combination of a decrease in PET/CT costs and TPs palliative treatment costs (ER-negative/HER2-negative and ER-positive/HER2-positive), and in the UK, by either a decrease in PET/CT costs (ER-positive/HER2-negative and ER-negative/HER2-positive), a combination of a decrease in PET/CT costs and TPs palliative treatment costs (ER-negative/HER2-negative), or a decrease in TPs palliative treatment costs (ER-positive/HER2-positive).

PART V

GENERAL DISCUSSION AND ANNEX

CHAPTER 9

General discussion

In view of the high research and development costs of new technologies [1], especially in the late phases of development [2], there has been growing interest in the use of economic evaluations in early development phases of medical technologies. However, despite this gain in popularity, its use in real-life applications is not fully exploited yet [3–5]. This thesis contributes to the literature on early cost-effectiveness (CE) analysis (CEAs), as well as on value of information (VOI) and resource modeling analysis, particularly applied to medical technologies for emerging breast cancer interventions. As breast cancer still remains the leading cause of cancer death in women, especially in advanced stages [6], the pursuance of new treatments for these patients is ongoing. Through the methodology applied in this thesis, our aim is to inform on development, further research and adoption decisions.

Main findings

Predictive biomarkers: personalize systemic treatment

In **chapter 2** we concluded that clinical translation of predictive biomarkers in neoadjuvant chemotherapy (NACT) for breast cancer is lacking, and we highlighted the underlying biological and clinical reasons that may underlie this (i.e., the existence of tumor heterogeneity or strict demands on study design to demonstrate clinical utility). Furthermore, we suggested that early health technology assessment (HTA) could be useful in helping decision-making during the biomarker development process. For instance on choosing optimal study design characteristics (via multi criteria decision analysis) or in informing on the cost-effectiveness of specific biomarker test characteristics (via CEA).

In **chapter 3** we developed an early cost-effectiveness model that simulates the clinical application of the BRCA1-like biomarker, by using the Multiplex Ligation-dependent Probe Amplification (MLPA) test. This model showed that treating triple negative breast cancer (TNBC) with personalized high dose alkylating chemotherapy (HDAC) based on the BRCA1-like predictive biomarker is not yet cost-effective. Furthermore, the minimum prevalence of the biomarker and positive predictive value of its diagnostic test for this biomarker strategy to become cost-effective are 58.5% and 73.0% respectively.

Chapter 4 was motivated by the discovery that by further characterizing BRCA1-like tumors with two other biomarkers, XIST and 53BP1, responses to HDAC could increase from 70% to a 100%. We thus compared the CE of treating TNBCs with the following biomarker strategies: 1) BRCA1-like measured by the MLPA test; 2) BRCA1-like measured by the array comparative genomic hybridization (aCGH) test; 3) strategy 1 combined with the XIST and 53BP1 biomarkers; and 4)

strategy 2 combined with the XIST and 53BP1 biomarkers, or by current practice. We concluded that there is excessive uncertainty around the CE outcomes to decide on a preferred treatment strategy for TNBCs. We subsequently determined that further research is valuable to reduce this uncertainty up to costs of €639. This information could optimally be gathered by setting up four simultaneous ancillary studies to ongoing randomized clinical trials (like the NCT01057069) with a total sample size of 3000 patients. These retrospective studies should separately collect data on 1) BRCA1-like prevalence, PPV and treatment response rates (TRRs) in biomarker negative patients - as determined by the MLPA test, and TRR in the whole population of TNBC patients; 2) same parameters as strategy 1 - as determined by the aCGH test alone and by the combination of the MLPA and aCGH tests with the XIST and 53BP1 biomarkers; 3) model costs; and 4) model utilities.

Imaging techniques: monitoring systemic treatment

In **chapter 5** we systematically reviewed literature on the performance of imaging for NACT response guidance separately per breast cancer subtype. We concluded that there is insufficient evidence to draw on subtype specific recommendations for NACT guidance. Further steps towards reaching consensus on specific study design characteristics (i.e., pCR definitions, imaging protocols or time intervals between baseline and response monitoring) are necessary before initiating well-designed studies that generate higher levels of evidence.

In **chapter 6** we calculated the cost-effectiveness of a response-guided NACT scenario for the treatment of hormone-receptor positive breast cancers. The scenario started with all patients being treated with two cycles of docetaxel, doxorubicin, and cyclophosphamide. After monitoring with ultrasound, patients that responded to the treatment continued with 6 cycles of the initial regimen, while non-respondents were switched to four cycles of vinorelbine and capecitabine. Results of our CEA indicated that this response-guided NACT scenario is cost-effective (vs conventional NACT). While prospective validation of the effectiveness of this scenario is advisable from a clinical perspective, we suggest that early CEAs are used to prioritize further research from a broader health economic perspective, by identifying which parameters contribute most to current decision uncertainty.

In **chapter 7** we calculated the cost-effectiveness and the resource demands of implementation for a response-guided NACT scenario for the treatment of hormone-receptor positive/HER2-negative breast cancers. The scenario started with all patients being treated with 3 cycles of dose dense doxorubicin and cyclophosphamide. After MRI monitoring, patients that responded to the treatment continued with 3 cycles of the same regimen, while non-respondents switched to 3 cycles of dose dense docetaxel and capecitabine. The novelty of this study is that outcomes

were calculated for a conventional and a full implementation scenario of this intervention in the Netherlands. This addition is important because the variation of emerging interventions' uptake can affect cost-effectiveness estimates. Conclusions of this study are that at current evidence levels, response-guided NACT is cost-effective under both scenarios. This means that response-guided NACT is less costly and more effective than conventional NACT and that at any uptake level cost-effectiveness is positively impacted. In terms of resource demands, we concluded that The Netherlands has sufficient personnel and MRI capacity for a future full implementation scenario.

Imaging techniques: screening for distant metastasis

In **chapter 8** we calculated the cost-effectiveness of distant metastases screening with PET/CT in stage II/III breast cancer patients for the four major breast cancer subtypes in three countries: the Netherlands, the United Kingdom (UK) and the United States (US). We concluded that PET/CT is expected cost-effective only for screening HER2-negative patients treated in the US. This is because in this subtype the costs of palliative treatment are higher in false positives (FP) than in true positives (TP), and as PET/CT increases TP but decreases FP, this results in cost-savings. PET/CT cost-effectiveness in the Netherlands and in the UK could be attained by reductions in PET/CT costs and by reductions in palliative treatment costs.

Determinants of the cost-effectiveness of personalized interventions

In line with previous literature [7–14], this thesis concluded that four main parameters define the CE of personalized interventions (PI): the performance of the diagnostic test, the effectiveness of the treatment (within the target group), the prevalence of the biomarker and the costs of treatment or the costs of diagnostic testing. It thus is important that these parameters are present in any economic evaluation of a PI [13,15]. This is particularly important in the case of performance, which has often been ignored in published CEAs [9,12,16,17].

From our thesis chapters, we gathered a set of observations on the behavior of these determinants, which are in line with other literature [7,12,14,18,19]:

- 1) with good diagnostic test performance and favorable treatment effect PIs are likely to be more cost-effective than all-comers strategies i.e., equal treatment to all patients ([7,18], chapter 3, 6, 7); even in low prevalent diseases ([7,14], chapter 3) and at low intervention uptake rates (chapter 7);
- 2) treatment effectiveness drives the effect part of CE vs. test performance (chapter 8);

- 3) treatment costs usually drive the cost part of CE vs. test costs. This is because costs of targeted treatments tend to be higher than test costs ([12,16], chapter 3,4, 6). Only when this relationship changes i.e. test costs are higher than drug costs, test costs drive the cost part of CE [10,8].

Methodological considerations

Early cost effectiveness analysis

An iterative process

A characteristic of early CEAs is that decision-analytical models need to be populated with available data at the time of analysis, which is likely scarce, and then are complemented with data derived from literature and/or assumptions (usually derived from expert elicitation). As early economic evaluation is not an on-off assessment of a technology, literature suggests that iterations of these models should be performed when more data becomes available [20,21]. This thesis research encompasses the first iteration of such models and provides the groundwork for next iterations. For example, the BRCA1-like biomarker (chapter 2) is an excellent case to illustrate the impact that adding additional effectiveness information has on model outcomes and decision uncertainty, as several additional clinical validation studies have been or are about to be published for this biomarker.

Cooperation with other stakeholders

During this thesis it became apparent that early CEAs to quantify an intervention's expected impact on survival, QALYs and/or costs, and to draw lessons for their improvement or for further research, were not always as influential as expected. Findings were sometimes met with resistance. This is not unique in this thesis work, as confirmed by the observations of the Clinical and Translational Science Awards (CTSA) Program of the National Institutes of Health (NIH) in the US [22]. Reasons that may underlie this are a "publish first", or otherwise protectionist attitude (of own projects and publications), or simply a lack of importance given to CEAs use in the scientific research process. Views on the benefits of using early CEAs vary in the scientific community [23]. Collaborations of clinicians and researchers in CEA-related projects are often limited of scope and rare as such [24]. Our chapter 2 shows an example of such collaboration to disseminate the use of early CEAs during predictive biomarker research. These collaborations require accurate selection of partners (i.e., stakeholders that belief on the importance of each others' work, that are willing to invest time on understanding each others' concerns and that

have a shared objective to improve the translation of promising technologies into practice) and a clear definition of roles and expectations from the outset.

Value of information analysis

Study designs for further research

In VOI literature, it is common that further research is calculated as derived from one single RCT, hence assuming that a new RCT is started to gather all necessary data. Our full VOI analysis was instead presented as a portfolio of retrospective studies to an ongoing RCT. This approach was chosen because it was unrealistic to assume that an RCT for a set of newly discovered biomarkers with limited evidence (chapter 4) would be started. As we are aware that the shortfall of using retrospective and uncontrolled data is its proneness to bias, we purposely choose that these studies were performed along an ongoing RCT, as this guarantees higher levels of evidence (LOE) [25].

A limitation we encountered in projecting further research with retrospective studies is that maximum studies size is restricted to that of the ongoing RCT trial. We suggest that further studies using this approach to calculate VOI overcome this limitation by either 1) finding similar RCTs than the one used for the VOI calculations to obtain the desired data by setting additional retrospective studies to it; or 2) by assuming that a new prospective RCT will be conducted to collect data for samples bigger than the ongoing RCT. This will demand accounting for extra costs in the ENBS calculations for these samples.

Personalized interventions

The way in which CEAs have traditionally been performed for drugs i.e. given to large populations is being challenged by its use in PIs. PIs have different characteristics than drugs and thus different demands. Some of these issues that arise when using CEAs in PI have been nicely illustrated by some [12,14,28,29], while others have generated recommendations [13,29]. Below, we highlight the most important issues we faced in this regard.

Incorporating performance

Performance is highly dependent on the assumptions that underlay its definition. Three main assumptions limit our CEAs: 1) the assumed effectiveness of the non-cross resistant treatment given in response-guided NACT interventions; 2) the follow-up time used to determine the responsive patients to a specific PI; and 3) the cut-off values to determine the biomarker positive population or the responsive patients to a specific PI.

The first assumption was forced by the absence of control groups in the group of patients that were switched to a non-cross resistant treatment after being classified as irresponsive to the initial treatment by imaging. This made it impossible to distinguish if irresponsiveness in this group of patients was due to non-cross resistant treatment ineffectiveness or due to a wrong classification by imaging (and patients should have continued with the initial treatment). Hence imaging performance could not be calculated unless an assumption on treatment effectiveness was made. The other two assumptions were required to apply current CEA methodology to PIs. CEA models incorporate performance in terms of sensitivity and specificity. These measures can only be derived if specific assumptions on thresholds and cut-offs are made.

Effectiveness data quality

PI narrow down the size of the relevant population, and as a consequence generating reliable effectiveness data from RCTs requires longer times and great expenses [29]. Furthermore, one needs to collect effectiveness data on both, the test detecting the biomarker, and the biomarker predicting response to the drug. In the course of these thesis, we only used RCT data for one model (chapter 6), the remaining were populated with data coming from single cohorts (chapter 7) or from several observational sources (chapter 3, 4, 8). The shortfall of using CEAs with lower LOE than RCTs is that this type of data is more prone to bias and can lead to cost-effectiveness recommendations with large degrees of uncertainty or even to decision-makers unwilling to make decisions based on these. These shortfalls can be minimized by collecting effectiveness evidence following best practices [30]; considering all relevant evidence, selecting those that fit best the model demands, while simultaneously aiming for the highest LOE. Furthermore, policies of the type of “coverage with evidence development” should be promoted. These policies contain an RCT to generate better evidence on a new technology/drug and a CEA to demonstrate its additional values, and in the meantime, the new technology/drug is already being reimbursed. A first example has recently started in the Netherlands (BRCA1-like biomarker for stage III breast cancer).

Capturing health related quality of life

The use of predictive testing can decrease patients health related quality of life (HRQoL) due to discomfort (while testing) or anxiety (while awaiting the test results). In this thesis we did not account for this temporary decrease in HRQoL. While discomfort was not really a concern in any chapter, anxiety could have been important in all of them. In chapters 3, 4, 6 and 7 due to the possibility (or not) of benefiting from a treatment, and in chapter 8, due to the presence (or absence) of metastatic disease. We do expect that accounting for this HRQoL decrease could have affected the results of chapters 6 and 7, as in these chapters none of the assigned utilities

was especially low. On the other hand, in chapters 3, 4 and 8 where low utilities were already present due to the use of toxic treatments or due to the severity of the disease, this omission is not expected to modify our conclusions. We suggest that CEAs of PI that have relatively high HRQoL i.e., less severe interventions or diseases, pay (more) attention to the possible impact that patient discomfort and/or anxiety caused by testing can have in HRQoL.

High levels of uncertainty

CEAs in the field of personalized medicine have increased uncertainty, in terms of both model structure (structural uncertainty) and input data (parameter uncertainty) [13]. This is due to the higher complexity of PIs models, which need to mimic more complex pathways than that of drugs. This is also consequence of the lack of large prospective studies on long-term effectiveness data, which requires extensive extrapolation of models costs and benefits. These limitations were present in all our CEAs. Parameter uncertainty was tackled by the standard probabilistic sensitivity analysis, while structural uncertainty was taken into account via additional one- and two- way SA [30]. Scenario analysis could also have been used to further explore these uncertainties. Furthermore, overall model uncertainty can be addressed by performing VOI analysis. We suggest that CEAs of PI consider these additional analysis to PSA, so decision-makers can understand the robustness of findings and draw adequate recommendations.

Wider organizational implications

The addition of a test into clinical practice has generally wide organizational implications i.e., the creation of new working pathways, of new infrastructures, the training of new personnel or the purchase of new diagnostic machinery [28]. CEAs do not always account for the additional resources that may be needed at the time of implementation [31]. This usual omission stems from CEAs origin in assessing the “one fits all” kind of drugs, where the only resource concern was the availability of the compound itself. For PI, accounting for additional resources becomes more relevant. In fact so relevant, that if ignored, it may jeopardize the translation of promising technologies. Methods like resource modeling analysis [31] can help anticipating these demands to facilitate PIs translation and eventual implementation.

Current clinical and economical value, implications and future research

In this section we elaborate on the current clinical and cost-effectiveness evidence available for each PI by making use of a medical value map (see Figure 1 [32]). As these two types of evidence

are essential to support decisions on adoption and coverage, we elaborate on the implications of their current evidence level and suggest directions for further HTA research.

Figure 1: Medical value map. Adapted from a report entitled “Articulating the value of diagnostics: Challenges and opportunities” from Panaxea b.v. [32]. This map shows the value of an intervention based on its clinical and cost effectiveness evidence. We suggested a position for each of our case studies in this map (using the chapter numbers). Notice that chapters 2 and 5 were clinical literature reviews and thus have no data on cost effectiveness. Also, that chapter 8 is placed in two different quadrants. This is because the CEA of the PI intervention was assessed from different country perspective and resulted in different outcomes. Furthermore, we highlight that the place of the numbers within the squares does not indicate any grading of evidence. Footnotes: * HER2-negative (US perspective), > ER-/HER2+ (US perspective), ^ ER+/HER2+ (US perspective) and all subtypes (NL and UK perspective).

Predictive biomarkers: personalize systemic treatment

The clinical effectiveness of the BRCA1-like biomarker for predicting response to HDAC has so far been demonstrated in three studies [33,34] and two prospective RCTs are currently ongoing. The clinical effectiveness of the BRCA1-like plus XIST and 53BP1 combination has so far been demonstrated in one small retrospective study (*Schouten et al submitted*) backed up by pre-clinical studies [35–38]. The first cost-effectiveness evidence on either of the biomarker combinations has been provided in this thesis (chapter 3 and 4). This evidence indicates that it is still uncertain whether personalized HDAC based on any of these biomarker strategies is more cost-effective than using standard practice.

Our study results imply that until the BRCA1-like biomarker becomes cost-effective vs. current practice (chapter 3), or the CEA with all biomarker strategies depicts a clear “winner” (chapter 4), coverage of these predictive biomarkers will not occur, and standard chemotherapy will continue as the gold standard. Furthermore, higher LOE of effectiveness for all the biomarkers are required for its clinical adoption.

Evidence from two RCTs validating the BRCA1-like biomarker are expected in the coming 5 to 10 years (one is ongoing and one is about to start). Their positive outcome is likely to facilitate the adoption of the BRCA1-like biomarker into clinical practice. In terms of coverage, the BRCA1-like biomarker has recently entered a ‘coverage with evidence development’ type of agreement through one of these RCTs. The data resulting from this trial is expected to be used for future coverage decisions. Our model of chapter 3 could be re-analyzed with this new data and serve as the final confirmation for its coverage.

Further evidence on the effectiveness of the BRCA1-like plus XIST and 53BP1 combination could be derived retrospectively from these two ongoing BRCA1-like RCTs. Furthermore, as suggested by the results of our chapter 4, additional data on costs, other effectiveness-related parameters and utilities could also be derived from these RCTs. Subsequently, our model of chapter 4 could be updated and re-analyzed with these data and that generated from the BRCA1-like RCTs.

Other factors than clinical and cost-effectiveness evidence are expected to influence these biomarkers’ adoption; 1) the need for stem cell transplantation upon administration of HDAC, which adds risks for patients [39]; 2) the organizational implications of the different tests’ logistics; and 3) the tests’ costs, which depend on the number of samples used per run, the turnaround time between runs and the technique used. We suggest examining scenarios on these and other aspects prior to formal adoption in order to facilitate biomarker translation.

Imaging techniques: monitoring systemic treatment

Our review revealed that clinical evidence on the performance of imaging for NACT response guidance separately per breast cancer subtype is lacking. All included studies are of low LOE. They are underpowered, with heterogeneous study designs and outcome measures. Furthermore, there is absence of studies on the effectiveness of the whole response-guided NACT approach, which suggests that this approach is still young for its adoption into clinical practice. The first cost-effectiveness evidence on response-guided NACT has been presented in this thesis (chapter 6 and 7). These two CEAs demonstrated that response-guided NACT is likely to be cost-effective when adopted in clinical practice. While these results could imply low payer barriers, this is challenged by the low LOE of the input effectiveness data. Furthermore, the two selected studies have limited application into clinical practice, consequence of the use of non-standard drug regimens.

This implies that so far there is not enough evidence to support neither the clinical application nor the reimbursement of response-guided NACT and thus conventional NACT should continue as standard practice.

Our suggestion is that well-designed studies that generate higher LOE on the effectiveness of imaging in monitoring NACT in breast cancer are undertaken. However, prior steps towards reaching consensus on specific study design characteristics are required (i.e., pCR definitions, imaging protocols or time intervals between baseline and response monitoring). Thereafter, RCTs that mimic the response-guided NACT approach can be started. These studies have the advantage to not only inform on the effectiveness of imaging in monitoring NACT, but also on suitable treatment switches for not responders at imaging. An example of such trial is the AVATAXHER [40] which applied response-guided NACT in HER2 breast cancers using taxanes, trastuzumab and bevacizumab containing regimens [40]. As accounting for breast cancer subtypes dramatically reduces sample sizes, we suggest that all future studies are conducted in multi centric trials.

Imaging techniques: screening for distant metastasis

The clinical effectiveness of PET/CT in detecting DM in breast cancer is of low LOE, as evidence so far comes from three observational studies [41,42]. The generated cost-effectiveness evidence in this thesis indicates that cost-effectiveness differs between countries and subtypes. So far PET/CT is only expected cost-effective for screening HER2-negative patients treated in the US. To attain PET/CT cost-effectiveness in the Netherlands and in the UK, reductions in PET/CT costs and reductions in palliative treatment costs are warranted.

Our CE results imply that PET/CT can only be recommended to US payers and only for screening HER2-negative subtypes. For all other cases, conventional imaging should remain current practice.

Our results suggest that further studies that explore PET/CT effectiveness are needed before any consideration for its clinical implementation can be made. Furthermore, evidence on the differential long term outcomes of early detected DM (at screening) vs. late detected DM (at follow up after being missed at screening) per subtype are needed. If early detection of DM significantly improves survival, this will be an additional argument supporting the use of PET/CT. As previously mentioned that generating subtype specific data in a single institution may be challenging, we suggest collecting these data via a multicentre studies.

Concluding remarks and future directions

Breast cancer is a highly prevalent disease [6] and still remains the leading cause of cancer death in women [6]. Personalized medicine is an emerging approach to patient care, whose aim is to find the right treatment for the right patient at the right time [43]. The implementation of PIs in breast cancer treatment is expected to improve current breast cancer survival rates. Through the use of early CEAs, the chances of successfully translating promising biomarkers and targeted treatments into clinical practice are expected to increase.

This thesis has contributed to the literature on early CEAs as well as value of information analysis and resource modeling analysis by using emerging personalized breast cancer intervention studies. The results of these studies have been informative to developers of these interventions with regard to 1) the likely cost-effectiveness of these interventions given current evidence (chapter 3, 4, 6, 7, 8); 2) the development targets needed (chapter 3) and the additional research required to make these intervention cost-effective (chapter 4 and 6); 3) the resource requirements for implementing these interventions (chapter 7); 4) the state of the art of predictive biomarkers for NACT in breast cancer and imaging techniques' performance in NACT monitoring (chapters 2 and 5); and 5) the usefulness of early HTA methods during predictive biomarker research decision-making (chapter 2).

This thesis concluded that the BRCA1-like biomarker is at present the only biomarker with likely sufficient clinical evidence and expected economical evidence to be accepted by payers and doctors in the near future. As expected from emerging PI, all remaining case studies either lacked of effectiveness data to be accepted in the clinic, and/or had unfavorable or uncertain cost-effectiveness outcomes (Figure 1).

The methods used in this thesis are still not incorporated into routine practice (chapter 2). However, given the speed of scientific advances, it is expected that early CEAs and VOI that will assess effectiveness data of non-randomized RCTs [29] will become more common. This will permit deciding early on whether research on a specific PI should be continued instead of investing those resources elsewhere. As payers may be reluctant to take decisions based on these low LOE's, 'wait and see' or 'coverage with evidence development' conclusions are likely to become more common in CEAs as a result [29]. Moreover, the use of resource modeling as an annex to CEAs can anticipate adoption demands and speed up translation. We expect its use to become more extended, especially in later stages of development.

While this thesis dealt with single biomarker testing, it is expected that multiple testing, the use of panels and even whole genome testing will be widely considered in the near future. This will increase the complexity of CEAs. Challenges will include developing methods to incorporate genomic effectiveness data into economic evaluation frameworks, establishing appropriate methods to cost platform diagnostics with multiple applications, development of innovative evaluation frameworks outside the traditional model-based CEA by combining methods to evaluate additional HTA aspects like clinicians and patient behavior, and agreements on appropriate health outcome measures that permit more individualization. Communication between researchers, clinicians, health-economists and decision-makers in all stages of the translational research process will be necessary to ensure that appropriate data and methods for addressing the economic value of these complex diagnostic testing methods associated with targeted therapies are being developed.

References

- [1] Dorsey ER, de Roulet J, Thompson JP, Remnick JJ, Thai A, White-Stellato Z, et al. Funding of US biomedical research, 2003-2008. *JAMA* 2010;303:137-43. doi:10.1001/jama.2009.1987.
- [2] Moses H, Dorsey ER, Matheson DHM, Thier SO. Financial Anatomy of Biomedical Research. *JAMA* 2005;294:1333. doi:10.1001/jama.294.11.1333.
- [3] Jönsson B. Bringing in health technology assessment and cost-effectiveness considerations at an early stage of drug development. *Mol Oncol* 2015;9:1025-33. doi:10.1016/j.molonc.2014.10.009.
- [4] Steuten LM, Ramsey SD. Improving early cycle economic evaluation of diagnostic technologies. *Expert Rev Pharmacoecon Outcomes Res* 2014;14:491-8. doi:10.1586/14737167.2014.914435.
- [5] Pham B, Tu HA, Han D, Pechlivanoglou P, Miller F, Rac V, et al. Early economic evaluation of emerging health technologies: protocol of a systematic review. *Syst Rev* 2014;3:81. doi:10.1186/2046-4053-3-81.
- [6] Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JWW, Comber H, et al. Cancer incidence and mortality patterns in Europe: Estimates for 40 countries in 2012. *Eur J Cancer* 2013;49:1374-403. doi:10.1016/j.ejca.2012.12.027.
- [7] Trusheim MR, Burgess B, Hu SX, Long T, Averbuch SD, Flynn AA, et al. Quantifying factors for the success of stratified medicine. *Nat Rev Drug Discov* 2011;10:817-33. doi:10.1038/nrd3557.
- [8] Vegter S, Boersma C, Rozenbaum M, Wilffert B, Navis G, Postma MJ. Pharmacoeconomic evaluations of pharmacogenetic and genomic screening programmes: a systematic review on content and adherence to guidelines. *Pharmacoeconomics* 2008;26:569-87.
- [9] Oosterhoff M, van der Maas ME, Steuten LMG. A Systematic Review of Health Economic Evaluations of Diagnostic Biomarkers. *Appl Health Econ Health Policy* 2015. doi:10.1007/s40258-015-0198-x.
- [10] Hatz MHM, Schremser K, Rogowski WH. Is individualized medicine more cost-effective? A systematic review. *Pharmacoeconomics* 2014;32:443-55. doi:10.1007/s40273-014-0143-0.
- [11] Lee J, Tollefson E, Daly M, Kiel E. A generalized health economic and outcomes research model for the evaluation of companion diagnostics and targeted therapies. *Expert Rev Pharmacoecon Outcomes Res* 2013;13:361-70. doi:10.1586/erp.13.23.
- [12] Doble B, Tan M, Harris A, Lorgelly P. Modeling companion diagnostics in economic evaluations of targeted oncology therapies: systematic review and methodological checklist. *Expert Rev Mol Diagn* 2015;15:235-54. doi:10.1586/14737159.2014.929499.
- [13] Annemans L, Redekop K, Payne K. Current methodological issues in the economic assessment of personalized medicine. *Value Health J Int Soc Pharmacoeconomics Outcomes Res* 2013;16:S20-6. doi:10.1016/j.jval.2013.06.008.
- [14] San Miguel L, Hulstaert F. The importance of test accuracy in economic evaluations of companion diagnostics. *J Comp Eff Res* 2015;4:569-77. doi:10.2217/cer.15.41.
- [15] Dolan JG, Boohaker E, Allison J, Imperiale TF. Patients' preferences and priorities regarding colorectal cancer screening. *Med Decis Mak Int J Soc Med Decis Mak* 2013;33:59-70. doi:10.1177/0272989X12453502.
- [16] San Miguel L, Hulstaert F. The importance of test accuracy in economic evaluations of companion diagnostics. *J Comp Eff Res* 2015;4:569-77. doi:10.2217/cer.15.41.
- [17] Frank M, Mittendorf T. Influence of Pharmacogenomic Profiling Prior to Pharmaceutical Treatment in Metastatic Colorectal Cancer on Cost Effectiveness: A Systematic Review. *Pharmacoeconomics* 2013;31:215-28. doi:10.1007/s40273-012-0017-2.
- [18] Flowers CR, Veenstra D. The role of cost-effectiveness analysis in the era of pharmacogenomics. *Pharmacoeconomics* 2004;22:481-93.
- [19] Davis JC, Furstenthal L, Desai AA, Norris T, Sutaria S, Fleming E, et al. The microeconomics of personalized medicine: today's challenge and tomorrow's promise. *Nat Rev Drug Discov* 2009;8:279-86. doi:10.1038/nrd2825.
- [20] Sculpher M, Drummond M, Buxton M. The iterative use of economic evaluation as part of the process of health technology assessment. *J Health Serv Res Policy* 1997;2:26-30.

- [21] Vallejo-Torres L, Steuten LMG, Buxton MJ, Girling AJ, Lilford RJ, Young T. Integrating health economics modeling in the product development cycle of medical devices: a Bayesian approach. *Int J Technol Assess Health Care* 2008;24:459–64. doi:10.1017/S0266462308080604.
- [22] Scott CS, Nagasawa PR, Abernethy NF, Ramsey BW, Martin PJ, Hacker BM, et al. Expanding assessments of translational research programs: supplementing metrics with value judgments. *Eval Health Prof* 2014;37:83–97. doi:10.1177/0163278713500984.
- [23] Esmail LC, Roth J, Rangarao S, Carlson JJ, Thariani R, Ramsey SD, et al. Getting our priorities straight: a novel framework for stakeholder-informed prioritization of cancer genomics research. *Genet Med Off J Am Coll Med Genet* 2013;15:115–22. doi:10.1038/gim.2012.103.
- [24] Concannon TW, Fuster M, Saunders T, Patel K, Wong JB, Leslie LK, et al. A systematic review of stakeholder engagement in comparative effectiveness and patient-centered outcomes research. *J Gen Intern Med* 2014;29:1692–701. doi:10.1007/s11606-014-2878-x.
- [25] Simon RM, Paik S, Hayes DF. Use of archived specimens in evaluation of prognostic and predictive biomarkers. *J Natl Cancer Inst* 2009;101:1446–52. doi:10.1093/jnci/djp335.
- [26] Briggs et al. *Decision modelling for Health Economic Evaluation*. Oxford: Oxford university press; 2006.
- [27] Griffin S, Welton NJ, Claxton K. Exploring the Research Decision Space: The Expected Value of Information for Sequential Research Designs. *Med Decis Making* 2010;30:155–62. doi:10.1177/0272989X09344746.
- [28] Sorenson C, Tarricone R, Siebert M, Drummond M. Applying health economics for policy decision making: do devices differ from drugs? *Eur Eur Pacing Arrhythm Card Electrophysiol J Work Groups Card Pacing Arrhythm Card Cell Electrophysiol Eur Soc Cardiol* 2011;13 Suppl 2:ii54–8. doi:10.1093/europace/eur089.
- [29] Buchanan J, Wordsworth S, Schuh A. Issues surrounding the health economic evaluation of genomic technologies. *Pharmacogenomics* 2013;14:1833–47. doi:10.2217/pgs.13.183.
- [30] Briggs AH, Weinstein MC, Fenwick EAL, Karnon J, Sculpher MJ, Paltiel AD. Model Parameter Estimation and Uncertainty: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-6. *Value Health* 2012;15:835–42. doi:10.1016/j.jval.2012.04.014.
- [31] Thokala P, Dixon S, Jahn B. Resource Modelling: The Missing Piece of the HTA Jigsaw? *PharmacoEconomics* 2014. doi:10.1007/s40273-014-0228-9.
- [32] Panaxea. *Articulating the value of diagnostics: Challenges and opportunities*. 2015.
- [33] Schouten PC, Marme F, Aulmann S, Sinn H-P, Van Essen DF, Ylstra B, et al. Breast cancers with a BRCA1-like DNA copy number profile recur less often than expected after high-dose alkylating chemotherapy. *Clin Cancer Res Off J Am Assoc Cancer Res* 2014. doi:10.1158/1078-0432.CCR-14-1894.
- [34] Vollebergh MA, Lips EH, Nederlof PM, Wessels LFA, Schmidt MK, van Beers EH, et al. An aCGH classifier derived from BRCA1-mutated breast cancer and benefit of high-dose platinum-based chemotherapy in HER2-negative breast cancer patients. *Ann Oncol Off J Eur Soc Med Oncol ESMO* 2011;22:1561–70. doi:10.1093/annonc/mdlq624.
- [35] Rottenberg S, Nygren AOH, Pajic M, van Leeuwen FWB, van der Heijden I, van de Wetering K, et al. Selective induction of chemotherapy resistance of mammary tumors in a conditional mouse model for hereditary breast cancer. *Proc Natl Acad Sci U S A* 2007;104:12117–22. doi:10.1073/pnas.0702955104.
- [36] Rottenberg S, Vollebergh MA, de Hoon B, de Ronde J, Schouten PC, Kersbergen A, et al. Impact of intertumoral heterogeneity on predicting chemotherapy response of BRCA1-deficient mammary tumors. *Cancer Res* 2012;72:2350–61. doi:10.1158/0008-5472.CAN-11-4201.
- [37] Bouwman P, Aly A, Escandell JM, Pieterse M, Bartkova J, van der Gulden H, et al. 53BP1 loss rescues BRCA1 deficiency and is associated with triple-negative and BRCA-mutated breast cancers. *Nat Struct Mol Biol* 2010;17:688–95. doi:10.1038/nsmb.1831.
- [38] Bunting SF, Callén E, Wong N, Chen H-T, Polato F, Gunn A, et al. 53BP1 inhibits homologous recombination in Brca1-deficient cells by blocking resection of DNA breaks. *Cell* 2010;141:243–54. doi:10.1016/j.cell.2010.03.012.
- [39] Rodenhuis S, Bontenbal M, Beex LVAM, Wagstaff J, Richel DJ, Nooij MA, et al. High-dose chemotherapy with hematopoietic stem-cell rescue for high-risk breast cancer. *N Engl J Med* 2003;349:7–16. doi:10.1056/NEJMoa022794.

- [40] Coudert B, Pierga J-Y, Mouret-Reynier M-A, Kerrou K, Ferrero J-M, Petit T, et al. Use of [(18)F]-FDG PET to predict response to neoadjuvant trastuzumab and docetaxel in patients with HER2-positive breast cancer, and addition of bevacizumab to neoadjuvant trastuzumab and docetaxel in [(18)F]-FDG PET-predicted non-responders (AVATAXHER): an open-label, randomised phase 2 trial. *Lancet Oncol* 2014;15:1493–502.
- [41] Koolen BB, Vrancken Peeters M-JTFD, Aukema TS, Vogel WV, Oldenburg HSA, van der Hage JA, et al. 18F-FDG PET/CT as a staging procedure in primary stage II and III breast cancer: comparison with conventional imaging techniques. *Breast Cancer Res Treat* 2012;131:117–26. doi:10.1007/s10549-011-1767-9.
- [42] Niikura N, Costelloe CM, Madewell JE, Hayashi N, Yu T-K, Liu J, et al. FDG-PET/CT compared with conventional imaging in the detection of distant metastases of primary breast cancer. *The Oncologist* 2011;16:1111–9. doi:10.1634/theoncologist.2011-0089.
- [43] Jackson SE, Chester JD. Personalised cancer medicine. *Int J Cancer J Int Cancer* 2015;137:262–6. doi:10.1002/ijc.28940.

ANNEX

Summary

Samenvatting

Acknowledgements

List of publications

Curriculum vitae

Summary

Even though the idea of starting economic evaluations early in the product life cycle of medical technologies has gained popularity in the past few years, its use has not been fully exploited yet. In this thesis, we aimed to contribute to the literature on early cost-effectiveness analysis (CEA), value of information analysis and resource modeling analysis, particularly applied to medical technologies for emerging breast cancer interventions.

After a short introduction (**chapter 1**), this thesis is divided in three parts, distinguished by the type of technologies assessed: The first part focuses on predictive biomarkers to personalize systemic treatment (**chapters 2, 3, 4**), the second part focuses on imaging techniques to guide the personalization of neoadjuvant chemotherapy (NACT) (**chapters 5, 6, 7**), and the third part focuses on imaging as a tool to detect distant metastases (**chapter 8**).

Predictive biomarkers: personalize systemic treatment

In **chapter 2** we investigate the current research status of predictive biomarkers in NACT for breast cancer and discuss the challenges for their translation into clinical practice. Furthermore, we explore the current use of early health technology assessment (HTA) methods in this field and provide concrete guidance on how its use could benefit predictive biomarker translation. We concluded that clinical translation of predictive biomarkers in neoadjuvant chemotherapy (NACT) for breast cancer is lacking, and we highlighted the underlying biological and clinical reasons that may underlie this (i.e., the existence of tumor heterogeneity or strict demands on study design to demonstrate clinical utility). Furthermore, we suggested that early health technology assessment (HTA) could be useful in helping decision-making during the biomarker development process. For instance on choosing optimal study design characteristics (via multi criteria decision analysis) or in informing on the cost-effectiveness of specific biomarker test characteristics (via CEA).

Chapters 3 and 4 focus on two predictive biomarker strategies for high dose alkylating chemotherapy (HDAC) in triple negative breast cancer: BRCA1-like biomarker testing, and BRCA1-like plus XIST and the 53BP1 biomarker testing. In **chapter 3**, we developed an early cost-effectiveness model that simulates the clinical application of the BRCA1-like biomarker, by using the Multiplex Ligation-dependent Probe Amplification (MLPA) test. This model showed that at current performance levels this biomarker strategy is not yet cost-effective. Furthermore, the minimum prevalence of the biomarker and positive predictive value of its diagnostic test for this biomarker strategy to become cost-effective are 58.5% and 73.0% respectively.

In **chapter 4**, we extended this cost-effectiveness model to include the possibility to personalize HDAC based on the two aforementioned biomarker strategies, using two different BRCA1-like tests. We thus compared the CE of treating TNBCs with the following biomarker strategies: 1) BRCA1-like measured by the MLPA test; 2) BRCA1-like measured by the array comparative genomic hybridization (aCGH) test; 3) strategy 1 combined with the XIST and 53BP1 biomarkers; and 4) strategy 2 combined with the XIST and 53BP1 biomarkers, or by current practice. Based on this model, we were not able to discern one biomarker strategy likely to be more cost-effective than current practice. Subsequently, a value of information analysis was performed, and we found that further research would be valuable to identify the most cost-effective biomarker strategy up to costs of €639 million. This information could optimally be gathered by setting up four simultaneous ancillary studies to ongoing randomized clinical trials (like the NCT01057069) with a total sample size of 3000 patients. These retrospective studies should separately collect data on 1) BRCA1-like prevalence, PPV and treatment response rates (TRRs) in biomarker negative patients - as determined by the MLPA test, and TRR in the whole population of TNBC patients; 2) same parameters as strategy 1 - as determined by the aCGH test alone and by the combination of the MLPA and aCGH tests with the XIST and 53BP1 biomarkers; 3) model costs; and 4) model utilities.

Imaging techniques: monitoring systemic treatment

In **chapter 5** we systematically reviewed literature on the performance of imaging for NACT response guidance separately per breast cancer subtype. We concluded that there is insufficient evidence to draw on subtype specific recommendations for NACT guidance. Further steps towards reaching consensus on specific study design characteristics (i.e., pCR definitions, imaging protocols or time intervals between baseline and response monitoring) are necessary before initiating well-designed studies that generate higher levels of evidence.

In chapter 6 and 7 we constructed two early CEAs to calculate the expected cost-effectiveness of two emerging 'response-guided NACT' interventions i.e., where NACT treatment is adapted according to response assessed by imaging. In **chapter 6** we calculated the cost-effectiveness of a response-guided NACT scenario for the treatment of hormone-receptor positive breast cancers. The scenario started with all patients being treated with two cycles of docetaxel, doxorubicin, and cyclophosphamide. After monitoring with ultrasound, patients that responded to the treatment continued with 6 cycles of the initial regimen, while non-respondents were switched to four cycles of vinorelbine and capecitabine. Results of our CEA indicated that this response-guided NACT scenario is cost-effective (vs conventional NACT). While prospective validation of the effectiveness of this scenario is advisable from a clinical perspective, we suggest that early CEAs are used to prioritize further research from a broader health economic perspective, by identifying which parameters contribute most to current decision uncertainty.

In **chapter 7** we calculated the cost-effectiveness and the resource demands of implementation for a response-guided NACT scenario for the treatment of hormone-receptor positive/HER2-negative breast cancers. The scenario started with all patients being treated with 3 cycles of dose dense doxorubicin and cyclophosphamide. After MRI monitoring, patients that responded to the treatment continued with 3 cycles of the same regimen, while non-respondents switched to 3 cycles of dose dense docetaxel and capecitabine. The novelty of this study is that outcomes were calculated for a conventional and a full implementation scenario of this intervention in the Netherlands. This addition is important because the variation of emerging interventions' uptake can affect cost-effectiveness estimates. Conclusions of this study are that at current evidence levels, response-guided NACT is cost-effective under both scenarios. This means that response-guided NACT is less costly and more effective than conventional NACT and that at any uptake level cost-effectiveness is positively impacted. In terms of resource demands, we concluded that The Netherlands has sufficient personnel and MRI capacity for a future full implementation scenario.

Imaging techniques: screening for distant metastases

In **chapter 8** we calculated the cost-effectiveness of distant metastases screening with PET/CT in stage II/III breast cancer patients for the four major breast cancer subtypes in three countries: the Netherlands, the United Kingdom (UK) and the United States (US). We concluded that PET/CT is expected cost-effective only for screening HER2-negative patients treated in the US. This is because in this subtype the costs of palliative treatment are higher in false positives (FP) than in true positives (TP), and as PET/CT increases TP but decreases FP, this results in cost-savings. PET/CT cost-effectiveness in the Netherlands and in the UK could be attained by reductions in PET/CT costs and by reductions in palliative treatment costs.

To conclude, this thesis has contributed to the literature on early CEAs as well as value of information analysis and resource modeling analysis by using emerging personalized breast cancer intervention studies. The results of these studies have been informative to developers of these interventions with regard to 1) the likely cost-effectiveness of these interventions given current evidence (chapter 3, 4, 6, 7, 8); 2) the development targets needed (chapter 3) and the additional research required to make these intervention cost-effective (chapter 4 and 6); 3) the resource requirements for implementing these interventions (chapter 7); 4) the state of the art of predictive biomarkers for NACT in breast cancer and imaging techniques' performance in NACT monitoring (chapters 2 and 5); and 5) the usefulness of early HTA methods during predictive biomarker research decision-making (chapter 2).

Samenvatting

Het idee om economische evaluaties reeds in een vroeg stadium van de productlevenscyclus van een medische technologie te starten, heeft de afgelopen jaren aan populariteit gewonnen. Ondanks de toename in populariteit, lijkt het gebruik van deze analyses nog niet volledig geëxploiteerd te worden. Met dit proefschrift hadden wij tot doel bij te dragen aan de literatuur met betrekking tot vroege kosten-effectiviteitsanalyses (cost-effectiveness analysis (CEA)), ‘*value of information*’ (VOI) analyses en ‘*resource modelling*’ analyses, specifiek op het gebied van medische technologieën voor nieuwe interventies voor de behandeling van borstkanker.

Na de introductie (**hoofdstuk 1**) is dit proefschrift verdeeld in drie delen gebaseerd op de technologie die onderzocht werd. Het eerste deel richt zich op predictieve biomarkers om systemische anti-kanker behandeling te personaliseren (vroege diagnostiek voor “therapie-op-maat”) (**hoofdstuk 2,3,4**); het tweede deel richt zich op beeldvormende technieken om de respons op neoadjuvante chemotherapie te meten (**hoofdstuk 5,6,7**) en het derde deel richt zich op het toepassen van beeldvorming om afstandsmetastasen te ontdekken (**hoofdstuk 8**).

Predictieve biomarkers: personaliseren van systemische anti-kanker behandeling

In hoofdstuk 2 evalueerden we de huidige stand van zaken in het onderzoek met betrekking tot predictieve biomarkers voor neoadjuvante chemotherapie tegen borstkanker en bediscussieerden we de uitdaging voor de translatie van deze biomarkers naar een klinische toepassing. Daarnaast onderzochten we het gebruik van vroege economische evaluaties van medische technologie (*‘health technology assessment’*, HTA) in dit onderzoeksveld, en gaven we aan hoe deze technieken toegepast dienen te worden om de translatie van predictieve biomarkers te verbeteren. We concludeerden dat klinische translatie van predictieve biomarkers voor neoadjuvante chemotherapie bij borstkanker gebrekkig is. We beschreven biologische en klinische oorzaken die daaraan ten grondslag kunnen liggen, bijv. de aanwezigheid van heterogeniteit binnen de kenmerken van borstkanker en de hoge eisen die gesteld worden aan de studieopzet om klinische ‘*utility*’ aan te tonen. Een vroege HTA kan nuttig zijn bij de besluitvorming tijdens het ontwikkelingsproces van de biomarker. Bijv. bij het kiezen van een optimale studieopzet gegeven aanwezige middelen (door middel van ‘*multi criteria decision analysis*’) of het schatten van de kosteneffectiviteit van de testkarakteristieken van een bepaalde biomarker test (door middel van CEA).

In hoofdstuk 3 en 4 onderzochten we twee biomarker testen die voorspellend lijken te zijn voor hoge dosis alkyliserende chemotherapie in hormoon-receptor-negatieve, HER2-negatieve borstkanker (*‘triple negatief’*: de BRCA1-like status en BRCA1-like status gecombineerd met XIST en 53BP1 status. In hoofdstuk 3 ontwikkelden we een vroeg kosteneffectiviteitsmodel

dat de klinische toepassing van BRCA1-like status heeft gemeten met Multiplex Ligation Probe dependent Amplification (MLPA). Dit model liet zien dat het toepassen van de test met de huidige testkarakteristieken nog niet kosteneffectief is. De minimale prevalentie en positief voorspellende waarde van de test om kosteneffectief te zijn, schatten wij respectievelijk op 58.5% en 73.0 %. In hoofdstuk 4 breidden we het kosteneffectiviteits model van BRCA1-like status uit met XIST en 53BP1. We vergeleken de volgende biomarker combinaties: 1) BRCA1-like gemeten met MLPA; 2) BRCA1-like gemeten met *array Comparative Genomic Hybridisation* (aCGH); 3) strategie 1 gecombineerd met XIST en 53BP1; 4) strategie 2 gecombineerd met XIST en 53BP1. Op basis van dit model concludeerden we dat, gebaseerd op de huidige resultaten, het niet mogelijk is een biomarker-strategie te onderscheiden die meer kosten-effectief is dan de huidige klinische praktijk. Vervolgens hebben we een VOI analyse uitgevoerd, waaruit bleek dat het de moeite waard is om vervolgonderzoek te doen met een kostenplafond van 639 miljoen euro om de meest kosten-effectieve strategie te identificeren. De benodigde informatie kan het beste verzameld worden door vier zijstudies met een totale steekproefgrootte van 3000 patiënten te doen in een reeds lopende gerandomiseerde gecontroleerde studies (zoals NCT01057069). In deze retrospectieve studies moeten gegevens worden verzameld over: 1) de prevalentie van BRCA1-like borstkanker, de positief voorspellende waarde en de responspercentages van behandeling in biomarker-negatieve patiënten (MLPA niet-BRCA1-like), en de responsepercentages in de hele triple negatieve borstkankerpopulatie; 2) dezelfde parameters als in strategie 1 maar BRCA1-like status bepaald met aCGH, en de MLPA en aCGH BRCA1-like status gecombineerd met XIST en 53BP1 status; 3) kosten; 4) utiliteiten.

Beeldvormende technieken: monitoren van systemische anti-kankerbehandeling

In **hoofdstuk 5** beschrijven we een systematische literatuurreview over de prestaties van beeldvorming om de respons op neoadjuvante chemotherapie te monitoren per borstkanker subtype. We concludeerden dat er te weinig bewijs is om subtype-specifieke aanbevelingen te doen voor het monitoren van neoadjuvante chemotherapie met beeldvorming. Het is nodig om consensus te bereiken met betrekking tot de studieopzet, bijv. definities van “pathologisch Complete Response”, protocollen voor de uitvoering van beeldvorming en de tijdsintervallen tussen de start van de behandeling en het meten van de respons, voordat goed opgezette studies die een hoog niveau bewijs kunnen leveren worden gestart.

In **hoofdstuk 6 en 7** hebben we twee modellen gebouwd om in een vroeg stadium de kosteneffectiviteit te berekenen van twee nieuwe respons-gestuurde neoadjuvante chemotherapie interventies, waarbij neoadjuvante chemotherapie gedurende de behandeling aangepast wordt op basis van de respons gemeten met beeldvorming. In **hoofdstuk 6** berekenden we de kosteneffectiviteit van een scenario voor de behandeling van hormoon-receptor positieve borstkanker. Dit scenario startte met de behandeling van alle patiënten met twee kuren docetaxel,

doxorubicine en cyclophosphamide. Na het beoordelen van de respons middels echografie kregen de patiënten die reageerden op de therapie nog 6 kuren met hetzelfde therapieschema. Patiënten die niet reageerden op de eerste behandeling kregen vier kuren vinorelbine en capecitabine. De resultaten van de kosteneffectiviteitsanalyse laten zien dat deze manier van therapiemonitoring kosteneffectief is vergeleken met het niet monitoren van therapie. Vanuit klinisch oogpunt is het nodig een prospectieve validatie van dit scenario uit te voeren; i.e. het opzetten van een prospectieve studie. De vroege kosteneffectiviteitsanalyses kunnen hiervoor gebruikt worden om vervolgonderzoek te prioriseren, door het identificeren van parameters die het meest bijdragen aan de onzekerheid met betrekking tot het nemen van een beslissing (i.e., het wel of niet implementeren van de nieuwe beeldvormingsstrategie).

In **hoofdstuk 7** berekenden we de kosteneffectiviteit en benodigde investeringen om een ander respons-geleid neoadjuvante chemotherapie scenario te implementeren. Dit maal betrof het hormoon-receptor positieve, HER2 negatieve borstkankers. Dit scenario startte met de behandeling van alle patiënten met drie kuren dose dense doxorubicine en cyclophosphamide. Na het monitoren van de respons middels MRI ontvingen de patiënten met een respons op de behandeling nog drie kuren van hetzelfde schema, en werd het schema voor niet-reagerende patiënten aangepast naar drie kuren dose dense docetaxel en capecitabine. Het innovatieve aspect van deze studie is dat de uitkomsten werden berekend voor een huidig scenario en voor een scenario bij invoering van deze interventie in heel Nederland. Deze toevoeging is belangrijk omdat de overstap naar een nieuwe technologie bij verschillende artsen wisselend verloopt, wat de kosteneffectiviteit kan beïnvloeden. De conclusie van deze studie is dat, gebaseerd op de huidige gegevens, respons-geleide neoadjuvante chemotherapie in beide scenario's kosteneffectief is. Dit betekent dat respons-geleide neoadjuvante chemotherapie goedkoper en effectiever is dan conventionele chemotherapie (i.e., zonder beeldvorming) ongeacht hoe snel de adoptie van de nieuwe techniek verloopt. Wat betreft investeringen in het onderzoek concludeerden we dat Nederland voldoende personeel en MRI capaciteit heeft om het scenario volledig te implementeren.

Beeldvormende techniek: screenen voor afstandsmetastasen

In **hoofdstuk 8** berekenden we de kosteneffectiviteit van het screenen voor afstandsmetastasen middels PET/CT in stadium II/III borstkanker patiënten van de vier grote borstkanker subtypes in drie landen, namelijk Nederland, Groot Brittannië en de Verenigde Staten (VS). We concludeerden dat PET/CT met hoge zekerheid kosteneffectief is in HER2-negatieve patiënten indien zij behandeld worden in de VS. De verklaring voor dit resultaat is dat de kosten voor de palliatieve behandeling in dit subtype hoger zijn de fout-positieve dan in de terecht-positieve patienten. PET/CT verhoogt het terecht-positieve percentage en verlaagt het fout-positieve percentage wat resulteert in een kostenbesparing. De kosteneffectiviteit van PET/CT in Groot Brittannie en Nederland kan bereikt

worden door het verlagen van de kosten van PET/CT en door het verlagen van de kosten van de palliatieve behandeling.

Samenvattend draagt dit proefschrift bij aan de literatuur met betrekking tot vroegtijdig toegepaste kosteneffectiviteitsanalyses, '*value of information*' analyses en '*resource modelling*' analyses. Hiertoe gebruikten we case studies waarin nieuwe interventies van gepersonaliseerde behandeling van borstkanker werden onderzocht. De uitkomsten van deze studies informeren onderzoekers over: 1) de kans dat de interventie op basis van het huidige bewijs kosteneffectief is (hoofdstuk 3,4,6,7,8); 2) de ontwikkelingsdoelen om de interventie kosteneffectief te maken (hoofdstuk 3); 3) het type onderzoek dat nodig is om de interventie kosteneffectief te maken (hoofdstuk 4 en 6); 4) de investeringen die nodig zijn om de interventie te implementeren (hoofdstuk 7); 5) de huidige stand van zaken van predictieve biomarkers voor neoadjuvante chemotherapie bij borstkanker en respons-geleide neoadjuvante chemotherapie (hoofdstuk 2 en 5); en 6) het nut van vroege HTA methoden bij beleidsbeslissingen tijdens het ontwikkelen van een biomarker (hoofdstuk 2).

Acknowledgements

There are many individuals who have contributed to the success of this thesis.

First and foremost, my gratitude goes to prof. dr. Wim van Harten for allowing me to perform my doctoral thesis under his supervision. During the course of my PhD you have taught me a lot. From you I learned to be more assertive, more confident of my own ideas, and to not give up in adversities. Thank you for being such an inspiring and supportive supervisor.

Secondly, I would like to express deep gratitude to dr. Lotte Steuten who has taught me so much about health economics. You have always been supportive and a great problem solver in challenging situations. Despite your transfer overseas (to the Fred Hutchinson Cancer Research Center), you have shown continuous commitment to the project. Without your expertise this thesis would certainly have been more trying.

Special thanks to prof. dr. Sjoerd Rodenhuis for sharing his excellent expertise in breast oncology. I would also like to thank my PhD committee members, including prof. dr. René Medema, prof. dr. Sabine Linn and prof. dr. Floor van Leeuwen for their time and valuable comments.

The results of this thesis would have not been possible without close collaboration with several colleagues. Deep gratitude goes to dr. Valesca Retèl in whom I could find inspiration and with whom I had very fruitful discussions, to dr. Bianca Lederer and prof. dr. von Minckwitz for sharing their valuable data of the GeparTrio trial, to Lisanne Rigter and Suzana Teixeira, who invested time in helping me construct realistic cost-effectiveness models, and to Melanie Lindenberg for being such an enthusiastic, fun and hard-working companion. Last but certainly not least, I would like to thank Philip Schouten for teaching me the real-life struggles of predictive biomarker research, and for being the greatest companion in life.

My gratitude also goes to those that helped in the successful completion of my thesis: prof. dr. Sabine Linn, dr. Esther Lips, dr. Petra Nederlof, dr. Valdés Olmos, prof. dr. Emiel Rutgers, Mirjam Franken, dr. Vincent van der Noort, dr. Gabe Sonke, dr. Marcel Stokkel and dr. Jelle Wesseling. Some projects did not end as chapters for this thesis. Nonetheless, I would like to thank the people that invested time in them: dr. Kenneth Pengel, dr. Kenneth Gilhuijs, dr. Marie-Jeanne Vrancken Peeters, prof. dr. Ruud Pijnapple, Claudette Loo and Erik van Werkhoven.

Also thanks to Jorrita Tuurenhout and Marianne Brocken for smoothening this journey. You were always supportive and available for us (PhD students). Thanks to my close colleagues from the PSOE department and from the Wim van Harten Research group: Wim, Wilma, Valesca, Abi,

Melanie, Anke, Bruno, Ann-Jean, Laura, Miranda, Heleen and Willem. With you I shared great laughs – and once in a while frustrations. Not less important is my appreciation to all people who I shared a beer with during the research Friday borrels. Thanks to you this process has been more fun!

A special thanks goes to Jacobien and Lisanne for being my paranympths. My days in the NKI would have been so boring without you! I loved our morning coffees, our non-existing lunches, and of course, our borrels. We have shared confidences and supported each other, but most of all, we have had a lot of fun. You have being super collaborative during the preparation of this thesis and the organization of my defense party. Sharing it with you has made it way more exciting.

My special gratitude goes to those working relations that grew into friendships: Jacobien, Lisanne, Hellen, Wilma, Rita, Daniela and Rui. The best times during these PhD years were with you guys. I hope we keep on collecting many more! A big thanks to my oldest friends from high school and university. Although the distance has prevented us to meet as often as we would like to, I have enjoyed the extremely fun and intense reunions throughout Europe. Another thanks goes to my family in-law. Thank you so much for welcoming me in the family and for the affection that one needs when living abroad.

My most special acknowledgments go to my (step-)parents. You have always been my biggest support and have encouraged me to follow my dreams, despite the distance. Thank you for loving me unconditionally (*Els agraïments més especials van als meus pares (i padrastres). Sempre heu estat el meu gran suport. Sempre m'heu recolzat perquè fes allò que és millor per a mi, encara que això representi viure separats. Gràcies per estimar-me incondicionalment*).

Last, I would like to dedicate this thesis to my granddads, who are no longer with us. I know that they would be endlessly proud of my achievement (*Per acabar, m'agradaria dedicar aquesta tesi al padrí, a l'avi i al Josep. Sé que tots tres estarien molt orgullosos de veure on he arribat*).

Anna

April 1st, 2016

List of publications included in this thesis

Miquel-Cases A & Schouten PC, Steuten LMG, Retèl VP, Linn S, van Harten WH. **(Very) early health technology assessment and translation of predictive biomarkers in breast cancer.**

Submitted for publication

Miquel-Cases A, Steuten LMG, Retèl VP, van Harten WH. **Early stage cost-effectiveness analysis of a BRCA1-like test to detect triple negative breast cancers responsive to high dose alkylating chemotherapy.**

The Breast. 2015 Aug;24(4):397-405.

Received the “Best new investigator podium presentation” award at the annual congress of the International Society for Pharmacoeconomics and Outcomes Research. 2014 Amsterdam.

Miquel-Cases A, Retèl VP, van Harten WH, Steuten LMG. **Decisions on further research for predictive biomarkers of high dose alkylating chemotherapy in triple negative breast cancer: A value of information analysis.**

Value in Health 2016, in press.

Presented at the annual congress of the International Society for Pharmacoeconomics and Outcomes Research. 2014 Amsterdam

Lindenberg M, Miquel-Cases A, Retèl VP, Sonke G, Stokkel M, Wesseling J, van Harten WH. **Imaging performance in guiding response to neoadjuvant therapy according to breast cancer subtypes: A systematic literature review**

Submitted for publication

Miquel-Cases A, Retèl VP, Lederer V, von Minckwitz G, Steuten LMG, van Harten WH. **Exploratory cost-effectiveness analysis of response-guided neoadjuvant chemotherapy for hormone positive breast cancer patients.**

Accepted with minor revisions

Miquel-Cases A, Steuten LMG, Rigner LS, van Harten WH. **Cost-effectiveness and resource use of implementing MRI-guided NACT in ER-positive/HER2-negative breast cancers.**

Revised submission

Presented at the annual congress of the International Society for Pharmacoeconomics and Outcomes Research. 2015 Milan.

Miquel-Cases A & Teixeira S, Retèl VP, Steuten LMG, Valdés Olmos RA, Rutgers EJT & van Harten WH. **¹⁸F-FDG-PET/CT for distant metastasis screening in stage II/III breast cancer patients: A cost-effectiveness analysis from a British, US and Dutch perspective.**

Submitted for publication

Received the “*Best new investigator podium presentation*” award at the annual congress of the International Society for Pharmacoeconomics and Outcomes Research. 2015 Milan.

Curriculum vitae

Anna Miquel-Cases was born on December 15, 1987 in Igualada, Barcelona (Spain). She completed a Bachelor and a Master's degree in Pharmacy at the Universitat of Barcelona, from which she graduated in 2010. During her Master's degree she took part in an European Erasmus program in the University of Leiden, where she coursed a Science Based Business course that stimulated her interest towards the managerial side of health-care. After pursuing an internship as a community pharmacist in Barcelona, she moved to Rotterdam where she started a second Masters on 'Health economics, policy and law' at the Erasmus University in Rotterdam. She graduated in 2011, and in that same year, she started her PhD research in the Netherlands Cancer Institute (NKI-AVL) in Amsterdam (supervised by prof. Dr. Wim van Harten) in collaboration with the University of Twente in Enschede (co-supervised by dr. Lotte M Steuten). Her thesis was part of the Center for Translational Molecular Medicine (CTMM) project and focused on performing early cost-effectiveness analysis to emerging technologies to personalize breast cancer treatment.