fluorocitrate when it chelates to the enzyme-bound ferrous iron in a manner similar to that of isocitrate.

The inactivation of aconitase with time is more complicated. It has been suggested that it is possibly due to alkylation at the active site following, for instance, cleavage of the C–F bond (Kun, 1969; Carrell *et al.*, 1970). Alternatively the central carboxyl group of fluorocitrate may hydrogen-bond to the hydrogen abstracting site. Further biochemical studies on this problem and on the effect of fluorocitrate on the tri-carboxylate carrier of mitochondria (Eanes *et al.*, 1972) are now in progress in various laboratories.

We are grateful to Professor Jerry Donohue for the use of the automatic diffractometer at the University of Pennsylvania, Philadelphia, Pa. We also thank Miss Carol Ann Casciato and Miss Joyce Dargay for technical assistance. The enzymatic assay was performed by Dr Joseph J. Villafranca.

This research was supported by U.S.P.H.S. grants CA-10925, CA-06927 and RR-05539 from the National Institutes of Health, and by an appropriation from the Commonwealth of Pennsylvania.

References

- BUSING, W. R., MARTIN, K. O. & LEVY, H. A. (1962). *ORFLS*. Report ORNL-TM-305, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- CAHN, R. S., INGOLD, C. & PRELOG, V. (1966). *Angew. Chem. Int. Ed.* **5**, 385–445.
- CARRELL, H. L., GLUSKER, J. P., VILLAFRANCA, J. J., MILDVAN, A. S., DUMMEL, R. J. & KUN, E. (1970). *Science*, **170**, 1412–1414.
- CROMER, D. T. & LIBERMAN, D. (1970). *J. Chem. Phys.* **53**, 1891–1898.
- DUMMEL, R. J. & KUN, E. (1969). *J. Biol. Chem.* **244**, 2966–2969.
- EANES, R. Z., SKILLETER, D. N. & KUN, E. (1972). *Biochem. Biophys. Res. Commun.* **46**, 1618–1622.
- FANSHIER, D. W., GOTTFELD, L. K. & KUN, E. (1964). *J. Biol. Chem.* **239**, 425–434.
- GANTZEL, P. K., SPARKS, R. A., LONG, R. E. & TRUEBLOOD, K. N. (1965). *UCLALS4* Program in Fortran IV.
- GLUSKER, J. P. (1968). *J. Mol. Biol.* **38**, 149–162.
- HARRIS, D. R. (1966). *BDLSQ* Program No. 398 in *World List of Crystallographic Computer Programs*, 2nd ed. Edited by D. P. SHOEMAKER. Utrecht: Oosthoek.
- International Tables for X-ray Crystallography* (1962). Vol. III, pp. 202–207. Birmingham: Kynoch Press.
- JOHNSON, C. K. (1965). *ORTEP*. Report ORNL-3794, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- KUN, E. (1969). In *Citric Acid Cycle. Control and Compartmentation*. Edited by J. M. LOWENSTEIN, chap. 6, p. 318. New York: Dekker.
- LOVE, W. E. & PATTERSON, A. L. (1960). *Acta Cryst.* **13**, 426–428.
- PETERS, R. (1954). *Endeavour*, **13**, 147–154.
- STEWART, J. M., KUNDELL, F. A. & BALDWIN, J. C. (1970). *The X-ray System of Crystallographic Programs*. Version of July 1970, updated.
- STEWART, R. F., DAVIDSON, E. R. & SIMPSON, W. T. (1965). *J. Chem. Phys.* **42**, 3175–3187.
- VILLAFRANCA, J. J. & MILDVAN, A. S. (1971). *J. Biol. Chem.* **246**, 772–779.

Acta Cryst. (1973). **B29**, 682

The Crystal Structure of Lithium Formate Monohydrate

BY A. ENDERS-BEUMER AND S. HARKEMA

Chemical Physics Laboratory, Twente University of Technology, P. O. Box 217, Enschede, The Netherlands

(Received 7 July 1972; accepted 14 November 1972)

The structure of $\text{LiHCOO} \cdot \text{H}_2\text{O}$ has been determined. The space group is $Pbn2_1$ with 4 molecules per unit cell. 1027 Mo $K\alpha$ reflexions gave a final R value of 4.9%. All hydrogen atoms were found.

Introduction

Though crystallographic data for lithium formate monohydrate were published long ago (Nitta, 1928), the structure was not known. Raman studies with polarized light (Cadène, 1970) suggested a tetrahedrally coordinated Li ion and formate ions in planes parallel to (100), forming hydrogen bonds 2.74 Å long. This is confirmed by the present work.

Experimental

Lithium formate was prepared by neutralizing a solution of formic acid in water with lithium carbonate.

After several days, long transparent needles separated from the solution. By cutting and etching with water a crystal approximately $0.3 \times 0.3 \times 0.4$ mm was obtained and sealed in a Lindemann-glass capillary tube to protect it from moisture. Data were collected on a Nonius automatic diffractometer at 20°C, using Zr-filtered Mo $K\alpha$ radiation. 1027 reflexions with $\theta < 40^\circ$ were measured using the $\theta/2\theta$ scanning technique; 166 had an intensity smaller than twice the standard deviation as estimated from counting statistics. The crystal was rotated about [001].

The systematic absences were: $0kl$ with $k = 2n + 1$; $h0l$ with $h + l = 2n + 1$, indicating the space group

All U values are given in Å². Standard deviations in the last digits are shown in parentheses.

Values are $\times 10^4$. The anisotropic temperature factor used was

$$\exp[-2\pi^2(h^2a^{*2}U_{11}+k^2b^{*2}U_{22}+l^2c^{*2}U_{33}+2hka^*b^*U_{12}+2hla^*c^*U_{13}+2klb^*c^*U_{23})].$$

	X/a	Y/b	Z/c	U_{11}	U_{22}	U_{33}	U_{12}	U_{13}	U_{23}
C	3876 (2)	3046 (1)	172 (3)	375 (6)	222 (4)	223 (4)	-26 (4)	-22 (5)	-26 (4)
O(1)	4001	2920	2719	579	231	222	-3	-1	27
O(2)	4220	4098	-1145	385	291	217	-61	-41	44
O(3)	3893	340	4500 (0)	328	259	306	-6	-21	43
Li	3844 (3)	4509 (2)	5012 (5)	346 (8)	268 (8)	236 (9)	11 (7)	-18 (8)	26 (7)

(b) Hydrogen atoms

Values are $\times 10^3$, except for those for B , which are $\times 1$.

The isotropic temperature factor was

$$\exp(-8\pi^2 U \sin^2 \theta / \lambda^2) = \exp(-B \sin^2 \theta / \lambda^2).$$

	X/a	Y/b	Z/c	U	B
H(1)	346 (2)	232 (2)	-71	41 (4)	3.2 (3)
H(2)	395	119	413	56 (5)	4.4 (4)
H(3)	454	30	578	31 (4)	2.4 (3)

A three-dimensional sharpened Patterson synthesis gave the coordinates of the carbon and oxygen atoms. After several cycles of refinement of positional and

Table 2. *Observed and calculated structure factors ($\times 10$)*[illegible]

isotropic temperature parameters a difference Fourier synthesis revealed the position of the Li ion. After further refinement with anisotropic temperature factors a second difference-Fourier map was calculated which showed the positions of the H atoms. Refinement was carried out with a modification of the program *ORFLS* (Busing, Martin & Levy, 1962). Atomic scattering factors for carbon and oxygen atoms were calculated (Harkema, 1971) from the atomic wave functions of Clementi (1965). The scattering factor for hydrogen was taken from Stewart, Davidson & Simpson (1965), and that for Li^+ from Hurst, Miller & Matsen (1958).

In the last stage of refinement a correction for secondary extinction was applied (Larson, 1969). As the linear absorption coefficient for Mo radiation was 1.55 cm^{-1} , no absorption correction was applied. The final *R* index was 4.9%. Positional and thermal parameters are given in Table 1. Structure factors are given in Table 2.

Discussion of the results

The most important bond distances and angles are given in Table 3.

Table 3. Bond distances (Å) and angles (°) with standard deviations in parentheses

C—O(1)	1.244 (3)
C—O(2)	1.248 (3)
C—H(1)	0.88 (3)
O(1)···O(3)	2.715 (3)
O(3)···O(3'')	2.897 (3)
O(1)—C—O(2)	125.5 (2)
O(1)—C—H(1)	115 (2)
O(2)—C—H(1)	120 (2)
O(3)—H(2)···O(1)	170 (3)
Li tetrahedron	
Li—O(2')	1.923 (4)
Li—O(2'')	1.954
Li—O(1)	1.938
Li—O(3''')	1.974

The lithium ion (Fig. 1) is tetrahedrally coordinated with 3 formate oxygens and one water oxygen. The angles vary between 104 and 112°, the Li—O distances between 1.923 and 1.974 Å, comparable with *e.g.* 1.89–2.04 Å in lithium acetate dihydrate (Galigné, Mouvet & Falgueirettes, 1970) and 1.935–2.076 Å in lithium oxalate (Beagley & Small, 1964). The water molecules form infinite zigzag chains parallel to the *c* axis, with hydrogen bonds of 2.897 Å, while the other water hydrogen atom forms a link with the formate ion. Muthukrishnan & Ramakrishna (1971) concluded from n.m.r. data that H(3) should be disordered, occupying two potential minima between 2 water oxygen atoms. Our data, however, indicate complete localization of H(3).

The formate group is probably planar: the distance of the H atom from the plane of the C and O atoms

Fig. 1. Arrangement of some molecules in the unit cell, showing the coordination of the Li ion and the water chain.

is 0.03 Å, comparable to the standard deviation. When our refinement was completed, we received the paper of Mohana Rao & Viswamitra (1971), who had also determined the structure of lithium formate monohydrate from 331 $\text{Cu K}\alpha$ reflexions with film methods. Their results agree reasonably with ours (though theirs are, of course, less accurate). Contrary to our results, however, they found a significant difference in the C—O bond lengths, C—O(1) = 1.213(12); C—O(2) = 1.279(12) Å, which they explained by the different environments of the two oxygen atoms. Indeed O(2) is coordinated to two Li ions, O(1) only to one, but it forms the hydrogen bond. The infrared (Harvey, Morrow & Shurvell, 1963; Donaldson, Knifton & Ross, 1964) and Raman (Ramanujam, 1969; Cadène, 1970) spectra show a slight splitting of some fundamentals, but the splitting is very small ($< 20 \text{ cm}^{-1}$) and each of the four spectra gives a different result. Our conclusion is that the effect of the hydrogen bond compensates for the loss of a Li coordination, so both bonds should be approximately equal.

The present investigation has been carried out under the auspices of the Foundation for Fundamental Research on Matter by Electron and X-rays (FOMRE) and with aid from the Netherlands Organization for the Advancement of Pure Research (ZWO).

References

- BEAGLEY, B. & SMALL, R. W. H. (1964). *Acta Cryst.* **17**, 783–788.
- BUSING, W. R., MARTIN, K. O. & LEVY, H. A. (1962). *ORFLS*. Report ORNL-TM-305, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A.
- CADÈNE, M. (1970). *C. R. Acad. Sci. Paris*, **B270**, 909–912.
- CLEMENTI, E. (1965). *IBM J. Res. Dev.* **9**, 2–19.
- DONALDSON, J. D., KNIFTON, J. F. & ROSS, S. D. (1964). *Spectrochim. Acta*, **20**, 847–851.
- GALIGNÉ, J. L., MOUVET, M. & FALGUEIRETTES, J. (1970). *Acta Cryst.* **B26**, 368–372.

- HARKEMA, S. (1971). Thesis, Twente University of Technology, Enschede, The Netherlands.
- HARVEY, K. B., MORROW, B. A. & SHURVELL, H. F. (1963). *Canad. J. Chem.* **41**, 1181–1187.
- HURST, R. P., MILLER, J. & MATSEN, F. A. (1958). *Acta Cryst.* **11**, 320–322.
- LARSON, A. C. (1969). In *Crystallographic Computing*, pp. 291–294. Edited by F. R. AHMED. Copenhagen: Munksgaard.
- MOHANA RAO, J. K. & VISWAMITRA, M. A. (1971). *Ferroelectrics*, **2**, 209–216.
- MUTHUKRISHNAN, K. & RAMAKRISHNA, J. (1971). *Curr. Sci.* **40**, 486–487.
- NITTA, I. (1928). *Sci. Pap. Inst. Phys. Chem. Res. Tokyo*, **9**, 151–163.
- RAMANUJAM, P. S. (1969). *Indian J. Pure Appl. Phys.* **7**, 420–423.
- STEWART, R. F., DAVIDSON, E. R. & SIMPSON, W. T. (1965). *J. Chem. Phys.* **42**, 3175–3187.

Acta Cryst. (1973). **B29**, 685

The Crystal Structure of 5-Bromo-12*S*-tetrahydroaustamide

BY J. COETZER AND P. S. STEYN

National Physical and National Chemical Research Laboratories, Council for Scientific and Industrial Research, P.O. Box 395, Pretoria, South Africa

(Received 13 November 1972; accepted 23 November 1972)

The crystal structure of 5-bromo-12*S*-tetrahydroaustamide has been determined by three-dimensional X-ray analysis. The crystals are monoclinic with space group *P*2₁, and two molecules in the unit cell of dimensions *a* = 7·241, *b* = 13·148, *c* = 11·556 Å and β = 106·68°. The absolute configuration of the molecule was derived from the known stereochemistry at the 12-position established in earlier chemical and spectroscopic studies.

Introduction

In our continuing studies on biologically active fungal metabolites, austamide (I) was isolated as a toxic component from *Aspergillus ustus* (Steyn, 1971).

Prolonged hydrogenation of I gave two diastereoisomeric tetrahydroaustamides IIa and IIb. Acid hydrolysis of the major product, IIa, liberated *S*-proline, therefore establishing the 12*S*-configuration at this newly created chiral centre. The stereochemistry at position 9 was based on the 9*S*, 12*S*-configuration of prolyl-2-(1'1'-dimethylallyl)tryptophyldiketopiperazine, an apparent bioprecursor of I. The configuration at the *spiro* atom, position 2 relative to that of position 9, was based on n.m.r. shift data (Steyn, to be published).

5-Bromo-12*S*-tetrahydroaustamide (IIc) was prepared for an X-ray crystallographic study in an attempt to substantiate the above conclusions.

Experimental

12*S*-Tetrahydroaustamide (50 mg) in tetrahydrofuran (2 ml) was treated with pyridinium hydrobromide perbromide (44 mg) in tetrahydrofuran (2 ml). The colourless mixture was concentrated in a stream of nitrogen and the residue partitioned between chloroform and water. The organic phase was concentrated and crystallized from ethanol to yield IIc. [Analysis: Found: C, 54·31; H, 5·60; Br, 17·25; N, 9·05%. C₂₁H₂₄BrN₃O₃·H₂O requires C, 54·63; H, 5·44; Br, 17·06; N, 8·97%].

The bright yellow rhombic crystals are monoclinic with space group *P*2₁ (0*kl*0 absent for *k* odd).

The other crystal data are:

Molecular formula C₂₁H₂₄BrN₃O₃
 $\lambda(\text{Mo } K\alpha) = 0\cdot70926 \text{ \AA}$

a = 7·241 (1) Å

b = 13·148 (2)

c = 11·556 (1)

β = 106·68 (30)°

*d*_x = 1·46 g cm⁻³

*d*_m = 1·46 (measured by flotation in a mixture of bromoform and methanol)

M.W. 446·34

Z = 2.

Unit-cell dimensions were calculated from the accurate spot positions measured on a Hilger & Watts four-circle automatic diffractometer using a least-