A Systematic Study to Neutral, Water Soluble Calix[4]arenes

Marcel H. B. Grote Gansey, Willem Verboom and David N. Reinhoudt*

Laboratory of Organic Chemistry, University of Twente, P. O. Box 217, 7500 AE
Enschede, The Netherlands

Abstract: A series of neutral water soluble calix[4]arenes was synthesized via direct chlorosulfonylation followed by reaction with hydroxyl group-containing amines. The solubility of these calix[4]arene sulfonamides in water, as determined by means of UV measurements, varies from $\sim 10^{-5}$ to 0.31 M.

Calix[4]arenes^{1,2} have become important hydrophobic building blocks in supramolecular chemistry. We have used (selective) functionalization³ of calix[4]arenes both at the phenolic OH groups (*lower rim*) and at the para positions of the phenol rings (*upper rim*) for the design of selective receptors for cations,⁴ anions,⁵ and neutral molecules.⁶ Subsequent application of these calix[4]arene based ionophores in ion sensors⁷ or ion transport through liquid membranes⁸ requires a high hydrophobicity. However, for *in vivo* application of appropriately functionalized calix[4]arenes water solubility is a prerequisite.⁹ Hitherto in most cases water soluble calixarenes have been obtained by the introduction of charged moieties. The first report of water soluble calixarenes by Ungaro *et al.*¹⁰ describes a *p-tert*-butylcalix[4]arene tetracarboxylic acid, of which the alkali and ammonium salts are soluble in water (concentrations between 5×10^{-4} and 5×10^{-3} M depending on the cation used). Shinkai *et al.*¹¹ reported the pK_a values of a water soluble *p*-tetrasulfonate tetrasodium calix[4]arene. *Lower rim* functionalized *upper rim* sulfonato calix[4]arenes were described by Casnati *et al.*¹² Other methods using charged species to achieve water solubility are the introduction of sulfonato groups at the *lower rim*,¹³ phosphonic acid groups¹⁴ or cationic trialkylammonium groups.¹⁵ To avoid unspecific binding of cations by anionic groups, or repulsion by cationic groups in the receptor molecule, we need **neutral**, water soluble, functionalized calix[4]arenes. To the best of our knowledge in literature only two neutral, water soluble calix[4]arenes have been described. Shinkai and Reinhoudt *et al.*¹⁶ reported the pK_a determination of

neutral, water soluble tetrakis[bis-(2-hydroxyethyl)-aminosulfonyl]calix[4]arene. Newkome *et al.*¹⁷ developed the *silvanols* with hydroxyl group-containing amides at the *upper rim*, of which the calix[4]arene derivative contains 36 hydroxyl groups. However, in both cases no quantitative data about the water solubilities are given.

Recently we reported that calix[4]arene sulfonamides can be easily obtained by chlorosulfonylation at the *upper rim* of calix[4]arenes, functionalized at the *lower rim*, followed by reaction with an appropriate amine.¹⁸ In this communication we present our preliminary results of a study towards water soluble calix[4]arenes by systematically increasing the number of hydroxyl groups in calix[4]arene sulfonamides.

As starting compounds we selected two different tetrakis(chlorosulfonylated)calix[4]arenes *viz.* the tetrakis(ethoxymethoxy)- (1) and the tetraethyl ester derivative (2).¹⁸ Reaction of 1 and 2 with ethanolamine in dichloromethane for 8 h gave the calix[4]arene sulfonamides 3a¹⁹ and 4a²⁰ both in 57% yield. From the corresponding reaction with diethanolamine compounds 3b²¹ and 4b²² were obtained in 42 and 49% yield, respectively. For solubility reasons the reactions of 1 and 2 with tris(hydroxymethyl)aminomethane were carried out in DMSO for 8 h to give compounds 3c²³ and 4c²⁴ in 56 and 42% yield, respectively. The ¹H-NMR spectra of all compounds 3 and 4, recorded in solvents more polar than CDCl₃, show singlets for the aromatic protons and one AB system for the protons of the methylene bridges, indicating that the substitution reaction of the chloro atoms of compounds 1 and 2 by the amines has taken place completely.

The water solubility of sulfonamides 3a-c and 4a-c was determined by means of UV measurements²⁶ (Table 1). These data clearly demonstrate that by variation of the number of hydroxyl groups the water solubility of calix[4]arenes can be dramatically altered. Introduction of one additional hydroxyl group per

aromatic unit of the calix[4]arene leads roughly to a hundred fold increase of the water solubility. This method may be useful for obtaining neutral, *lower rim* functionalized, water soluble calix[4]arenes that can be applied in biological systems.

Table 1: Water solubility of neutral calix[4]arenes at 25 °C

Compound	Water solubility (M)
3a	$\sim 10^{-5}$
3b	$(2.5 \pm 0.2) \times 10^{-3}$
3c	0.23 ± 0.02
4a	$\sim 10^{-5}$
4b	$(8.8 \pm 0.1) \times 10^{-4}$
4c	0.31 ± 0.02

These investigations are supported by the Netherlands Foundation for Chemical Research (SON) with financial aid from the Netherlands Technology Foundation (STW).

REFERENCES AND NOTES

1. Gutsche, C. D. *Calixarenes, Monographs in Supramolecular Chemistry*, Vol. 1; Stoddart, J. F., Ed.; The Royal Society of Chemistry; Cambridge, 1989.
2. *Calixarenes, a Versatile Class of Macrocyclic Compounds*; Vicens, J., Böhmer, V., Eds.; Kluwer Academic Publishers; Dordrecht, the Netherlands, 1991.
3. Van Loon, J.-D.; Verboom, W.; Reinhoudt, D. N. *Org. Prep. Proc. Int.* **1992**, *24*, 437.
4. Iwema Bakker, W. I.; Haas, M.; Khoo-Beattie, C.; Ostaszewski, R.; Franken, S. M.; Den Hertog, Jr., H. J.; Verboom, W.; De Zeeuw, D.; Harkema, S.; Reinhoudt, D. N. *J. Am. Chem. Soc.* **1994**, *116*, 123.
5. Rudkevich, D. M.; Verboom, W.; Reinhoudt, D. N. *J. Org. Chem.* **1994**, in press.
6. Van Loon, J.-D.; Heida, J. F.; Verboom, W.; Reinhoudt, D. N. *Recl. Trav. Chim. Pays-Bas* **1992**, *111*, 353.
7. Cobben, P. L. H. M.; Egberink, R. J. M.; Bomer, J. G.; Bergveld, P.; Verboom, W.; Reinhoudt, D. N. *J. Am. Chem. Soc.* **1992**, *114*, 10573.
8. Nijenhuis, W. F.; Buitenhuis, E. G.; De Jong, F.; Sudhölter, E. J. R.; Reinhoudt, D. N. *J. Am. Chem. Soc.* **1991**, *113*, 7963.
9. Iwema Bakker, W. I.; Haas, M.; Den Hertog, Jr., H. J.; Verboom, W.; De Zeeuw, D.; Bruins, A. P.; Reinhoudt, D. N. *J. Org. Chem.* **1994**, *59*, 972.
10. Arduini, A.; Pochini, A.; Reverberi, S.; Ungaro, R. *J. Chem. Soc., Chem. Commun.* **1984**, 981.
11. Shinkai, S.; Araki, K.; Koreishi, H.; Tsubaki, T.; Manabe, O. *Chem. Lett.* **1986**, 1351.
12. Casnati, A.; Ting, Y.; Berti, D.; Fabbi, M.; Pochini, A.; Ungaro, R.; Sciotto, D.; Lombardo, G. G.

- Tetrahedron* **1993**, *49*, 9815.
13. Shinkai, S.; Kawabata, H.; Arimura, T.; Matsuda, T.; Satoh, H.; Manabe, O. *J. Chem. Soc., Perkin Trans. 1* **1989**, 1073.
 14. Almi, M.; Arduini, A.; Casnati, A.; Pochini, A.; Ungaro, R. *Tetrahedron* **1989**, *45*, 2177.
 15. Shinkai, S.; Shirahama, Y.; Tsubaki, T.; Manabe, O. *J. Chem. Soc., Perkin Trans. 1* **1989**, 1859.
 16. Shinkai, S.; Araki, K.; Grootenhuys, P. D. J.; Reinhoudt, D. N. *J. Chem. Soc., Perkin Trans. 2* **1991**, 1883.
 17. Newkome, G. R.; Hu, Y.; Saunders, M. J.; Fronczek, F. R. *Tetrahedron Lett.* **1991**, *32*, 1133.
 18. Morzherin, Y.; Rudkevich, D. M.; Verboom, W.; Reinhoudt, D. N. *J. Org. Chem.* **1993**, *58*, 7602.
 19. m.p. 167-168 °C; ¹H-NMR (DMSO-*d*₆): δ 7.27 (s, 8H, ArH), 7.00 (t, 4H, NH), 4.54 (t, 4H, OH), 4.52 (part of ABq, 4H, ArCH₂Ar),²⁵ 4.16 (t, 8H, OCH₂), 3.79 (t, 8H, OCH₂), 3.5-3.4 (m, 12H, CH₂OH and part of ABq, 4H, ArCH₂Ar),²⁵ 3.31 (s, 12H, OCH₃), 2.56 (q, 8H, NCH₂); ¹³C-NMR (DMSO-*d*₆): δ 158.8 (s, ArC-O), 135.0 (s, ArC-SO₂), 134.4 (s, ArC-CH₂), 126.8 (d, ArC-H), 73.6 (t, OCH₂), 71.2 (t, OCH₂), 60.0 (t, CH₂OH), 58.0 (q, OCH₃), 44.9 (t, NC), 29.8 (t, ArCH₂Ar); FAB mass spectrum, *m/e* 1149.6 [(M + H)⁺, calcd 1149.3].
 20. m.p. 237-239 °C; ¹H-NMR (DMSO-*d*₆): δ 7.29 (s, 8H, ArH), 7.07 (bs, 4H, NH), 4.84 (s, 8H, OCH₂C(O)), 4.79 and 3.49 (ABq, 8H, *J* = 13.5 Hz, ArCH₂Ar), 4.57 (bs, 4H, OH), 4.15 (t, 8H, OCH₂Me), 3.35 (t, 8H, CH₂OH), 2.57 (bt, 8H, NCH₂), 1.22 (t, 12H, CH₃); ¹³C-NMR (DMSO-*d*₆): δ 169.1 (s, C(O)), 157.8 (s, ArC-O), 135.0 (s, ArC-SO₂), 134.5 (s, ArC-CH₂), 127.0 (d, ArC-H), 71.1 (t, OCH₂C(O)), 60.5 (t, OCH₂Me), 59.9 (t, CH₂OH), 44.9 (t, NC), 30.7 (t, ArCH₂Ar), 13.9 (q, CH₃); FAB mass spectrum, *m/e* 1261.7 [(M + H)⁺, calcd 1261.3].
 21. m.p. 215-216 °C; ¹H-NMR (DMSO-*d*₆): δ 7.33 (s, 8H, ArH), 4.80 (t, 8H, OH), 4.47 and 3.55 (ABq, 8H, *J* = 12.5 Hz, ArCH₂Ar), 4.16 (t, 8H, OCH₂), 3.80 (t, 8H, OCH₂), 3.48 (q, 16H, CH₂OH), 3.32 (s, 12H, OCH₃), 2.86 (t, 16H, NCH₂); ¹³C-NMR (DMSO-*d*₆): δ 159.0 (s, ArC-O), 135.1 (s, ArC-SO₂), 132.1 (s, ArC-CH₂), 127.3 (d, ArC-H), 73.6 (t, OCH₂), 71.1 (t, OCH₂), 60.1 (t, CH₂OH), 58.0 (q, OCH₃), 51.5 (t, NC), 30.0 (t, ArCH₂Ar); FAB mass spectrum, *m/e* 1325.4 [(M + H)⁺, calcd 1325.4].
 22. m.p. 211-213 °C; ¹H-NMR (DMSO-*d*₆): δ 7.36 (s, 8H, ArH), 4.86 (s, 8H, OCH₂C(O)), 4.78 (t, 8H, OH), 4.75 (ABq, 8H, *J* = 13.4 Hz, ArCH₂Ar), 4.16 (q, 8H, OCH₂Me), 3.48 (bt, 16H, CH₂OH), 2.86 (bt, 16H, NCH₂), 1.22 (t, 12H, CH₃); ¹³C-NMR (DMSO-*d*₆): δ 169.0 (s, C(O)), 158.1 (s, ArC-O), 134.8 (s, ArC-SO₂), 132.7 (s, ArC-CH₂), 127.6 (d, ArC-H), 71.0 (t, OCH₂C(O)), 60.5 (t, OCH₂Me), 60.1 (t, CH₂OH), 51.5 (t, NC), 30.5 (t, ArCH₂Ar), 13.9 (q, CH₃); FAB mass spectrum, *m/e* 1438.6 [(M + H)⁺, calcd 1438.7].
 23. m.p. 153-155 °C; ¹H-NMR (D₂O): δ 7.29 (s, 8H, ArH), 4.51 and 3.44 (ABq, 8H, *J* = 13.3 Hz, ArCH₂Ar), 4.25 (t, 8H, OCH₂), 3.92 (t, 8H, OCH₂), 3.68 (s, 24H, CH₂OH), 3.38 (s, 12H, OCH₃); ¹³C-NMR (D₂O): δ 158.5 (s, ArC-O), 137.3 (s, ArC-SO₂), 135.1 (s, ArC-CH₂), 126.1 (d, ArC-H), 73.2 (t, OCH₂), 72.1 (t, OCH₂), 61.7 (t, CH₂OH), 59.8 (s, NC), 58.1 (q, OCH₃), 30.8 (t, ArCH₂Ar); FAB mass spectrum, *m/e* 1099.4 [(M - SO₂NHC(CH₂OH)₃ - C(CH₂OH)₃)⁺, calcd 1099.3].
 24. m.p. 166-168 °C; ¹H-NMR (D₂O): δ 7.32 (s, 8H, ArH), 4.90 (s, 8H, OCH₂C(O)), 4.72 and 3.49 (ABq, 8H, *J* = 13.6 Hz, ArCH₂Ar), 4.22 (q, 8H, OCH₂Me), 3.68 (s, 24H, CH₂OH), 1.26 (t, 12H, CH₃); ¹³C-NMR (D₂O): δ 172.0 (s, C(O)), 157.4 (s, ArC-O), 138.1 (s, ArC-SO₂), 134.7 (s, ArC-CH₂), 126.5 (d, ArC-H), 71.6 (t, OCH₂C(O)), 62.4 (t, OCH₂Me), 61.7 (s, NC), 59.6 (t, CH₂OH), 31.3 (t, ArCH₂Ar), 13.6 (q, CH₃); FAB mass spectrum, *m/e* 1133.4 [(M - 2 x SO₂NHC(CH₂OH)₃ + H)⁺, calcd 1133.3].
 25. The coupling constant could not be determined due to overlap of signals.
 26. Solutions of the compounds in doubly distilled water were measured at 25 °C. The concentrations were chosen to give absorbances between 0.2 and 0.8 and a plot was drawn of the absorbance against the concentration. A saturated solution was filtered and diluted with a known amount of water to give an absorbance in the same region and the concentration of the saturated solution could be calculated using the plot.

(Received in UK 13 July 1994; accepted 29 July 1994)