Synthesis of new macrocyclic ligands for hetero-multinuclear transition-metal complexes

Wilhelm T.S. Huck, Frank C.J.M. van Veggel and David N. Reinhoudt *

Laboratory of Organic Chemistry, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands

(Received November 4, 1994)

Abstract. The [3 + 3] macrocyclic compounds **2a–2c** were easily prepared by cyclization of the dialdehyde **8** with the appropriate diamines in refluxing CH₃CN/MeOH (1/1), in the presence of Ba²⁺ as a template ion. These macrocycles contain three salen units and are therefore, in combination with the crown ether interior, potentially tetranucleating. The length of the diamine bridge and the temperature at which the reaction is carried out, appear to influence the size of the macrocycle formed. In this way, the [4 + 4] macrocycle **2d** could be formed selectively over the [3 + 3] macrocycles by carrying out the reaction at room temperature.

Introduction

Schiff-base-containing macrocyclic ligands designed for complexing more than one transition metal, are widely used in coordination chemistry^{1–4}. The distinctive properties of these multinuclear complexes are related to the close proximity of two or more transition metal cations. This is of theoretical interest because of metal-metal interactions⁵, and also plays an important role when the complexes are used in studies as enzyme mimics or as homogeneous catalysts. The metals can function as multiple coordination sites for substrate binding and/or activation. Complexes containing two different transition metals with a short intermetallic distance, have interesting redox properties⁶. Furthermore, Floriani et al.⁷ have shown that an alkaline cation (Lewis acid) can assist in the activation and binding of small substrate molecules, like CO₂, to a Co(I) center (Lewis base) by stabilizing the CO₂[–] intermediate. Previously we have reported on hetero-di- and -trinuclear complexes combining transition metal cations and alkali-metal or alkaline-earth-metal cations⁸ (Chart 1).

These macrocyclic ligands contain a salen unit which forms stable transition-metal complexes⁹, and a crown-ether unit which complexes various alkali- or alkaline-earth-metal cations¹⁰. In these hetero-multinuclear complexes it is possible to tune the redox behavior of the transition metals by co-complexation of various 'hard' cations¹¹.

The electrocatalytic properties of the Ni cation in complexes **1a** and **1b** are influenced by co-complexation of the Ba²⁺ cation. The Ni/Ba complex converts benzyl chloride and acetic anhydride into 1-phenylpropan-2-one (11%), whereas the Ni complex does not show any catalytic activity¹².

In this paper we describe an extension to a tetranucleating macrocycle by combining the 'hard' polyether cavity with three 'soft' salen units. We report the synthesis of a macrocyclic ligand, in which six phenolic OH groups are

Chart 1.

bridged intramolecularly by hydrogen bonds to the imine nitrogens¹³. In this way the oxygens resemble the arrangement in 18-crown-6 with the ethylene bridges replaced by salen units (Chart 2).

Results and discussion

The synthesis of the dialdehyde **8**, required for the macrocyclization with an appropriate diamine, is depicted in Scheme 1^{14,15}. Firstly, naphthalene-2,3-diol was diallylated with allyl bromide in acetone, using K₂CO₃ as a base¹⁶. Subsequently a Claisen rearrangement at 170°C of the 2,3-diallyloxynaphthalene gave the 1,4-di(prop-1-enyl)naphthalenediol. The product was not isolated but immediately reacted with acetyl chloride, in order to prevent oxidation reactions at the phenolic OH groups in the subsequent steps. Isomerization of the double bonds to the 2-alkenes was carried out using RhCl₃ as catalyst in THF/EtOH at 50°C¹⁷. After this isomerization, ozonolysis was carried out in CH₂Cl₂ at –70°C, followed by reductive work-up with dimethyl sulfide to give the dialde-

Chart 2.

hyde 7 in 90% yield. Deprotection of the phenolic OH groups gave 2,3-dihydroxy-1,4-naphthalenedialdehyde. The macrocyclic ligand was prepared by a [3 + 3] macro-

cyclic condensation reaction of dialdehyde 8 and several diamines, using Ba^{2+} as a template cation. The macrocyclization with diamines 9a–9c, in the presence of 0.33 mol% of $\text{Ba}(\text{CF}_3\text{SO}_3)_2$, gave the highly colored [3 + 3] macrocycles 2a–2c in approximately 70% yield (Scheme 2). $\text{Cs}(\text{CF}_3\text{SO}_3)$ and $\text{Pb}(\text{ClO}_4)_2$ did not produce cyclic products, but, when the Ba^{2+} cation was used, a macrocyclic product was formed. The role of the template cation was shown unambiguously by the following control experiment. When the dialdehyde 8 and an appropriate diamine were refluxed in the absence of $\text{Ba}(\text{CF}_3\text{SO}_3)_2$, no cyclic products were obtained according to FAB-MS, and only a high molecular weight residue was formed. When this was redissolved and refluxed again in the presence of the Ba^{2+} salt, a cyclic product was formed, albeit in somewhat lower yields than in the templated reaction of dialdehyde and diamine.

Compounds 2 were characterized by ^1H - and ^{13}C -NMR, fast atom bombardment (FAB) MS and Fourier-Transform (FT) IR spectroscopy. The simplicity of the ^1H NMR spectrum of macrocycle 2a is a good indication that a macrocycle is formed. Because of the high symmetry, all phenolic and imine protons are equivalent and give only one singlet each. The ^1H -NMR spectrum of 2b shows two

Scheme 1.

Scheme 2.

imine protons at room temperature (δ 9.35 and 9.38 ppm). This means that this macrocycle is not completely planar. When the spectrum is recorded at 75°C, the two signals coalesce at δ 9.36 ppm. In the FT-IR spectra the imine C=N stretch frequency for all macrocycles was present at approximately 1620 cm^{-1} . In addition, FAB-MS shows only peaks corresponding to the molecular ion peak (M^+) and the subsequent loss of one or two counter-anions (CF_3SO_3^-). The largest intensity in the positive FAB-MS spectrum is the $(M-\text{CF}_3\text{SO}_3)^+$ cation, but in the negative FAB-MS there is a small peak corresponding to M^- . With propane-1,3-diamine as the diamine, a mixture of [3 + 3] and [4 + 4] macrocycles was formed. When the reaction was carried out at room temperature instead of reflux temperature, the [4 + 4] macrocycle formed selectively over the [3 + 3] product¹⁸. In the FAB-MS spectrum there were also peaks present that could be attributed to a [2 + 2] macrocycle. The [4 + 4] macrocycle **2d** was purified by column chromatography. The ^1H -NMR spectrum shows a highly symmetric pattern with two imine protons at δ 9.90 and 9.84 ppm which indicate some deviation from planarity.

Preliminary FAB-MS results indicate that, with transition metals like Cu(II), Ni(II) and Zn(II), tetranuclear complexes are formed. The FT-IR spectra show a shift in the imine C=N stretch to lower wavelengths, indicating complexation of the transition metals in the salen moiety¹⁹. Plasma-desorption (PD) and electrospray-ionisation (EI) MS indicate intermolecular stacking of the macrocyclic complexes.

Conclusions

Macrocyclic ligands are easily formed when dialdehyde **8** is refluxed with various diamines in the presence of Ba^{2+} as a template ion. The macrocycles **2a–2c** are made up of three dialdehydes and three diamines. Lowering the temperature and using propane-1,3-diamine resulted in a [4 + 4] instead of a [3 + 3] macrocyclization. Preliminary FAB-MS data show that the corresponding transition metal complexes can be formed.

Experimental

General methods

NMR spectra were recorded on a Bruker AC 250 spectrometer in CDCl_3 , unless otherwise stated. The residual protons of the solvents were used as internal standard. Mass spectra were obtained with a Finnigan MAT 90 spectrometer. FAB-MS spectra were obtained with 3-nitrobenzyl alcohol as a matrix. IR spectra were recorded with a Biorad Digilab FTS 60 IR-spectrometer. Melting points were determined using a Reichert melting point apparatus and are uncorrected. Elemental analyses were carried out with a Model 1106 Carlo Erba Strumentazione elemental analyzer. CH_2Cl_2 was distilled over CaCl_2 before use. CH_3CN (p.a., Merck) was stored over molecular sieves (4 Å) prior to use. THF was distilled from sodium/benzophenone. Other chemicals were of reagent grade and used without further purification. All reactions were carried out under an argon atmosphere.

1,4-Bis(prop-2-enyl)-2,3-diacetoxynaphthalene (**5**)

2,3-Bis(prop-2-enyloxy)naphthalene (100.0 g, 0.417 mol) was heated neat at 170°C for 1 h. When the rearrangement was complete on NMR, the temperature was lowered to room temperature and 500 ml of CH_2Cl_2 and 100 ml of Et_3N were added. At 0°C, acetyl chloride (76.5 g, 1.0 mol) was added over a period of 1 h. After stirring overnight, the reaction mixture was washed twice with H_2O and dried over MgSO_4 . Removal of the solvent and trituration with MeOH yielded 95.1 g (71%) of pure **5**; m.p. 128–129°C. ^1H -NMR: δ 8.15, dd, 2H, J 3.2 Hz, H6 and H7; 7.05, dd, 2H, J 3.2 Hz, H5 and

H8; 6.1–5.9, m, 2H, =CH; 5.1–5.0, m, 4H, =CH₂; 3.73, d, 4H, J 5.9 Hz, ArCH_2 ; 2.35, s, 6H, CH_3 . ^{13}C -NMR: δ 168.5, C=O; 139.5, ArO; 135.2, =CH₂; 131.1, 127.0, Ar; 126.0, 125.0, Ar; 30.7, ArCH_2 . IR (KBr): 1770, C=O, cm^{-1} . MS (EI): m/z 324.1, M^+ , calcd. 324.1. Anal. calcd. for $\text{C}_{20}\text{H}_{20}\text{O}_4$: C 74.06, H 6.21; found C 73.98, H 6.21%.

1,4-Bis(prop-2-enyl)-2,3-diacetoxynaphthalene (**6**)

1,4-Bis(prop-1-enyl)-2,3-diacetoxynaphthalene (78.0 g, 0.24 mol) was dissolved in a mixture of 100 ml of EtOH and 100 ml of THF. RhCl_3 (0.5 g) was added and the reaction mixture was stirred at 50°C, until conversion was complete by NMR (48 h); the multiplet at δ 5.1–5.0 disappeared and was replaced by a double doublet at δ 1.96. After filtration over silica gel to remove the catalyst, removal of the solvent and subsequent recrystallization from MeOH yielded 42.9 g (55%) of **6**; m.p. 144–145°C. ^1H -NMR: δ 8.07, dd, 2H, J 3.0 Hz, H6 and H7; 7.46, dd, 2H, J 3.0 Hz, H-5,8; 6.56, dd, 2H, J 16 Hz, 1.4 Hz, ArCH_2 ; 6.1–5.9, m, 2H, =CH; 2.75, s, 6H, OAc; 1.96, dd, 6H, J 6.2 Hz, 1.4 Hz, CHCH_3 . ^{13}C -NMR: δ 168.9, C=O; 138.4, Ar; 133.7, ArCH_2 ; 130.7, 128.1, Ar; 126.0, =CH; 125.6, 124.6, Ar; 20.5, $\text{C}(\text{O})\text{CH}_3$; 9.2, CHCH_3 . IR (KBr): 1769, C=O, cm^{-1} . MS (EI): m/z 324.1, M^+ , calcd. 324.0. Anal. calcd. for $\text{C}_{20}\text{H}_{20}\text{O}_4$: C 74.06, H 6.21; found C 74.12, H 6.32%.

2,3-Diacetoxy-1,4-naphthalenedialdehyde (**7**)

Compound **6** (5.0 g, 0.015 mol) was dissolved in 100 ml dry CH_2Cl_2 and cooled to -70°C . Ozone was passed through the solution until the color changed to slightly greenish-blue. The excess of ozone was removed by leading N_2 through the solution. Subsequently 20 ml dimethyl sulfide (DMS) was added and the mixture was allowed to warm and stand for 48 h. DMS and solvent were evaporated and the crude product was triturated with propan-2-ol to give 4.2 g (90%) of **7**; m.p. 194–195°C. ^1H -NMR: δ 10.64, s, 2H, CHO; 8.87, dd, 2H, J 3.3 Hz, H6 and H7; 7.61, dd, 2H, J 3.3 Hz, H5 and H8; 2.42, s, 6H, CH_3 . ^{13}C -NMR ($\text{DMSO}-d_6$): δ 174.0, CH=O; 168.9, C=O; 144.8, ArO; 133.5, Ar; 129.5, 124.8, Ar; 20.3, CH_3 . IR (KBr): 1774, COO; 1698, CH=O, cm^{-1} . MS (EI): m/z 300.063, M^+ , calcd. 300.063. Anal. calcd. for $\text{C}_{16}\text{H}_{12}\text{O}_6 \cdot 0.3\text{H}_2\text{O}$: C 62.87, H 4.15; found: C 62.80, H 4.13%.

2,3-Dihydroxy-1,4-naphthalenedialdehyde (**8**)

To a degassed solution of NaOH (1 g, 25 mmol) in 55 ml of MeOH was added 3.74 g (12.5 mmol) of 2,3-diacetoxy-1,4-naphthalenedialdehyde and 20 ml of THF was added to give a homogeneous dark-green solution. After 30 min, the solvent was evaporated and THF (90 ml) and H_2O (60 ml) were added. 4-Toluenesulfonic acid was added until the pH was 3 and the reaction mixture was stirred overnight. Compound **8** (1.97 g, 73%) precipitated as yellow crystals and was collected by filtration and dried in air; m.p. 158–159°C. ^1H -NMR: δ 13.0, bs, 2H, OH; 10.87, s, 2H, CH=O; 8.38, dd, 2H, J 3.3 Hz, H6 and H7; 7.54, dd, 2H, J 3.3 Hz, H5 and H8. ^{13}C -NMR ($\text{DMSO}-d_6$): δ 194.1, CH=O; 154.2, ArOH; 126.3, 125.4, 122.3, 117.1, Ar. IR (KBr): 1671, CH=O, cm^{-1} . MS (EI): m/z 216.0, M^+ , calcd. 216.0. Anal. calcd. for $\text{C}_{12}\text{H}_8\text{O}_4 \cdot 0.75\text{H}_2\text{O}$: C 62.75, H 3.84; found: C 62.75, H 3.76%.

General procedure for the synthesis of macrocycles **2a–2c**

Dialdehyde **8** (0.20 g, 0.926 mmol) and $\text{Ba}(\text{CF}_3\text{SO}_3)_2$ (0.134 g, 0.309 mmol) were refluxed in a mixture of 50 ml CH_3CN and 50 ml MeOH for 15 min. To the slightly yellow solution, 0.926 mmol of diamine was added in one portion, resulting in an immediate color change from yellow to (orange)red. After 1 h reflux the solution was cooled to room temperature and 50 ml diisopropyl ether were added to induce precipitation of the macrocyclic complex, in typical yields of 70% and pure by ^1H -NMR, FAB-MS and elemental analysis.

Barium(2 +), (5,42 : 14,19 : 28,33-triethenohexabenzo[*b,g,l,q,v,a_1*] [1,4,11,14,21,24]hexaazacyclotriacontine-43,44,45,46,47,48-hexol-*O*, *O'*, *O''*, *O'''*, *O''''*), (OC-6-11)-, salt with trifluoromethanesulfonic acid (1 : 2) (**2a**) Dark red solid; m.p. > 300°C. ^1H -NMR ($\text{DMSO}-d_6$, 60°C): δ 14.4, bs, 6H, OH; 8.75, s, 6H, CH=N; 7.58, bs, 6H, PhenH; 7.40, bs, 6H, PhenH; 7.20, m, 6H, ArH; 7.09, m, 6H, ArH. IR (KBr): 1616, C=N, cm^{-1} . FAB-MS (+VE): m/z 1151.1 ($M-\text{CF}_3\text{SO}_3$)⁺; 1002.1 ($M-2\text{CF}_3\text{SO}_3$)⁺. Anal. calcd. for $\text{C}_{56}\text{H}_{36}\text{BaF}_6\text{N}_6\text{O}_{12}\text{S}_2$: C 51.72, N 6.46, H 2.79; found: C 51.84, N 6.29, H 2.94%.

Barium(2 +), (8,9,20,21,32,33-hexahydro-5,36 : 12,17 : 24,29-triethenotribenzo[*g,q,a_1*] [1,4,11,14,21,24]hexaazacyclotriacontine-

37,38,39,40,41,42-hexol-O,O',O'',O''',O''''-, (OC-6-11)-, salt with trifluoromethanesulfonic acid (1:2) (**2b**) Yellow solid; m.p. > 300°C. ¹H-NMR (DMSO-*d*₆, 75°C): δ 15.00, bs, 6H, OH; 9.36, s, 6H, CH=N; 7.95, s, 6H, ArH; 7.26, s, 6H, ArH; 4.10, s, 12H, NCH₂. ¹³C-NMR (DMSO-*d*₆): δ 174.7, C=N; 162.3, ArOH; 128.0, 125.5, 123.9, 119.3, ArH; 106.9, NCH₂. IR (KBr): 1620, C=N, cm⁻¹. FAB MS (-VE): *m/z* 1155.3 (M⁻); (+VE): 1006.9 (M-CF₃SO₃)⁺; 857.9 (M-2CF₃SO₃)⁺. Anal. calcd. for C₄₄H₃₆BaF₆N₆O₁₂S₂: C 45.71, N 7.27, H 3.14; found: C 46.27, N 7.62, H 3.79%.

Barium(2+), (7a,8,9,10,11,11a,21a,22,23,24,25,25a,35a,36,37,38,39,39a-octadecahydro-5,42:14,19:28,33-triethenohexabenzob[b,g,l,q,v,a₁]/[1,4,11,14,21,24]hexaazacyclotriacontine-43,44,45,46,47,48-hexol-O,O',O'',O''',O''''-, (OC-6-11)-, salt with trifluoromethanesulfonic acid (1:2) (**2c**) Orange solid; m.p. > 300°C. ¹H-NMR (DMSO-*d*₆, 75°C): δ 15.09, bs, 6H, OH; 9.30, s, 6H, CH=N; 8.01, m, 6H, ArH; 7.39, m, 6H, ArH; 4.38, m, 6H, NCH; 1.95, m, 6H, CH₂; 1.61, m, 18H, CH₂. ¹³C-NMR (DMSO-*d*₆): δ 174.1, C=N; 160.8, ArOH; 128.0, 125.4, 124.0, 119.5, ArH; 107.2, NCH; 59.2, 26.9, CH₂. IR (KBr): 1623, C=N, cm⁻¹. FAB-MS (-VE): *m/z* 1317 (M⁻); (+VE): 1169.1 (M-CF₃SO₃)⁺; 1019.2 (M-2CF₃SO₃)⁺. Anal. calcd. for C₅₆H₅₄BaF₆N₆O₁₂S₂: C 51.01, N 6.37, H 4.13; found: C 51.25, N 6.93, H 4.53%.

Barium(2+), (8,9,10,21,22,23,34,35,36,47,48,49-dodecahydro-5,52:13,18:26,31:39,44-tetraethenotetrazabenzob[h,s,d₁,o₁]/[1,5,12,16,23,27,34,38]octaazacyclotetratetracontine-53,54,55,56,57,58,59,60-octol-O,O',O'',O''',O''''-, salt with trifluoromethanesulfonic acid (1:2) (**2d**) The same general procedure was followed as for macrocycles **2a**–**2c**, with the same molar equivalent of Ba(CF₃SO₃)₂, but stirring at room temperature instead of refluxing. Precipitation with diisopropyl ether gave 0.35 g (60%) of a mixture containing [4+4] macrocycle **2d** and a product deriving from [2+2] cyclization. Purification by column chromatography (Sephadex® LH 20–100, eluens CH₃CN) gave the pure orange macrocycle **2d**; m.p. > 300 °C. ¹H-NMR (DMSO-*d*₆): δ 13.8, bs, 8H, OH; 9.90, s, 4H, CH=N; 9.84, s, 4H, CH=N; 7.5, m, 8H, ArH; 6.8, m, 8H, ArH; 4.1, bs, 16H, NCH₂; 2.3, s, 3H CH₃CN; 2.1, bs, 8H, CH₂. ¹³C-NMR (DMSO-*d*₆): δ 180.0, C=N; 165.1, ArOH; 133.2, 130.7, 130.5, 128.0, 123.7, ArH; 111.8, NCH₂; 55.6, CH₂. IR (KBr): 1621, C=N, cm⁻¹. FAB-MS (+VE): *m/z* 1303.6 (M-CF₃SO₃)⁺; 1154.5 (M-2CF₃SO₃)⁺. Anal. calcd. for C₆₂H₅₆BaF₆N₈O₁₄S₂·6H₂O·CH₃CN: C 48.00, N 7.88, H 4.44; found: C 48.05, N 8.10, H 4.47%.

Acknowledgments

We gratefully acknowledge Dr. *W. Niessen*, Gorlaeus Laboratoria, Leiden University and *R. Fokkens*, Institute of Mass Spectrometry, University of Amsterdam, for the Plasma Desorption and Electrospray Ionisation MS experiments, respectively. We also thank the Dutch Foundation for Chemical Research (SON) for financial support.

References and notes

- ^{1a} *S.M. Nelson*, Pure Appl. Chem. **52**, 2461 (1980);
- ^b *P. Guerreiro, P.A. Vigato, D.E. Fenton and P.C. Hellier*, Acta Chem. Scand. **46**, 1025 (1992).
- ^{2a} *S.S. Tandon, L.K. Thompson and J.N. Bridson*, Inorg. Chem. **32**, 32 (1993);
- ^b *S. Brooker and P.D. Croucher*, J. Chem. Soc., Chem. Commun. 1278, (1993);
- ^c *B.F. Hoskins, R. Robson and P. Smith*, J. Chem. Soc., Chem. Commun. 488, (1990);
- ^d *D.E. Fenton, S.J. Kitchen, C.M. Spencer, P.A. Vigato and S. Tamburini*, Inorg. Chim. Acta **139**, 55 (1987).
- ^{3a} *M.J. Grannas, B.F. Hoskins and R. Robson*, Inorg. Chem. **33**, 1071 (1994);
- ^b *S.S. Tandon, L.K. Thompson, J.N. Bridson and M. Bubenik*, Inorg. Chem. **32**, 4621 (1993);
- ^c *D. Coucouvanis, S.G. Jonasdottir, D. Christodoulou, C.G. Kim and J.W. Kampf*, Inorg. Chem. **32**, 2987 (1993);
- ^d *C.S. Velazquez, T.F. Baumann, M.M. Olmstead, H. Hope, A.G.M. Barrett and B.M. Hoffmann*, J. Am. Chem. Soc. **115**, 9997 (1993).
- ⁴ *F.C.J.M. van Veggel, W. Verboom and D.N. Reinhoudt*, Chem. Rev. **94**, 279 (1994).
- ⁵ *B. Srinivas and P.S. Zacharias*, Can. J. Chem. **70**, 2906 (1992).
- ^{6a} *D.G. McCollum, L. Hall, C. White, R. Ostrander, A. Rheingold, J. Whelan and B. Bosnich*, Inorg. Chem. **33**, 924 (1994);
- ^b *D.G. McCollum, C. Fraser, R. Ostrander, A.L. Rheingold and B. Bosnich*, Inorg. Chem. **33**, 2383 (1994).
- ⁷ *F. Arena, C. Floriani, A. Chiesi-Villa and C. Guastini*, Inorg. Chem. **25**, 4589 (1986).
- ^{8a} *F.C.J.M. van Veggel, S. Harkema, M. Bos, W. Verboom, G. Klein Woolhuis and D.N. Reinhoudt*, J. Org. Chem. **54**, 2351 (1989);
- ^b *F.C.J.M. van Veggel, M. Bos, S. Harkema, W. Verboom and D.N. Reinhoudt*, Angew. Chem. Int. Ed. Engl. **28**, 746 (1989).
- ⁹ *D. Dabrowska and W. Rzoska*, Acta Pol. Pharm. **35**, 69 (1978).
- ¹⁰ *L.F. Lindoy*, The Chemistry of Macrocyclic Ligand Complexes, Cambridge University Press, Cambridge, 1989.
- ¹¹ *F.C.J.M. van Veggel, S. Harkema, M. Bos, W. Verboom, C.J. van Staveren, G.J. Gerritsma and D.N. Reinhoudt*, Inorg. Chem. **28**, 1133 (1989).
- ¹² *F.C.J.M. van Veggel, M. Bos, W. Verboom and D.N. Reinhoudt*, Recl. Trav. Chim. Pays-Bas **109**, 515 (1990).
- ¹³ *C.J. van Staveren, J. van Eerden, F.C.J.M. van Veggel, S. Harkema and D.N. Reinhoudt*, J. Am. Chem. Soc. **110**, 4994 (1988).
- ¹⁴ Although the dialdehyde **8** has been reported previously by reacting naphthalene-2,3-diol under Reimer-Tiemann conditions¹⁵, this route failed in our hands.
- ¹⁵ *I.P. Singh, V.K. Shukla, A. K. Dwivedi and N.M. Khanna*, Ind. J. Chem. **28b**, 692 (1989).
- ¹⁶ *E. Gojow, P. Dubourdeaux, N. Gon and J.M. Latour* New. J. Chem. **12**, 931 (1988).
- ¹⁷ Pd/C (10%) in EtOH at reflux temperature gave varying results, because the reaction mostly stopped at less than 100% conversion.
- ¹⁸ To our knowledge, this is only the third example of the formation of a [4+4] Schiff base macrocycle. In all cases the macrocycle contained (CH₂)₃ bridges between the two nitrogens of the diamine.
- ^a *S. Brooker, R.J. Kelly and G.M. Sheldrick*, J. Chem. Soc., Chem. Commun. 487, (1994);
- ^b *V. McKee and W.B. Shepard*, J. Chem. Soc., Chem. Commun. 158, (1985).
- ¹⁹ *A.M.A. Hassaan and M. A. Khalifa*, Monatsh. Chem. **124**, 803 (1993).