# Life review therapy for older adults with moderate depressive symptomatology: a pragmatic randomized controlled trial

# J. Korte<sup>1</sup>\*, E. T. Bohlmeijer<sup>1</sup>, P. Cappeliez<sup>2</sup>, F. Smit<sup>3,4</sup> and G. J. Westerhof<sup>1</sup>

<sup>1</sup> University of Twente, Department of Health, Psychology and Technology, Enschede, The Netherlands

<sup>2</sup> University of Ottawa, School of Psychology, Ottawa, Ontario, Canada

<sup>8</sup> Trimbos Institute (Netherlands Institute of Mental Health and Addiction), Centre of Prevention and Early Intervention, Utrecht, The Netherlands

<sup>4</sup> EMGO + Institute for Health and Health Care Research, Department of Epidemiology and Biostatistics, VU University Medical Centre, Amsterdam, The Netherlands

**Background.** Although there is substantial evidence for the efficacy of life review therapy as an early treatment of depression in later life, its effectiveness in natural settings has not been studied. The present study evaluates an intervention based on life review and narrative therapy in a large multi-site, pragmatic randomized controlled trial (RCT).

**Method.** Life review therapy was compared with care as usual. The primary outcome was depressive symptoms; secondary outcomes were anxiety symptoms, positive mental health, quality of life, and current major depressive episode (MDE). To identify groups for whom the intervention was particularly effective, moderator analyses were carried out (on sociodemographic variables, personality traits, reminiscence functions, clinically relevant depressive and anxiety symptoms, and past MDEs).

**Results.** Compared with care as usual (n=102), life review therapy (n=100) was effective in reducing depressive symptoms, at post-treatment (d=0.60, B=-5.3, p<0.001), at 3-month follow-up (d=0.50, B=-5.0, p<0.001) and for the intervention also at 9-month follow-up (t=5.7, p<0.001). The likelihood of a clinically significant change in depressive symptoms was significantly higher [odds ratio (OR) 3.77, p<0.001 at post-treatment; OR 3.76, p<0.001 at the 3-month follow-up]. Small significant effects were found for symptoms of anxiety and positive mental health. Moderator analyses showed only two significant moderators, the personality trait of extraversion and the reminiscence function of boredom reduction.

**Conclusions**. This study shows the effectiveness of life review therapy as an early intervention for depression in an ecologically valid context, supporting its applicability to a broad target group. The intervention is also effective in reducing anxiety symptoms and strengthening positive mental health.

Received 28 March 2011; Revised 11 July 2011; Accepted 30 August 2011; First published online 14 October 2011

Key words: Depressive symptomatology, life-review therapy, older adults, randomized controlled trial.

## Introduction

In older adults, clinical depression is a significant health problem, carrying a poor prognosis (Cole *et al.* 1999; Beekman *et al.* 2002; Licht-Strunk *et al.* 2007). Regarding the prevalence of depression, rates between 8.8% and 23.6% have been reported (Copeland *et al.* 1999; McDougall *et al.* 2007). The most important risk factor for developing late-life clinical depression is the presence of depressive symptoms not (as yet) meeting

diagnostic criteria (Cuijpers *et al.* 2004; Smit *et al.* 2006). Hence, effective early interventions directed at older adults with mild to moderate depressive symptomatology are of utmost importance. Meta-analytic evidence indicates that early interventions for older adults with depressive symptoms are indeed promising in preventing depressions (Cuijpers *et al.* 2007).

Besides cognitive behavioural therapy (Cuijpers *et al.* 2007), life review has been established as an evidence-based early treatment of depression in later life (Bohlmeijer *et al.* 2003, 2007; Pinquart *et al.* 2007). Life review involves a structured evaluation of one's own life, on the one hand aiming at coping with negative experiences and conflicts and on the other hand at giving a positive meaning to life (Haight, 1992; Wong,

<sup>\*</sup> Address for correspondence: J. Korte, M.Sc., University of Twente, Department of Health, Psychology and Technology, Drienerlolaan 5, PO Box 217, 7500 AE Enschede, The Netherlands.

<sup>(</sup>Email: j.korte@utwente.nl)

1995; Westerhof *et al.* 2010). Life review seems especially suited for older adults who experience loss of meaning in life and who hold a negative view of themselves. Critical life events may increase the use of specific negative functions of reminiscence such as reviving bitter experiences or escaping past problems (Wong, 1995; Cappeliez, 2002; Wink & Schiff, 2002). Associations between these functions and poorer mental health have been found recently (Cully *et al.* 2001; Cappeliez *et al.* 2005; Cappeliez & O'Rourke, 2006; Korte *et al.* 2011; O'Rourke *et al.* 2011).

The above underscores the importance of developing early interventions based on life review. Recently, attempts have been made to combine life review with other interventions aimed at people with clinically relevant mental distress (Watt & Cappeliez, 2000; Cappeliez, 2002; Westerhof *et al.* 2010), for example narrative therapy (Bohlmeijer *et al.* 2008, 2009). In narrative therapy, life stories are considered as a reconstruction of autobiographical memories (Atwood & Ruiz, 1993; Bluck & Levine, 1998). The most important element in life review combined with narrative therapy is to develop alternative, more agentic life stories, in which clients take responsibility for their own preferred ways of living (Bohlmeijer *et al.* 2008, 2009).

To summarize, substantial evidence supports the efficacy of life review as a treatment of depression in later life. However, most trials have not studied the effectiveness of life review in its natural setting (Bohlmeijer et al. 2003, 2007; Pinquart et al. 2007). To fill that gap, we have conducted a large, multi-site, pragmatic randomized controlled trial (RCT) in adults aged  $\geq$  55 years with moderate depressive symptoms, in which we compared the effectiveness of an intervention based on life review and narrative therapy (life review therapy) in its natural setting with care as usual. To study the effects in a natural setting, the Dutch health-care system was mimicked as closely as possible in terms of patient recruitment and offering the intervention. We specifically aimed at young older adults, who are in relatively good health. Life review might help to cope with age-related losses and declines, and to find meaning in this new phase of life (Haight, 1992; Wong, 1995; Westerhof et al. 2010). Most studies on life review interventions have included relatively small sample sizes (Bohlmeijer et al. 2003, 2007; Pinquart et al. 2007). These small sample sizes have precluded addressing important questions about the moderating influence of client characteristics such as sociodemographic attributes, personality traits, patterns of reminiscence and level of depressive symptomatology. Here we sought to identify subgroups for which the intervention might be particularly effective. This should help to determine the suitability of the intervention for different target groups.

# Method

# Design

The design of the current study is described in detail by Korte *et al.* (2009). We carried out a pragmatic, multi-site RCT in which participants were randomly assigned to either life review therapy or care as usual.

## Participants and procedure

Participants were recruited from September 2008 to September 2009, in collaboration with Dutch mental health-care services, through advertisements in newspapers and information booklets, plus a radio interview and commercial. Applicants were referred to a specifically developed website, where they could find more information about the study. Thereafter, applicants were assessed for eligibility and had to sign an informed consent form.

In total, 14 Dutch mental health-care services, in both urban and rural areas, participated in this study. They were responsible for the intake procedure, in which inclusion and exclusion criteria were examined. Therapists at these services, experienced in working with older adults, implemented the intervention. Before being allowed to offer the intervention, they were required to participate in a 2-day training program, supervised by a narrative therapist and a life review therapist. They also participated in two half-day follow-up discussion sessions and a booster training.

Inclusion criteria were age  $\geq$  55 years and the presence of moderate depressive symptoms, operationalized by a score of  $\geq 10$  on the Center for Epidemiologic Studies Depression Scale (CES-D; Radloff, 1977; Beekman et al. 1997). Applicants were excluded if diagnosed with current severe major depressive episode (MDE; eight or nine out of a total of nine symptoms) or with a moderate to high suicide risk according to the Mini International Neuropsychiatric Interview (MINI; Sheehan et al. 1998; Van Vliet & De Beurs, 2007). Furthermore, applicants were excluded if they had a score of  $\leq 9$  on the CES-D (Radloff, 1977), had started taking antidepressant medication or benzodiazepines recently (within the previous 2 months), were currently receiving any psychological treatment, or if the health-care professionals assessed the presence of other serious psychopathology, in which case they were referred for psychological treatment.

Participants were randomly assigned either to life review therapy or to care as usual, by means of a centrally conducted, independent randomization process. The randomization was stratified by gender and presence of current MDE (no current MDE or mild to moderate current MDE), using a computer-generated sequence of numbers.

#### Power analysis

To demonstrate an effect size of at least 0.35 (Cohen's *d*) in a one-tailed test at  $\alpha = 0.05$  and a power of  $(1 - \beta) = 0.80$ , a minimum of 80 participants in each condition was required at follow-up, estimated by power calculation in Stata 7.0 (StataCorp LD, USA). Anticipating a drop-out rate of 20% between baseline and the end of treatment, we needed to include 100 participants for each condition at baseline. This requirement was met, as we included 100 participants in the life review therapy condition and 102 participants in the care as usual condition.

## Measures

All participants completed measures at baseline (t0), post-intervention (t1: 3 months after baseline) and follow-up (t2; 3 months after the end of intervention). Only the intervention condition completed an extended follow-up (t3; 9 months after the intervention) whereas the control condition received the intervention after 6 months.

The primary outcome measure was depressive symptoms (CES-D; Radloff, 1977; Beekman et al. 1997;  $\alpha = 0.83$  in this study). Secondary outcome measures were anxiety symptoms measured on the Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983; Spinhoven et al. 1997;  $\alpha = 0.77$  in this study), positive mental health (Keyes et al. 2008; Lamers *et al.* 2011;  $\alpha = 0.87$  in this study), quality of life assessed by the Euroqol questionnaire (EQ-5D; Euroqol Group, 1990), and current mild to moderate MDE (five to seven out of nine symptoms on the MINI; Sheehan et al. 1998; Van Vliet & De Beurs, 2007). Moderating variables were sociodemographic attributes (age, gender and educational level), personality (neuroticism, extraversion, openness to experience, agreeableness and conscientiousness; NEO-FFI; Costa & McCrae, 1992; Hoekstra et al. 1996;  $\alpha = 0.70$ -0.78,  $\alpha = 0.50$  for openness to experience), reminiscence functions (identity, problem solving, bitterness revival and boredom reduction; Webster, 1997;  $\alpha = 0.79 - 0.85$  in this study), past MDEs (MINI; Sheehan et al. 1998; Van Vliet & De Beurs, 2007), chronic medical conditions (Statistics Netherlands, 1989) and critical life events (Social Readjustment Rating Scale; Holmes & Rahe, 1967). A more detailed description of the measures can be found in Korte et al. (2009).

## Intervention and control condition

The intervention, 'The stories we live by', is conducted in groups of four to six participants and consisted of eight similarly structured sessions of 2 h each. The intervention has three core elements. First, the integration of difficult life events from the past; second, the development of agentic life stories, which helps the participants to cope with present life events and to formulate new goals; third, the retrieval of specific positive memories, which can serve as building blocks of the new life stories.

The first two elements were developed by integrating life review into a narrative therapeutic framework that connects to theories about the role of life stories in the articulation of identity and meaning (Gergen, 1985; Freedman & Combs, 1996). A narrative therapeutic framework enhances the integration of negative events and the restoration of meaning in life in several ways. It stimulates clients to adopt an attitude of curiosity and not-knowing. In this way, ample space is created for clients to explore alternative stories and preferences (White & Epston, 1990). Furthermore, therapists have an arsenal of questions that may be helpful in constructing alternative, more agentic stories about negative life events and life periods (e.g. How were you able to cope with this situation? Were there any pleasant moments in this difficult time? Now, at a much later date, can you say that you have also learned from that period?). In addition, by always focusing on preferences of clients and relating them to other memories, clients are continually invited to express their values and past experiences (White, 2007). Finally, alternative stories are further elaborated by relating them to identity (e.g. What does this say about the person you are?) and to future goals and action (What can you do in the near future to live by this value or meaning?). The third core element was the attention to specific positive memories, special and unique for a certain period in the participants' lives, which are expected to activate nearly forgotten memories, especially in people with depressive symptoms (Serrano et al. 2004; Brewin, 2006; Williams et al. 2007).

The first five intervention sessions were focused on different life themes: origin, youth, work and care, love and conflicts, and loss and difficult times. Before each session, participants had to answer questions about those life themes. For each theme the participants had to describe one difficult life event they were still struggling with. Then they had to answer questions that helped to develop alternative stories that would help to integrate this life event. For each session they had to describe a specific positive memory as well. During the intervention sessions, participants had the opportunity to exchange and discuss the experiences with the other participants. In the last three sessions there was attention to some overarching themes that focused on creating an overview and on the near future. For example, participants had to recall an image that symbolized their lives and explain why (metaphor), they had to make an overview of ups and downs in their lives (course of life), and they had to think about how they wanted to arrange their lives with the knowledge they had gained during the intervention (new beginning).

To check whether the intervention was implemented as intended, one session was audiotaped by all participating mental health-care services. Two researchers independently scored a treatment integrity measure and solved a few disagreements by consensus. The measure addressed five criteria that were crucial in correctly delivering the intervention. First, structure involved the degree to which therapists followed the structure as described in the intervention protocol. Second, life review therapy skills encompassed skills that therapists had in asking the right therapeutic questions. Third, empathy measured whether therapists acknowledged participants' feelings and experiences. Fourth, curiosity assessed whether therapists had an attitude of curiosity and not-knowing. Fifth, group process involved the stimulation of a constructive interaction between participants, to create and elaborate alternative life stories. All criteria were assigned one (unsatisfactory), two (satisfactory) or three points (good), with a total score ranging from 5 to 15 (5-8=unsatisfactory, 9-11 = satisfactory, 12-15 = good). The therapists scored satisfactory (60%: eight out of 14 sessions) or good (40%: six out of 14 sessions).

Participants in the control condition received no intervention. However, they had unrestricted access to care as usual, that is they were not withheld from any treatment (e.g. they may receive psychological treatment). After conclusion of the RCT, the intervention was offered to them.

## Analyses

Analyses were performed according to the intentionto-treat (ITT) approach on missing data, using Missing Value Analysis in PASW 18 (SPSS Inc., USA) to impute all missing data on the continuous measures with expectation–maximization (EM). Missing values were replaced by estimates, which were based on maximum likelihood estimates using observed data in an iterative process (Dempster *et al.* 1977). As a comparison of results based on the imputed ITT sample *versus* the observed data revealed similar outcomes, only results from the ITT analyses are reported. Statistical tests were one-tailed because the hypothesis was unidirectional (assuming superiority of the intervention under the alternative hypothesis) and deemed significant at p < 0.05.

First, independent-samples t tests and  $\chi^2$  tests were conducted to examine differences between the

conditions at baseline. Second, to examine differences between the conditions on outcome measures, regression analysis was used in the case of continuous response variables, and logistic regression was used for dichotomous variables. In both cases, the dependent variables were the post-treatment and follow-up outcome measures with the baseline outcome measures and the treatment dummy (intervention versus control condition) serving as the independent variables. Following Jacobson & Truax (1991), the proportion of participants was determined who showed a clinically significant change on the CES-D from baseline to post-treatment, defined as a reduction on the CES-D of at least 5 points between t1 and t2 while crossing the cut-off value of 16 for clinically relevant depressive symptoms (Beekman et al. 1997; cf. Smit et al. 2006; Steunenberg et al. 2006). Participants manifesting a clinically significant improvement were coded 1 (success) or 0 (failure). This dichotomous outcome was used to calculate the odds ratio (OR) using logistic regression for participants who had a score of  $\geq 16$  at *t*0. The number needed to treat (NNT) was calculated based on the clinically significant change proportions. To investigate whether the effects in the intervention condition were maintained at t2, paired-sample t tests were carried out, comparing the scores on the second follow-up with those at baseline. All analyses took into account the fact that the observations were clustered because participants were 'nested' in each of the 14 mental health services. Therefore, robust standard errors were computed using the first-order Taylor-series linearization method as implemented in Stata version 8.2. Standardized mean differences (Cohen's d) at postintervention and follow-up were calculated as the difference between the means of the treatment and control condition divided by the standard error of the control condition.

Finally, to identify whether sociodemographic variables (age, gender, educational level), personality traits, reminiscence functions, clinically relevant depressive and anxiety symptoms, past MDEs, chronic medical conditions and critical life events moderated the effect of the intervention on the change of depressive symptoms at post-intervention, linear regression analyses were used. The cut-off scores for clinically relevant symptoms of depression ( $\geq 16$ ) and anxiety  $(\geq 11)$  and the medians on the scores on personality traits, past MDEs, chronic medical conditions and critical life events at baseline were used as cut-off scores for recoding them as dichotomous variables. In linear regression models, we entered the change score of t1-t0 of depressive symptoms as a dependent variable, and the intervention dummy (intervention condition = 1 versus control condition = 0), the potential



Fig. 1. Flow of participants. Analyses were performed according to the intention-to-treat (ITT) approach.

moderator and their interaction as independent variables.

#### Results

## Enrolment, treatment adherence and drop-out

Figure 1 shows the flow of participants. In total, 274 older adults expressed an interest to participate in the intervention. Based on the exclusion criteria, 49 applicants were excluded before the randomization (no depressive symptoms: n=23, current severe MDE: n=7, moderate to high suicide risk: n=3, recent antidepressant medication: n=5, currently receiving psychological treatment: n=3, and other serious psychology: n=8). Eighteen additional participants decided not to participate because they lacked motivation, and five applicants were excluded for other reasons. The remaining 202 participants were randomly assigned to the intervention (n=100) or the

control condition (n = 102). After randomization, one participant decided not to start with the intervention, and seven others discontinued the intervention due to lack of motivation (n = 5) or health problems (n = 2). Of the participants, 85.1% filled out the questionnaires at all measurement times (80.4% in the intervention group). The total percentage of missing data was 15.1%, due to unanswered items (2.3%) or incomplete assessments (12.8%). These missing values were imputed at baseline, post-intervention and follow-up. Participants who completed the assessments did not differ significantly at baseline from those who did not complete all measures.

## **Baseline characteristics**

Participants had a mean age of 63 years (s.D. =6.5, range =55-83) and were predominantly female (Table 1). Their education was distributed evenly

### 1168 J. Korte et al.

Table 1.	Baseline	characteristics	of partic	ipants for	total	sample,	life revi	iew th	erapy	and
care as us	sual									

	Total sample $(n=202)$	Life review therapy ( <i>n</i> =100)	Care as usual (n=102)	
Age <sup>a</sup>				
Mean (s.D.)	63.3 (6.5)	63.3 (6.2)	63.3 (6.8)	
Range	55-83	55-83	55-83	
Sex <sup>b</sup>				
Female	76.7 80.0		73.5	
Male	23.3	20.0	26.5	
Education <sup>b</sup>				
<11 years	36.6	37.0	36.3	
11–14 years	33.7	35.0	32.4	
>14 years	29.7	28.0	31.4	
Chronic medical conditions <sup>a</sup>				
Mean (s.D.)	1.5 (1.4)	1.4 (1.3)	1.5 (1.4)	
Range	0–6	0–6	0–6	
Critical life events <sup>a</sup>				
Mean (s.D.)	2.3 (1.5)	2.4 (1.5)	2.3 (1.5)	
Range	0–7	0–7	0–7	
Employment <sup>b</sup>				
Paid work	15.8	19.0	12.7	
Retired	30.7	33.0	28.4	
Housekeeping	15.8	16.0	15.7	
Volunteer work	18.3	19.0	17.6	
Disability pension	15.8	10.0	21.6	
Unemployed	3.5	3.0	3.9	

s.D., Standard deviation.

<sup>a</sup> No significant differences between intervention and control condition (*t* test

with p > 0.05).

<sup>b</sup> No significant differences between intervention and control condition ( $\chi^2$  test with p > 0.05).

across three categories (<11 years, 11–14 years, and >14 years). About one-third of the participants were retired. On average, participants had one to two chronic medical conditions (mean =1.5, s.D. =1.4) and they had experienced two to three critical life events in the past 3 years (mean =2.3, s.D. =1.5). Table 2 shows that participants at baseline scored an average of 20.5 (s.D. =8.6) on the CES-D, reflecting clinically relevant depressive symptoms.

#### Treatment effects

Table 2 presents the means and standard errors for the outcome measures at post-treatment, first and second follow-ups, the results of the regression analysis, and the effect sizes. Compared to the control condition, participants in the intervention condition reported significantly decreased depressive symptoms at post-treatment (B = -5.3, p < 0.001) and first followup (B = -5.0, p < 0.001). Effect sizes at post-treatment (d = 0.60) and first follow-up (d = 0.50) are medium (Cohen, 1992). The effects of the intervention condition were maintained at the second follow-up [t(99) = 5.7, p < 0.001]. The proportion of participants who reached a clinically significant change at post-treatment on the CES-D in the life review therapy condition was 31/68 (45.6%) *versus* 12/71 (16.9%) in the care-as-usual condition [OR 3.77, 95% confidence interval (CI) 1.38– 10.30, p < 0.001, NNT = 3.5]. At follow-up, in the intervention condition, 35/68 (51.5%) participants *versus* 15/71 (21.1%) participants in the control condition showed a successful outcome (OR 3.76, 95% CI 1.45– 9.72, p < 0.001, NNT = 3.3).

Compared to the control condition, participants in the intervention condition reported a significant decrease in anxiety symptoms (t1: B = -1.1, p < 0.01,

		Life review therapy $(n=100)$		Care as usual $(n = 102)$					
Measures		Mean	S.E.	Mean	S.E.	В	S.E.	d	
Primary outcome									
CES-D (0-60)	Baseline	20.5	1.1	20.6	0.74				
	Post-treatment	15.8	1.2	21.2	0.90	-5.3***	0.99	0.60	
	Follow-up	15.3	1.1	20.4	1.0	$-5.0^{***}$	1.3	0.50	
	Second follow-up	15.4	1.0			t=5.7***			
Secondary outcomes									
HADS-A (0-21)	Baseline	8.3	0.39	8.4	0.39				
	Post-treatment	6.8	0.34	8.0	0.42	$-1.1^{**}$	0.49	0.28	
	Follow-up	6.8	0.34	7.7	0.38	$-0.89^{*}$	0.36	0.25	
	Second follow-up	6.7	0.34			$t = 5.0^{***}$			
MHC-SF (0-70)	Baseline	34.2	0.97	34.0	1.2				
	Post-treatment	38.0	1.5	33.8	1.5	4.0***	1.3	0.29	
	Follow-up	36.9	1.1	33.7	1.2	2.9***	0.80	0.26	
	Second follow-up	39.0	1.1		$t = -4.2^{*}$				
EQ-5D (0-1)	Baseline	0.83	0.02	0.81	0.01				
	Post-treatment	0.81	0.02	0.78	0.02	0.02	0.02	0.17	
	Follow-up	0.81	0.02	0.80	0.01	-0.00	0.02	0.07	
	Second follow-up	0.83	0.02			t = 0.36			
MINI-MDE (%)	Baseline	19.0		17.6					
	Post-treatment	10.0		15.7		OR 0.57	0.22		

**Table 2.** Means and standard deviations for all outcome measures (at baseline, post-treatment, follow-up and second follow-up) and results of the GLM repeated measures and Cohen's d for intervention effects (taking clustering into account)

GLM, General Linear Model; CES-D, Center for Epidemiologic Studies Depression Scale; HADS-A, Hospital Anxiety and Depression Scale – Anxiety Score; MHC-SF, Mental Health Continuum – Short Form; EQ-5D, EuroQol Questionnaire; MINI-MDE, Mini International Neuropsychiatric Interview – mild to moderate major depressive episode; S.E., standard error; OR, odds ratio.

\*\*\* p < 0.001, \*\* p < 0.01, \* p < 0.05.

*t*2: *B* = -0.89, *p* < 0.05) and a significant improvement in positive mental health (*t*1: *B* = 4.0, *p* < 0.001, *t*2: *B* = 2.9, *p* < 0.001). Effect sizes for anxiety symptoms (*t*1: *d* = 0.28, *t*2: *d* = 0.25) and positive mental health (*t*1: *d* = 0.29, *t*2: *d* = 0.26) were small (Cohen, 1992). The effects were maintained at the second follow-up in the intervention condition [anxiety symptoms: *t*(99) = 5.0, *p* < 0.001, positive mental health: *t*(99) = -4.2, *p* < 0.05]. No significant effects were found for quality of life (*t*1: *B* = 0.02, *p* = 0.20, *t*2: *B* = -0.00, *p* = 0.41). Improvement on current mild to moderate MDE was not statistically significant (*p* = 0.08, OR 0.57).

## Moderator analyses

We investigated whether certain groups of participants derived more benefit from the intervention with respect to depressive symptoms. Analyses showed only two significant moderators, the personality trait of extraversion (B = -4.1, p = 0.03) and the use of reminiscence to reduce boredom in the present (B = 4.6, p = 0.03), in the sense that people with higher

levels of extraversion and lower levels of boredom reduction obtained higher benefits from the intervention. No significant interactions were found for the other putative moderators, that is age, gender, educational level, traits of neuroticism, openness to experiences, agreeableness, conscientiousness, reminiscence functions of identity, problem solving, bitterness revival, clinically relevant depressive and anxiety symptoms at baseline, past MDEs, chronic medical conditions and critical life events at baseline.

#### Discussion

#### Main findings

The present study evaluated an early intervention that integrates life review and narrative therapy under real-life conditions, using a large, pragmatic, multi-site RCT in older adults with moderate depressive symptoms. The results indicate that such a life review therapy is effective in reducing depressive symptoms. These effects were maintained at 3- and 9-month

follow-ups. We found medium effect sizes for depressive symptoms, both post-treatment (d = 0.60) and at follow-up (d = 0.50). These effect sizes are comparable to the effect size of 0.42 that was found for early interventions on psychological distress by Cuijpers et al. (2007). Moreover, the likelihood of a clinically significant change in depressive symptoms was significantly and substantially higher in the intervention condition than in the care as usual condition. These results support life review therapy as an effective early intervention for older adults with depressive symptomatology. The effects on major depression at the 3-month follow-up were not statistically significant at p = 0.08, but the effects on clinical cases of preventive and early interventions are most prominently found after longer periods of time (Cuijpers et al. 2008). Our finding that the average level of depressive symptoms remained below the cut-off score for possible caseness of depression at the 9-month follow-up is promising, but it needs to be substantiated by longer follow-up measurements under controlled conditions.

In addition, significant, albeit small, effects were observed for symptoms of anxiety and positive mental health. It thus seems that life review therapy has the potential not only to reduce symptoms of psychopathology but also to improve positive mental health, that is emotional, psychological and social well-being. There is growing recognition that enhancing positive mental health is an important objective in public mental health (Jané-Llopis & Barry, 2005; Keyes, 2005; Keyes et al. 2010; Fledderus et al. 2011). No significant differences were found for quality of life, however. This might be explained by our method of measuring quality of life. The EQ-5D (Brooks, 1996) focuses on health-related quality of life across various health dimensions, which makes it unlikely that effects in a single domain are sufficient to cause a change on the EQ-5D. After all, factors such as mobility and self-care are probably difficult to influence with a psychological intervention.

Overall, the results showed that life review therapy is applicable to a broad target group of adults of 55 years and over experiencing moderate depressive symptomatology. Moderator analyses on depressive symptoms identified only two significant moderators, the personality trait of extraversion and the reminiscence function of boredom reduction. Our results demonstrate that the intervention is more effective for people having higher levels of extraversion. In addition to being more energetic, more extraverted individuals may be more able and inclined to share their feelings, emotions and thoughts with others and thus may benefit more from an intervention conducted in groups. Indeed, it has been reported that extraversion predicts a higher frequency of reminiscence in general, and in particular for conversational and relational purposes (Cappeliez & O'Rourke, 2002). More extraverted individuals may also be more inclined to look at the 'silver lining' and thus benefit more from the components of the intervention directed at positive memories. Furthermore, older adults demonstrating higher levels of boredom reduction profit less from the intervention. High levels of boredom reduction reminiscence reflect taking solace in the past in front of an unstimulating environment or an unrewarding future (Webster, 1997). Indeed, this function of reminiscence has been linked with the difficulty in finding meaning in the present and in seeking new goals in the future (Cappeliez & O'Rourke, 2002). Importantly, the reminiscence function of boredom reduction has been strongly linked with negative physical and psychological health outcomes, including depression and anxiety (Cappeliez & O'Rourke, 2006; O'Rourke et al. 2011; Korte et al. 2011). It is possible that older adults who have difficulty finding meaning in the present may profit more from interventions that focus on the here and now.

There has been some criticism that participating in a life review intervention would stimulate bitterness and rumination in people with depression (Cully et al. 2001; Coleman, 2005). This study shows that this is not the case. Effects of the intervention were independent of level of depressive symptoms and prior MDEs. It is especially noteworthy that the personality trait of neuroticism and the use of reminiscence for bitterness revival were not found to moderate the effects of life review therapy. Bitterness revival is the repeated recall of memories about personal crisis and conflicts (Webster, 1997). It is related to neuroticism and depression (Webster, 1994; Cully et al. 2001; Molinari et al. 2001; Cappeliez & O'Rourke, 2002, 2006; Cappeliez et al. 2005; Korte et al. 2011). It seems that participants with higher levels of depressive symptoms, neuroticism and bitterness revival are also capable of developing alternative stories on their lives that are more positive, meaningful and empowering. The general lack of moderating effects suggests that the intervention is broadly applicable to a variety of participants.

This is the first study to investigate the beneficial effects of combining life review with narrative therapy. Although the usefulness of narrative therapy with older adults has been advanced (Kroph & Tandy, 1998), its effects have not been examined experimentally. A central focus in narrative therapy is to develop alternative stories about one's life that are more agentic (Polkinghorne, 1996), richer and more inclusive of experience (White & Epston, 1990; Payne, 2000). A recent qualitative study on the same intervention has identified these transformations in the

self-narrative (Bohlmeijer *et al.*, unpublished observations). One interesting line of investigation would be to collect and analyze stories about therapy to ascertain whether the increase of agency in life-stories is indeed a working mechanism (Adler *et al.* 2008).

## Limitations

Some limitations of this study must be acknowledged. First, the control condition being a care-as-usual waiting list condition, there was no control for the possible influence of non-specific factors, such as attention and social interaction. It would have been optimal to compare the targeted intervention with an active group intervention, or at least with an attention placebo comparison group. Second, given the relatively long follow-up period, it was not possible for ethical reasons to compare the intervention with care as usual during the second follow-up. Although the effects in the intervention group were maintained at the 9-month follow-up, we cannot claim that these effects were uniquely caused by the intervention.

Third, this study used self-report outcome measures, which might lead to biased findings. Future research could use more physiological outcome measures. A recent study shows, for example, that life review is also effective in reducing high blood pressure (Houston *et al.* 2011). Fourth, as the participants in our study were young older adults, generalizations regarding age should be made with some caution. However, in line with meta-analytic evidence (Bohlmeijer *et al.* 2007), we found that, with respect to depressive symptoms, young older adults.

#### Conclusions

This study shows that life review therapy is effective as an early intervention for older adults with depressive symptomatology, that life review therapy is effective under ecologically valid conditions, and that life review therapy is effective for a large variety of older adults (with the possible exception of introverted people and those who make considerable use of reminiscence to escape the present). Although the effects of life review therapy were maintained at the 9-month follow-up, it is recommended that the longterm effects be studied in randomized controlled conditions and that active control conditions be used to control for non-specific factors.

# Acknowledgments

This study was approved by the METiGG, a medical ethics committee for research in mental health-care

settings in The Netherlands. In addition, this study has been registered in the Netherlands Trial Register, the primary Dutch register for clinical trials (TC = 1860). The study was funded by ZonMw, the Netherlands organization for health research and development (grant no. 120610003). We acknowledge with many thanks therapists and participants for taking part in this study.

## **Declaration of Interest**

None.

# References

- Adler JM, Skalina LM, McAdams DP (2008). The narrative reconstruction of psychotherapy and psychological health. *Psychotherapy Research* **18**, 719–734.
- Atwood JD, Ruiz J (1993). Social constructionist therapy with the elderly. *Journal of Family Psychotherapy* **4**, 1–32.
- Beekman ATF, Deeg DJ, Van Limbeek J, Braam AW, De Vries MZ, Van Tilburg W (1997). Criterion validity of the Center for Epidemiologic Studies Depression Scale (CES-D): results from a community-based sample of older subjects in The Netherlands. *Psychological Medicine* 27, 231–235.
- Beekman ATF, Penninx BWJH, Deeg DJH, De Beurs E, Geerings SW, Van Tilburg W (2002). The impact of depression on the well-being, disability and use of services in older adults: a longitudinal perspective. *Acta Psychiatrica Scandinavica* **105**, 20–27.
- Bluck S, Levine LJ (1998). Reminiscence as autobiographical memory: a catalyst for reminiscence theory development. *Ageing & Society* 18, 185–208.
- Bohlmeijer ET, Kramer J, Smit F, Onrust S, van Marwijk H (2009). The effects of integrative reminiscence on depressive symptomatology and mastery of older adults. *Community Mental Health Journal* **45**, 476–484.
- Bohlmeijer ET, Roemer M, Cuijpers P, Smit F (2007). The effects of reminiscence on psychological well-being in older adults: a meta-analysis. *Aging & Mental Health* 11, 291–300.
- Bohlmeijer ET, Smit F, Cuijpers P (2003). Effects of reminiscence and life review on late-life depression: a meta-analysis. *International Journal of Geriatric Psychiatry* 18, 1088–1094.
- Bohlmeijer ET, Westerhof GJ, Emmerik-De Jong M (2008). The effects of integrative reminiscence on meaning in life: results of a quasi-experimental study. *Aging & Mental Health* **12**, 639–646.
- **Brewin CR** (2006). Understanding cognitive behaviour therapy: a retrieval competition account. *Behaviour Research and Therapy* **44**, 765–784.
- Brooks R (1996). EuroQol: the current state of play. *Health Policy* **37**, 53–72.
- **Cappeliez P** (2002). Cognitive-reminiscence therapy for depressed older adults in day hospital and long-term care. In *Critical Advances in Reminiscence Work: From Theory*

to Application (ed. J. D. Webster and B. K. Haight), pp. 300–313. Springer: New York.

Cappeliez P, O'Rourke N (2002). Personality traits and existential concerns as predictors of the functions of reminiscence in older adults. *Journals of Gerontology: Psychological Sciences* 57, 116–123.

Cappeliez P, O'Rourke N (2006). Empirical validation of a model of reminiscence and health in later life. *Journals of Gerontology. Series B, Psychological Sciences and Social Sciences* **61**, 237–244.

Cappeliez P, O'Rourke N, Chaudhury H (2005). Functions of reminiscence and mental health in later life. *Aging and Mental Health* 9, 295–301.

Cohen J (1992). A power primer. *Psychological Bulletin* **112**, 155–159.

**Cole MG, Bellavance F, Masour A** (1999). Prognosis of depression in elderly community and primary care populations: a systematic review and meta-analysis. *American Journal of Psychiatry* **156**, 1182–1189.

Coleman PG (2005). Uses of reminiscence: functions and benefits. *Aging & Mental Health* 9, 291–294.

Copeland JR, Beekman AT, Dewey ME, Hooijer C, Jordan A, Lawlor BA, Lobo A, Magnusson H, Mann AH, Meller I, Prince MJ, Reischies F, Turrina C, De Vries MW, Wilson KC (1999). Depression in Europe. Geographical distribution among older people. *British Journal of Psychiatry* 174, 312–321.

Costa Jr. PT, McCrae RR (1992). Revised NEO Personality Inventory (NEO PI-R) and NEO Five-Factor Inventory (NEO-FFI). Psychological Assessment Resources: Odessa, FL.

Cuijpers P, De Graaf R, Van Dorsselaer S (2004). Minor depression: risk profiles, functional disability, health care use and risk of developing major depression. *Journal of Affective Disorders* **79**, 71–79.

Cuijpers P, Smit F, Van Straten A (2007). Psychological treatments of subthreshold depression: a meta-analytic review. *Acta Psychiatrica Scandinavica* **115**, 434–441.

Cuijpers P, Van Straten A, Smit F, Mihalopoulos C, Beekman A (2008). Preventing the onset of depressive disorders: a meta analytic review of psychological interventions. *American Journal of Psychiatry* 165, 1272–1280.

Cully JA, LaVoie D, Gfeller JD (2001). Reminiscence, personality, and psychological functioning in older adults. *Gerontologist* **41**, 89–95.

**Dempster AP, Laird NM, Rubin DB** (1977). Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society* **39**, 1–38.

Euroqol Group (1990). EuroQol – a new facility for the measurement of health-related quality of life. *Health Policy* 16, 199–208.

Fledderus M, Bohlmeijer ET, Smit F, Westerhof GJ (2011). Mental health promotion as a new goal in public mental health care: a randomized controlled trial of an intervention enhancing psychological flexibility. *American Journal of Public Health* **100**, 2372–2378.

**Freedman J, Combs G** (1996). *Narrative Therapy: The Social Construction of Preferred Realities*. W. W. Norton: New York.

Gergen KJ (1985). The social constructionist movement in modern psychology. *American Psychologist* **40**, 266–275. Haight BK (1992). Long-term effects of a structured life review process. *Journal of Gerontology* **47**, 312–315.

Hoekstra HA, Ormel H, de Fruyt F (1996). NEO Personality Questionnaires: NEO-PI-R and NEO-FFI [in Dutch]. Swets & Zeitlinger: Lisse, The Netherlands.

Holmes TH, Rahe RH (1967). The Social Readjustment Rating Scale. *Journal of Psychosomatic Research* **11**, 213–218.

Houston TK, Allison JJ, Sussman M, Horn W, Holt CL, Trobaugh J, Salas M, Pisu M, Cuffee YL, Larkin D, Person SD, Barton B, Kiefe CI, Hullet S (2011). Culturally appropriate storytelling to improve blood pressure: a randomized trial. *Annals of Internal Medicine* **154**, 77–84.

Jacobson NS, Truax P (1991). Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology* **59**, 12–19.

Jané-Llopis E, Barry MM (2005). What makes mental health promotion effective? *Promotion and Education* **12**, 47–54.

Keyes CLM (2005). Mental illness and/or mental health? Investigating axioms of the complete state model of health. *Journal of Consulting and Clinical Psychology* 73, 539–548.

Keyes CLM, Dhingra SS, Simoes EJ (2010). Change in level of positive mental health as a predictor of future risk of mental illness. *American Journal of Public Health* 100, 2366–2371.

Keyes CLM, Wissing M, Potgieter JP, Temane M, Kruger A, Van Rooy S (2008). Evaluation of the mental health continuum-short form (MHC-SF) in Setswana-speaking South Africans. *Clinical Psychology and Psychotherapy* **15**, 181–192.

Korte J, Bohlmeijer ET, Smit F (2009). Prevention of depression and anxiety in later life: design of a randomized controlled trial for the clinical and economic evaluation of a life-review intervention. *BMC Public Health* 9, 250.

Korte J, Bohlmeijer ET, Westerhof GJ, Pot AM (2011). Reminiscence and adaptation to critical life events in older adults with mild to moderate depressive symptoms. *Aging & Mental Health* **15**, 638–646.

Kroph NP, Tandy C (1998). Narrative therapy with older clients: the use of a 'meaning-making' approach. *Clinical Gerontologist* 18, 3–16.

Lamers SMA, Westerhof GJ, Bohlmeijer ET, ten Klooster PM, Keyes CLM (2011). Evaluating the psychometric properties of the Mental Health Continuum-Short Form (MHC-SF). *Journal of Clinical Psychology* **67**, 99–110.

Licht-Strunk E, van der Windt DA, van Marwijk HW, de Haan M, Beekman AT (2007). The prognosis of depression in older patients in general practice and the community. A systematic review. *Family Practice* 24, 168–180.

McDougall FA, Kvaal K, Matthews FE, Paykel E, Jones PB, Dewey ME, Brayne C (2007). Prevalence of depression in older people in England and Wales: the MRC CFA Study. *Psychological Medicine* **37**, 1787–1795.

Molinari V, Cully JA, Kendjelic EM, Kunik ME (2001). Reminiscence and its relationship to attachment and personality in geropsychiatric patients. *International Journal of Aging and Human Development* **52**, 173–184.

O'Rourke N, Cappeliez P, Claxton A (2011). Functions of reminiscence and the psychological well-being of young-old and older adults over time. *Aging & Mental Health* **15**, 272–281.

Payne M (2000). Narrative Therapy. An Introduction for Counsellors. Sage: London.

Pinquart M, Duberstein PR, Lyness JM (2007). Effects of psychotherapy and other behavioral interventions on clinically depressed older adults: a meta-analysis. *Aging & Mental Health* **11**, 645–657.

Polkinghorne DE (1996). Transformative narratives: from victimic to agentic life plots. *American Journal of Occupational Therapy* **50**, 299–305.

Radloff LS (1977). The CES-D scale: a self-report depression scale for research in the general population. *Journal of Applied Psychological Measurement* 1, 385–401.

Serrano JP, Latorre JM, Gatz M, Montanes J (2004). Life review therapy using autobiographical retrieval practice for older adults with depressive symptomatology. *Psychology and Aging* **19**, 272–277.

Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R, Dunbar GC (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* **59**, 22–33.

Smit F, Ederveen A, Cuijpers P, Deeg D, Beekman A (2006). Opportunities for cost-effective prevention of late-life depression: an epidemiological approach. *Archives of General Psychiatry* 63, 290–296.

Spinhoven PH, Ormel J, Sloekers PPA, Kempen GIJM, Speckens AEM, Van Hemert AM (1997). A validation study of the Hospital Anxiety and Depression Scale (HADS) in different groups of Dutch subjects. *Psychological Medicine* 27, 363–370.

**Statistics Netherlands** (1989). *Health Interview Questionnaire*. Statistics Netherlands: Heerlen, The Netherlands. Steunenberg B, Beekman ATF, Deeg DJH, Kerkhof AJFM (2006). Personality and the onset of depression in late life. *Journal of Affective Disorders* **92**, 243–251.

Van Vliet IM, De Beurs E (2007). The MINI-International Neuropsychiatric Interview (M.I.N.I.). A brief structured diagnostic psychiatric interview for DSM-IV and ICD-10 psychiatric disorders [in Dutch]. *Tijdschrift voor Psychiatrie* 49, 393–397.

Watt LM, Cappeliez P (2000). Integrative and instrumental reminiscence therapies for depression in older adults: intervention strategies and treatment effectiveness. *Aging & Mental Health* **4**, 166–177.

Webster JD (1994). Predictors of reminiscence: a lifespan perspective. *Canadian Journal on Aging* **13**, 66–78.

Webster JD (1997). The reminiscence function scale: a replication. *International Journal of Aging and Human Development* 44, 137–148.

Westerhof GJ, Bohlmeijer ET, Webster JD (2010). Reminiscence and mental health: a review of recent progress in theory, research, and intervention. *Aging & Society* **30**, 697–721.

White M (2007). *Maps of Narrative Therapy*. W. W. Norton: New York.

White M, Epston D (1990). Narrative Means to Therapeutic *Ends*. W. W. Norton : New York.

Williams JM, Barnhofer T, Crane C, Herman D, Raes F, Watkins E, Dalgleish T (2007). Autobiographical memory specificity and emotional disorder. *Psychological Bulletin* 133, 122–148.

Wink P, Schiff B (2002). To review or not to review? The role of personality and life events in life review and adaptation to older age. In *Critical Advances in Reminiscence Work: From Theory to Application* (ed. J. D. Webster and B. K. Haight), pp. 44–60. Springer: New York.

Wong PTP (1995). The processes of adaptive reminiscence. In *The Art and Science of Reminiscing : Theory, Research, Methods and Applications* (ed. B. K. Haight and J. D. Webster), pp. 22–35. Taylor & Francis : Washington, DC.

Zigmond AS, Snaith RP (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica* **67**, 361–370.