

## **Predicting presence of macrovascular causes in non-traumatic intracerebral hemorrhage; the DIAGRAM prediction score**

Nina A Hilkens, MD\*<sup>1</sup>, Charlotte JJ van Asch, MD, PhD \*<sup>2,3</sup>, David J Werring, MD, PhD\*<sup>4</sup>, Duncan Wilson, MD<sup>4</sup>, Gabriël JE Rinkel, MD, FRCPE<sup>2</sup>, Ale Algra, MD, PhD<sup>1,2</sup>, Birgitta K Velthuis, MD, PhD<sup>5</sup>, Gérard AP de Kort, MD<sup>5</sup>, Theo D Witkamp, MD, PhD<sup>5</sup>, Koen M van Nieuwenhuizen, MD<sup>2</sup>, Frank-Erik de Leeuw, MD, PhD<sup>6</sup>, Wouter J Schonewille, MD, PhD<sup>7</sup> Paul LM de Kort, MD, PhD<sup>8</sup>, Diederik WJ Dippel, MD, PhD<sup>9</sup>, Theodora WM Raaymakers, MD<sup>10</sup>, Jeannette Hofmeijer, MD, PhD<sup>11</sup>, Marieke JH Wermer, MD, PhD<sup>12</sup>, Henk Kerckhoff, MD, PhD<sup>13</sup>, Korné Jellema, MD, PhD<sup>14</sup>, Irene M Bronner, MD, PhD<sup>15</sup>, Michel JM Remmers, MD<sup>16</sup>, Henri Paul Bienfait, MD<sup>17</sup>, Ron JGM Witjes, MD<sup>18</sup>, H Rolf Jäger, MD, FRCR<sup>19</sup>, Jacoba P Greving, PhD<sup>1</sup>, Catharina JM Klijn, MD, PhD<sup>2,6</sup>; the DIAGRAM study group

\* These authors contributed equally to the manuscript

1. Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, the Netherlands
2. Department of Neurology and Neurosurgery, Brain Center Rudolf Magnus, University Medical Center, Utrecht, the Netherlands
3. Kempenhaeghe, Academic Centre for Epileptology, Heeze, the Netherlands
4. Stroke Research Centre, Department of Brain Repair and Rehabilitation, UCL Institute of Neurology and The National Hospital for Neurology and Neurosurgery, Queen Square, London, UK
5. Department of Radiology, University Medical Center Utrecht, Utrecht, the Netherlands
6. Department of Neurology, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, Nijmegen, the Netherlands
7. Department of Neurology, St. Antonius Hospital, Nieuwegein, the Netherlands
8. Department of Neurology, St. Elisabeth Hospital, Tilburg, the Netherlands
9. Department of Neurology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands
10. Department of Neurology, Meander Medical Center, Amersfoort, the Netherlands
11. Department of Neurology, Rijnstate Hospital, Arnhem, the Netherlands

12. Department of Neurology, Leiden University Medical Center, Leiden, the Netherlands
13. Department of Neurology, Albert Schweitzer Hospital, Dordrecht, the Netherlands
14. Department of Neurology, MCH Westeinde, The Hague, the Netherlands
15. Department of Neurology, Flevo Hospital, Almere, the Netherlands
16. Department of Neurology, Amphia Hospital, Breda, the Netherlands
17. Department of Neurology, Gelre Hospital, Apeldoorn, the Netherlands
18. Department of Neurology, Tergooi Hospitals, Blaricum, the Netherlands
19. Neuroradiological Academic Unit, Department of Brain Repair and Rehabilitation, UCL Institute of Neurology and The National Hospital for Neurology and Neurosurgery, Queen Square, London, UK

**Corresponding author**

Prof. Catharina JM Klijn, MD

Department of Neurology, Donders Institute for Brain, Cognition and Behaviour,

Center for Neuroscience

Radboud University Medical Center Nijmegen

PO box 9101, 6500 HB Nijmegen

The Netherlands

Tel.: +31(0)243613394

Fax: +31(0)243655135

Email: [Karin.Klijn@radboudumc.nl](mailto:Karin.Klijn@radboudumc.nl)

**Word count:** 3022

**Number of references:** 20

**Key words:** intracerebral hemorrhage, CT angiography, digital subtraction angiography, arteriovenous malformation

## **ABSTRACT**

**Objective** A substantial part of non-traumatic intracerebral hemorrhages (ICH) arises from a macrovascular cause, but there is little guidance on selection of patients for additional diagnostic work-up. We aimed to develop and externally validate a model for predicting the probability of a macrovascular cause in patients with non-traumatic ICH.

**Methods** The DIAGRAM study (N=298; 69 macrovascular cause; 23%) is a prospective, multicenter study, assessing yield and accuracy of CTA, MRI/MRA and intra-arterial catheter angiography in diagnosing macrovascular causes in patients with non-traumatic ICH. We considered pre-specified patient and ICH characteristics in multivariable logistic regression analyses as predictors for a macrovascular cause. We combined independent predictors in a model, which we validated in an external cohort of 173 ICH patients (78 macrovascular cause, 45%).

**Results** Independent predictors were younger age, lobar or posterior fossa (versus deep) location of ICH and absence of small vessel disease (SVD). A model that combined these predictors showed good performance in the development data (c-statistic 0.83; 95% CI 0.78-0.88) and moderate performance in external validation (c-statistic 0.66; 0.58-0.74). When CTA results were added, the c-statistic was excellent (0.91; 0.88-0.94), and good after external validation (0.88; 0.83-0.94). Predicted probabilities varied from 1% in patients aged 51-70 years with deep ICH and SVD, to more than 50% in patients aged 18-50 years with lobar or posterior fossa ICH without SVD.

**Conclusion** The DIAGRAM scores help to predict the probability of a macrovascular cause in patients with non-traumatic ICH based on age, ICH location, SVD and CTA.

## INTRODUCTION

Intracerebral hemorrhage (ICH) accounts for 15-20% of all strokes and is the most devastating stroke subtype.[1,2] Around 15-25% of ICHs are caused by an underlying macrovascular cause, such as an arteriovenous malformation (AVM), aneurysm, dural arteriovenous fistula (dAVF), cavernoma, and cerebral venous sinus thrombosis.[3-5] Among young adults, macrovascular causes are the leading cause of ICH.[6]

Early diagnosis of underlying macrovascular lesions can influence clinical management and prognosis, as timely intervention might prevent recurrent haemorrhage.[7,8] Intra-arterial digital subtraction angiography (IADSA) is the gold standard for detection of macrovascular abnormalities, but is an invasive procedure associated with some risk of complications.[9] MRI/MRA is less invasive, but has lower diagnostic accuracy for macrovascular causes than IADSA.

Currently, there is little guidance on which patients to select for (invasive) angiographic imaging and clinical practice thus varies widely.[10] Several factors have been associated with a higher likelihood of finding a macrovascular cause, including younger age, lobar location and absence of hypertension.[11] Early risk stratification of patients with ICH might help physicians to make swift, well-informed decisions about who to select for further angiographic imaging.

We aimed to develop and externally validate a prediction model to estimate the probability of finding a macrovascular cause in patients with non-traumatic ICH, based on patient characteristics, hemorrhage characteristics and, optionally, CTA.

## **METHODS**

### **Development cohort**

We used data from the DIagnostic AngioGRAphy to find vascular Malformations (DIAGRAM) study, a prospective, multicenter cohort study that assessed yield and diagnostic accuracy of angiographic imaging (CTA, MRA, IADSA) in patients with non-traumatic ICH.[12] Between 2008 and 2014, 298 patients aged 18-70 years were included in 22 participating centers across the Netherlands. Patients over 45 years of age with hypertension and ICH in the basal ganglia, thalamus or posterior fossa were excluded, because of the low probability of finding an underlying macrovascular cause.[13] Also, patients with a known macrovascular abnormality, brain tumor or patients who used oral anticoagulants and had an INR of >2.5 at the time of ICH were excluded. All patients underwent CTA within seven days of the ICH, followed by MRI/MRA within four to eight weeks if the CTA was negative. Patients underwent subsequent IADSA if the results of CTA or MRI/MRA were inconclusive or negative. CTA or MRI/MRA were considered inconclusive if a macrovascular cause was suspected but a definite diagnosis could not yet be established. Scans were read both locally and centrally. In case of a new diagnosis, local centres were informed. One additional arteriovenous fistula was detected at central reading.

Two hundred ninety-one patients had a CTA of sufficient quality for assessment (98%). MRI/MRA was performed in 255 patients (86%), of whom 214 patients with a negative or inconclusive CTA and IADSA in 154 patients (52%), of whom 106 patients with a negative or inconclusive CTA (Supplemental Figure I). Quality of IADSA was insufficient for assessment in three patients. One hundred twenty-six patients had a negative or inconclusive CTA, but did not

undergo subsequent IADSA. The main reason for not performing IADSA in patients with a negative CTA was an alternative diagnosis on MRI/MRA, or reluctance of either patients or their treating physicians. Four patients with a negative CTA died before MRI/MRA could be performed. The outcome was presence of a macrovascular cause (AVM, aneurysm, dAVF, cavernoma, cerebral venous sinus thrombosis and developmental venous anomaly (DVA)) as cause of the hemorrhage, and was based on best available evidence from all findings (CTA, MRA, DSA) during one year follow-up. The DIAGRAM study was approved by the medical ethics committee of the University Medical Center Utrecht, the Netherlands, and local approval was obtained from all participating hospitals. All participants gave written informed consent.

### **Model development**

Candidate predictors were pre-selected based on the literature and included age, hypertension (defined as a history of hypertension, use of antihypertensive drugs before ICH or evidence of left ventricular hypertrophy on admission ECG), smoking, high alcohol intake (defined as four or more units per day), location of ICH (lobar, deep or posterior fossa), presence of small vessel disease (SVD) on non-contrast CT (NCCT) (defined as presence of white matter lesions, or a lacunar infarct in basal ganglia, thalamus or posterior fossa, irrespective of whether it had been symptomatic or was an asymptomatic finding (see Supplemental Methods for a detailed description of SVD assessment and Supplemental Figure II)) and CTA. We developed two models; one model based on patient characteristics and NCCT (DIAGRAM score) and another model based on patient characteristics, NCCT and results from CTA imaging for use in higher resource settings (DIAGRAM+ score), which may help to estimate the probability of a macrovascular cause given that CTA is negative. For the current analysis, inconclusive CTAs were joined with positive results, because a CTA suggesting a macrovascular cause, yet

inconclusive, will always trigger further diagnostic tests. Given the one in ten rule with one predictive variable for every ten outcome events, we could study a maximum of seven predictors.[14,15]

## **Statistical Analysis**

Missing values for alcohol consumption (1%), smoking (1%) and CTA (2%) in the development cohort were imputed with single imputation. We used restricted cubic spline functions and graphs to assess whether age could be analyzed as linear term or needed transformation. We performed multivariable logistic regression analysis to study the association between candidate predictors and the presence of a macrovascular cause. The full model containing all candidate predictors was simplified by performing backward selection based on Akaike's Information Criterion (AIC). We internally validated the model by performing bootstrapping. A shrinkage factor was estimated from the bootstrap procedure and regression coefficients were multiplied by this shrinkage factor to correct for overfitting. Model performance was assessed with discrimination and calibration. Discrimination refers to the ability of the model to distinguish between someone with and without a macrovascular cause and was assessed with the c-statistic. Calibration assesses the correspondence between observed and predicted risk and was studied with a calibration plot. As a sensitivity analysis, we examined the performance of the models in a subset of patients (n=171), excluding those who did not undergo IADSA following a negative or inconclusive CTA. We generated prediction charts with predicted probabilities of finding a macrovascular abnormality for each combination of risk factors. Additionally, we created two prediction scores based on regression coefficients of the final multivariable regression models. For the prediction charts and scores, age was dichotomized at a value close to the mean.

## **External validation**

For external validation, we used a cohort of 173 patients with non-traumatic ICH.[16] Consecutive patients who underwent IADSA at the National Hospital for Neurology and Neurosurgery in London between 2010 and 2014 were retrospectively reviewed. Patients with non-traumatic ICH with available NCCT and CTA were included. NCCT and CTA were routinely performed in all patients with acute ICH presenting to the hyperacute stroke unit, unless there were contra-indications. The necessity of IADSA performance was judged in a weekly neuroradiological meeting, and was based on age, ICH location and medical history. MRI was performed according to clinical care, but was not systematically undertaken in all patients. The reference standard in the validation cohort was IADSA. All CTAs were reviewed blinded to IADSA result. The study was approved by the Clinical Governance Committee of the National Hospital and the UCL Institute of Neurology and National Hospital Joint Research Ethics Committee.

We applied the original regression equation to the validation data and calculated the predicted probability of finding a macrovascular cause for each patient. We assessed model performance with the c-statistic and calibration plots. As calibration is known to be strongly influenced by the incidence of the outcome in the validation population, we recalibrated the prediction models. Recalibration was performed by logistic regression analysis in the validation data with the linear predictor (the combination of regression coefficients with covariate values) as offset in the model. The resulting intercept was combined with the original regression coefficients to obtain predicted probabilities for the validation population. We present calibration of the models after recalibration, as in practice it is also advised to recalibrate a model before putting it to use.

Calibration results before recalibration are provided in the online supplement. Analyses were performed with R version 3.3.2. Results are reported in accordance with the TRIPOD statement.[17]

## RESULTS

Table 1 shows the baseline characteristics of the development and validation cohorts.

**Table 1. Baseline characteristics of development and external validation cohort**

	Development population		Validation population	
	Vascular malformation (n=69)	No vascular malformation (n=229)	Vascular malformation (n=78)	No vascular malformation (n=95)
Age, mean (SD), years	47 (12.7)	55 (10.5)	49 (17)	50 (13)
Male sex	45 (65)	140 (61)	39 (50)	54 (57)
Smoking (current)	20 (29)	52 (23)	-	-
High alcohol intake	4 (6)	32 (14)	-	-
Hypertension	16 (23)	79 (35)	16 (21)	37 (39)
Location of ICH				
Deep	5 (7)	80 (35)	14 (18)	46 (48)
Lobar	49 (71)	129 (56)	46 (59)	37 (39)
Posterior fossa	15 (22)	20 (9)	13 (17)	15 (16)
IVH	-	-	6 (8)	3 (3)
Signs of small vessel disease	4 (6)	116 (51)	12 (15)	35 (37)
CTA				
Positive	47 (68)	12 (5)	53 (68)	0 (0)
Inconclusive	4 (6)	8 (4)	11 (14)	7 (7)

Values are numbers (percentage), unless otherwise stated. ICH, intracerebral hemorrhage; IVH, intraventricular hemorrhage; SVD, small vessel disease; CTA, computed tomography angiography

Among 298 patients included in the development cohort, 69 (23%) had an underlying macrovascular cause. In the validation cohort a macrovascular cause was found in 78 of 173 patients (45%). Patients in the development cohort were slightly older (mean age 53 years, SD

11.5 versus 50 years, SD 15.0 in the validation cohort). The frequency of underlying vascular etiologies in each cohort is presented in Table 2.

**Table 2. Macrovascular causes underlying ICH in development and validation cohort**

	<b>Development cohort</b>	<b>Validation cohort</b>
	<b>n (%)</b>	<b>n (%)</b>
Arteriovenous malformation	34 (49)	68 (87)
Dural arteriovenous fistula	13 (19)	7 (9)
Cavernoma	10 (14)	-
Cerebral venous sinus thrombosis	4 (6)	-
Aneurysm	7 (10)	2 (3)
Developmental venous anomaly*	1 (1)	-
Carotid cavernous fistula	-	1 (1)
<b>Total</b>	<b>69</b>	<b>78</b>

ICH, intracerebral hemorrhage; \* this patient had a large developmental venous anomaly with partial thrombosis, which was clearly the cause of the ICH.

In multivariable analysis younger age, location of ICH, absence of signs of SVD and a positive or inconclusive CTA were independent predictors for presence of an underlying macrovascular cause (Table 3).

**Table 3. Odds ratios for presence of a macrovascular cause from multivariable models in the development cohort**

	Patient characteristics and NCCT OR (95% CI)	Patient characteristics, NCCT and CTA OR (95% CI)
Age	0.95 (0.93-0.98)	0.97 (0.94-1.00)
Location		
Deep	1 [Ref]	1 [Ref]
Lobar	7.2 (2.8-22.4)	4.0 (1.3-14.2)
Posterior fossa	19.3 (5.8-75.4)	9.9 (2.5-44.9)
Absence of SVD	11.8 (4.4-41.2)	11.8 (3.7-48.6)
Positive or inconclusive CTA	-	15.9 (7.5-35.5)

NCCT: non-contrast CT, CTA: computed tomography angiography, SVD: small vessel disease, CI: confidence interval, OR: odds ratio, [Ref]: reference

A simple model based on age, location of ICH and signs of SVD had a c-statistic of 0.83 (95% CI 0.78 to 0.88) in the development cohort after shrinkage. The predictive performance of the model increased if CTA was included as predictor (c-statistic 0.91; 95% CI 0.88 to 0.94).

Calibration of both models was accurate, as shown by the calibration plots (Figure 1). The original regression equations are provided in Supplemental Table I. When we excluded patients in whom IADSA was not performed following a negative or inconclusive CTA, c-statistics were similar to those of the full cohort analysis. Calibration plots and c-statistics are presented in Supplemental Figure II.

Figure 2 shows risk charts with estimated probabilities of finding a macrovascular cause according to age, ICH location, presence of SVD, and for the same predictors combined with CTA. The probability of finding a macrovascular cause ranged from 1% in patients aged 51 to 70 years with deep ICH and signs of SVD, up to more than 50% in patients aged 18 to 50 years with lobar or posterior fossa ICH and no signs of SVD. Two simple risk scores are presented in Supplemental Table II, which can be used in combination with Supplemental Figure III to obtain predicted probabilities for individual patients.

### **External validation**

External validation of the models showed a c-statistic of 0.66 (95% CI 0.58 to 0.74) for the model based on patient characteristics and NCCT, and a c-statistic of 0.88 (95% CI 0.83 to 0.94) for the model with additional CTA. The calibration plots show that the likelihood of finding a macrovascular cause increased along the range of predicted probabilities, with moderate

calibration for the model with patient characteristics and NCCT (Figure 1A) and good calibration for the model with additional CTA (Figure 1B). Before recalibration, both models systematically underestimated the probability of finding a macrovascular cause (Supplemental Figure IV).

## **DISCUSSION**

Our study shows that younger age, lobar or posterior fossa location of ICH, absence of signs of SVD, and a positive or inconclusive CTA are independent predictors for presence of a macrovascular cause in patients with non-traumatic ICH. We combined predictors in two practical prediction charts, which we externally validated. Estimated risks vary from 1% in patient aged 51 to 70 with deep ICH and signs of SVD, to more than 50% in patients aged 18 to 50 with lobar or posterior fossa ICH and no signs of SVD. Both models showed good discriminatory ability and calibration in the development cohort, whereas performance in external validation was moderate for the model with NCCT and good for the model including CTA.

Previously, two other prediction models have been described to predict the probability of a macrovascular cause in patients with non-traumatic ICH (Supplemental Table III). The simple ICH score was developed in a retrospective cohort of 160 patients with non-traumatic ICH in which presence of a macrovascular cause was determined with IADSA.[18] Performance of the risk score was moderate in both the development and external validation cohort. This model was derived from a high-risk population, as represented by the relatively young age (mean age 41 years) and high proportion of patients with a macrovascular cause (51%). The results may therefore not be generalizable to all patients with ICH suspected of having a vascular malformation, and the prediction model will likely overestimate the probability of finding a macrovascular cause. The secondary intracerebral hemorrhage score (SICH) was developed in a retrospective cohort of 623 patients with ICH in the US.[11] Presence of a macrovascular cause was determined with CTA. The model was based on patient characteristics and NCCT characteristics, which included enlarged vessels or calcifications along ICH margins and

hyperattenuation within a dural venous sinus or cortical vein. Independent validation in the US showed good performance of the model,[19] performance was moderate in an external validation study in the Netherlands.[3] NCCT categorization was a strong predictor for macrovascular causes, but characteristics were not always easy to recognize on NCCT,[3] which may limit easy application of the model in clinical practice. The DIAGRAM prediction score is the first model developed in a prospective cohort, excluding patients in whom yield of angiographic imaging has been shown to be very low (patients older than 45 years with a history of hypertension and a deep or posterior fossa bleed).[13] Next to known predictors for a vascular malformation, we were able to add signs of SVD as important predictor of absence of a macrovascular cause. To our knowledge, this is the first prediction model that also incorporated results from CTA imaging. This can be useful in healthcare settings where CTA is often or routinely used, and clinicians have to decide whether or not to perform MRI/MRA and/or IADSA after a negative CTA. The DIAGRAM prediction score may help to weigh the probability of finding a macrovascular cause against the risk of complications of IADSA.

Performance of the model based on patient characteristics and NCCT diminished in the external validation cohort. This is likely due to differences between the development and validation cohorts in terms of patient selection and choice of reference standard. Selection of patients influences prevalence of macrovascular causes and may affect predictor outcome associations, which in turn affect model performance. By selection of patients who underwent IADSA in the validation cohort, the prior probability of finding a macrovascular abnormality in this cohort was higher, which resulted in a systematically underestimated risk of finding a macrovascular cause by the prediction models. Simple recalibration improved correspondence between observed and predicted risks, supporting the hypothesis that differences in outcome incidence were the main

source of miscalibration. Selection of more high-risk patients may also have altered predictor-outcome associations. As a consequence, the discriminatory ability of the model may have decreased. Given differences between development and validation cohorts, validation of the DIAGRAM prediction model in a prospective cohort is necessary to further establish the robustness of the model.

Strengths of our study include the prospective nature of the development cohort and the standardized radiological work-up. Another strength is the external validation in a setting outside of the Dutch healthcare system. Our study also has limitations. First, the models were developed in a preselected group of patients with a relatively high likelihood of finding a macrovascular cause, excluding those older than 70 years of age, and patients over the age of 45 years with hypertension and deep ICH or ICH in the posterior fossa. This preselected group represents patients in whom the diagnostic dilemma is most pressing in clinical practice. Generalizability to older patients with non-traumatic ICH remains to be established. In the elderly, diagnostic tests to search for macrovascular causes of ICH are often performed in only a small proportion of patients.[20] Second, not all patients in the development cohort underwent IADSA. As a consequence, small AVMs or dAVFs may have been missed. However, patients were followed-up for one year to assess occurrence of re-bleeds and register possible causes of ICH identified during follow-up. Third, the association between CTA and presence of a macrovascular cause may have been overestimated, as CTA was also part of the reference standard. However, when we restricted our analyses in the development cohort to the patients who underwent IADSA, the discriminatory performance of the model remained similar. Fourth, MRI/MRA was not systematically performed in the validation cohort, which may have led to underestimation of the number of patients in whom a cavernoma was the cause of ICH.

The current models may facilitate selection of patients for further diagnostic work-up. The results of the model based on patient characteristics and NCCT suggest that in the absence of SVD, some form of angiographic imaging (CTA/MRA/IADSA) should be performed in all patients under 70 years of age, regardless of ICH location. If signs of SVD are seen on NCCT, CTA should still be considered in young patients (18-50 years old) with lobar and posterior fossa ICH, and in elderly patients (51-70 years old) with posterior fossa ICH. In settings where it is feasible to perform CTA in all patients shortly after ICH, the DIAGRAM+ score is particularly useful in patients in whom CTA was negative to guide the decision to perform these additional tests. Following a negative CTA, there is still a substantial chance of finding a macrovascular cause in patients without signs of SVD, both in young and in older patients. In these patients, performance of MRI/MRA and IADSA deserves consideration, especially in patients with lobar and posterior fossa ICH. It should be noted that also in patients with a deep ICH who do not have SVD nor hypertension (as defined by the inclusion criteria), there is an around 9% (in those 18 to 50 years) and 3% (in those 51 to 70 years) chance of finding a macrovascular cause of the ICH after a negative CTA. Whether or not in these patients further imaging will be performed should be determined as part of a shared decision making process between the patient and the team responsible for their care. Because the AVMs or dAVFs that are sought for with IADSA after a negative CTA will be small, IADSA should be performed in centers with ample experience in detecting such lesions. Although the prediction charts can provide guidance in decision-making, it should be noted that there is a degree of uncertainty around the presented estimates, as shown by the confidence intervals in Supplemental Figure III.

In conclusion, the DIAGRAM prediction charts can help to predict the probability of finding a macrovascular cause in both low and high resource settings. External validation of the models in

other prospective cohorts and in elderly patients is needed to gain further insight in the robustness of the models.

## **FUNDING**

This study was supported by a Dutch Heart Foundation grant (No 2007B048 to CJMK). CJMK is also supported by a clinical established investigator grant of the Dutch Heart Foundation (No 2012T077), and an Aspasia grant from the Netherlands Organisation for Health Research and Development, ZonMw (015008048). JPG and NAH are supported by a grant from the Dutch Heart Foundation (No 2013T128 to JPG).

## **DISCLOSURES**

None

## **THE DIAGRAM STUDY GROUP**

F E de Leeuw, H B Boogaarts, and E J van Dijk (Radboud University Nijmegen Medical Center, Nijmegen; 27 patients enrolled), W J Schonewille, W M J Pellikaan, and C Puppels-de Waard (St. Antonius hospital, Nieuwegein; 22), P L M de Kort, J P Peluso, and J H van Tuijl, (St. Elisabeth Hospital Tilburg; 20), J Hofmeijer, F B M Joosten (Rijnstate Hospital, Arnhem; 16), D W Dippel, L Khajeh (Erasmus MC University Medical Center, Rotterdam; 16), T W M Raaijmakers (Meander Medical Center, Amersfoort; 16), M J Wermer and M A van Walderveen (Leiden University Medical Center, Leiden; 14), H Kerkhoff, E Zock (Albert Schweitzer Hospital, Dordrecht; 14), K Jellema, G J Lycklama à Nijeholt (Medical Center Haaglanden, The Hague; 12), I M Bronner (Flevo Hospital, Almere; 12), M J M Remmers (Amphia Hospital, Breda; 9), R J G M Witjes (Tergooi Hospital, Blaricum; 8), H P Bienfait, K E Droogh-Greve (Gelre Hospital, Apeldoorn; 8), R C J M Donders (Diakonessen Hospital, Utrecht; 6), V I H Kwa (now: Onze Lieve Vrouwe Gasthuis, Slotervaart Hospital, Amsterdam; 4), T H Schreuder and C L Franke (Atrium Medisch Centrum, Heerlen; 4), J S Straver (Hofpoort Hospital, Woerden; 2), C Jansen (Gelderse Vallei Hospital, Ede; 1), S L M Bakker and C C Pleiter (Sint Franciscus Gasthuis, Rotterdam; 1), M C Visser (Free University Medical Center, Amsterdam; 1), C J J van Asch, B K Velthuis, G J E Rinkel, K M van Nieuwenhuizen, C J M Klijn (University Medical Center Utrecht; 90).

## REFERENCES

- [1] Feigin VL, Lawes CM, Bennett DA, et al. Worldwide stroke incidence and early case fatality reported in 56 population-based studies: a systematic review. *Lancet Neurol*2009;8:355-69.
- [2] Qureshi AI, Mendelow AD, Hanley DF. Intracerebral haemorrhage. *Lancet*2009;373:1632-44.
- [3] van Asch CJ, Velthuis BK, Greving JP, et al. External validation of the secondary intracerebral hemorrhage score in The Netherlands. *Stroke*2013;44:2904-06.
- [4] Bekelis K, Desai A, Zhao W, et al. Computed tomography angiography: improving diagnostic yield and cost effectiveness in the initial evaluation of spontaneous nonsubarachnoid intracerebral hemorrhage. *J Neurosurg*2012;117:761-66.
- [5] Delgado Almandoz JE, Schaefer PW, Forero NP, et al. Diagnostic accuracy and yield of multidetector CT angiography in the evaluation of spontaneous intraparenchymal cerebral hemorrhage. *AJNR Am J Neuroradiol*2009;30:1213-21.
- [6] Al-Shahi R, Warlow C. A systematic review of the frequency and prognosis of arteriovenous malformations of the brain in adults. *Brain*2001;124:1900-26.
- [7] Ohtani R, Kazui S, Tomimoto H, et al. Clinical and radiographic features of lobar cerebral hemorrhage: hypertensive versus non-hypertensive cases. *Intern Med* 2003;42:576-80.
- [8] Yamada S, Takagi Y, Nozaki K, et al. Risk factors for subsequent hemorrhage in patients with cerebral arteriovenous malformations. *J Neurosurg*2007;107:965-72.
- [9] Dawkins AA, Evans AL, Wattam J, et al. Complications of cerebral angiography: a prospective analysis of 2,924 consecutive procedures. *Neuroradiology*2007;49:753-59.
- [10] Cordonnier C, Klijn CJ, van Beijnum J, et al. Radiological investigation of spontaneous intracerebral hemorrhage: systematic review and trinational survey. *Stroke*2010;41:685-90.
- [11] Delgado Almandoz JE, Schaefer PW, Goldstein JN, et al. Practical scoring system for the identification of patients with intracerebral hemorrhage at highest risk of harboring an underlying vascular etiology: the Secondary Intracerebral Hemorrhage Score. *AJNR Am J Neuroradio* 2010;31:1653-60.
- [12] van Asch CJ, Velthuis BK, Rinkel GJ, et al. Diagnostic yield and accuracy of CT angiography, MR angiography, and digital subtraction angiography for detection of macrovascular causes of intracerebral haemorrhage: prospective, multicentre cohort study. *BMJ*2015;351:h5762.

- [13] Zhu XL, Chan MS, Poon WS. Spontaneous intracranial hemorrhage: which patients need diagnostic cerebral angiography? A prospective study of 206 cases and review of the literature. *Stroke*1997;28:1406-09.
- [14] Peduzzi P, Concato J, Kemper E, et al. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol*1996;49:1373-79.
- [15] Moons KG, de Groot JA, Bouwmeester W, et al. Critical appraisal and data extraction for systematic reviews of prediction modelling studies: the CHARMS checklist. *PLoS Med*2014;11:e1001744.
- [16] Wilson D, Ogungbemi A, Ambler G, et al. Developing an algorithm to identify patients with intracerebral haemorrhage secondary to a macrovascular cause. *European Stroke Journal*2017.
- [17] Collins GS, Reitsma JB, Altman DG, et al. Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis (TRIPOD). *Ann Intern Med*2015;162:735-36.
- [18] Olavarria VV, Bustamante G, Lopez MJ, et al. Diagnostic accuracy of a simple clinical score to screen for vascular abnormalities in patients with intracerebral hemorrhage. *J Stroke Cerebrovasc Dis*2014;23:2069-74.
- [19] Delgado Almandoz JE, Jagadeesan BD, Moran CJ, et al. Independent validation of the secondary intracerebral hemorrhage score with catheter angiography and findings of emergent hematoma evacuation. *Neurosurgery*2012;70:131-40.
- [20] Arboix A, Vall-Llosera A, Garcia-Eroles L, et al. Clinical features and functional outcome of intracerebral hemorrhage in patients aged 85 and older. *J Am Geriatr Soc*2002;50:449-54.

**Figure 1. Calibration plots of DIAGRAM prediction models in the development and validation cohort**

Figure legend: Model based on patient characteristics and NCCT (A), model based on patient characteristics, NCCT and CTA (B). The triangles indicate the observed frequencies with 95% confidence intervals by quintiles of predicted probability

**Figure 2. Prediction charts with absolute probabilities (%) of an underlying macrovascular cause in individual patients with ICH**

Figure legend: ICH: intracerebral hemorrhage, NCCT: non contrast CT, SVD: small vessel disease, CTA: computed tomography angiography