

## Biscalix[4]arene-Zn-tetraarylporphyrins

Dmitry M. Rudkevich, Willem Verboom, and David N. Reinhoudt\*

Laboratory of Organic Chemistry, University of Twente, P. O. Box 217, 7500 AE Enschede,  
The Netherlands

**Abstract:** By covalent combination of two calix[4]arene and one Zn-tetraarylporphyrin building blocks novel, large multifunctional receptors for molecular recognition have been prepared.

Design and synthesis of nanosize (1-10<sup>2</sup> nm) receptor molecules is one of the challenges in supramolecular chemistry.<sup>1</sup> We are currently exploring an approach based on the combination of different rigid building blocks (e.g. calix[4]arenes,<sup>2</sup> cavitands,<sup>3</sup> metallocalixarenes,<sup>4</sup> etc.) for the preparation of host molecules with *large hydrophobic surfaces*. Recently, we have described the preparation of a holand, a molecule consisting of two calix[4]arene and two resorcin[4]arene fragments covalently connected to generate a preorganized surface with a size of 1.5 x 2.0 nm.<sup>3</sup>


We report here the synthesis of large metalloreceptors **1** and **2** which are formed by combining one Zn-tetraarylporphyrin with two calix[4]arene molecules. Calix[4]arene is a versatile building block in supramolecular chemistry;<sup>5</sup> its rigid and lipophilic platform provides unique possibilities to build up the binding sites for ionic<sup>6</sup> and neutral<sup>7</sup> species. Zn-porphyrins are being used in biomimetic chemistry for recognition of biologically significant structures such as nucleic bases, amino acids, as well as anions.<sup>8</sup> Covalent combination of these two structural modules in one *preorganized* molecular shape can lead to receptors with a highly specific multifunctional complexing ability.<sup>9</sup>

The synthesis of **1** and **2** is based on the condensation of pyrrole with calix[4]arene based dialdehydes **3** and **5**, respectively; the aldehyde moieties were immobilized at either the "lower" or the "upper" rim of the calixarene platform (Scheme 1). Dialdehyde **3** was prepared in 74% yield by diametrical alkylation of the "lower" rim of *p-tert-butylcalix[4]arene*<sup>10</sup> with *o*-(4-bromobutoxy)benzaldehyde in refluxing acetonitrile using K<sub>2</sub>CO<sub>3</sub> as a base. Reaction of **3** with pyrrole in refluxing propionic acid<sup>11</sup> for 4-5 h afforded biscalix[4]arene-porphyrin **4** in 3-5% yield after purification by column chromatography (silica gel, CH<sub>2</sub>Cl<sub>2</sub>). Although, due to the hindered rotation of the aromatic rings connected to the porphyrin moiety, three different isomers are theoretically possible,<sup>11-13</sup> only one was isolated.

The positive ion FAB mass spectrum of compound **4** shows a peak at *m/z* 2192.2 with an intensity of 96%, which corresponds to the structure containing one porphyrin and two calix[4]arene moieties. The <sup>1</sup>H NMR spectrum of **4** in CDCl<sub>3</sub> shows two characteristic singlets for the porphyrin rings at 8.84 and 8.77 ppm;

the calix[4]arene aromatic region consists of four singlets with equal intensities at 6.93, 6.77, 6.67, and 6.51 ppm. These data clearly prove the adjacent-linked structure<sup>11</sup> of **4**.

Zn-complex **1**<sup>4</sup> was synthesized in 96% yield by refluxing free base porphyrin **4** with  $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$  in chloroform-methanol, 1:1 (For the <sup>1</sup>H NMR spectrum see Figure 1). In a temperature range of +30 - -95 °C the <sup>1</sup>H NMR spectra ( $\text{CD}_2\text{Cl}_2$ ) of the **1**-pyridine complex show only one set of signals for both receptor and guest molecules. No splitting was observed at low temperatures which proves the *adjacent-cis-linked* structure<sup>11</sup> of **1**; the two calix[4]arene fragments are equivalent with respect to the complexed pyridine molecule.



**Scheme 1.** Reagents and Conditions: i) pyrrole, propionic acid, reflux 4-5 h; ii)  $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$ ,  $\text{CHCl}_3\text{-CH}_3\text{OH}$ , 1:1, reflux 2 h.

Calixarene **5** which has two aldehyde moieties at the "upper" rim was obtained by reaction of 1,3-bis(chloroacetamido)tetrapropoxycalix[4]arene<sup>4</sup> with *m*-hydroxybenzaldehyde in 64% yield. Compound **6** in which two calix[4]arene cavities are directed towards the porphyrin plane, was prepared analogously to **4** by reaction of **5** and pyrrole in propionic acid [yield after column chromatography (silica gel,  $\text{CH}_2\text{Cl}_2$ -diethyl

ether, 5:1) 4%]. Metallation of **6** with  $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$  gave metalloporphyrin **2**<sup>14</sup> in a quantitative yield.

In compounds **2** and **6** the aromatic rings connected to the porphyrin can freely rotate;<sup>13</sup> the <sup>1</sup>H NMR spectra of **2** and **6** show two singlets for the porphyrin rings at 9.11, 8.89 and 9.03, 8.62, respectively, proving the *adjacent-linked* structures<sup>11</sup> of their cavities.

Biscalix[4]arene-Zn-tetraarylporphyrins **1** and **2** are interesting models for *multifunctional* recognition. Besides the active metalcenter which is able to complex anions and neutral species, compounds **1** and **2** contain two lipophilic calix[4]arene moieties in a large preorganized cavity with a size of 1.0-1.5 nm (according to CPK models). These calixarene fragments possessing OH, C(O)-NH groups may serve as additional receptor sites for hydrogen bonding and dipole-dipole interactions. The complexation properties of receptors **1** and **2** towards organic anions and polar neutral molecules are currently under investigation.




Figure 1. <sup>1</sup>H NMR spectrum of Zn-porphyrin **1** (400 MHz,  $\text{CDCl}_3$ ).

## References and Notes

1. (a) Cram, D. J. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1009; (b) Lehn, J.-M. *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 1304; (c) Whitesides, G. M.; Mathias, J. P.; Seto, C. T. *Science* **1991**, *254*, 1312; (d) Anderson, S.; Anderson, H. L.; Sanders, J. K. M. *Acc. Chem. Res.* **1993**, *26*, 469; (e) Bryant, J. A.; Knobler, C. B.; Cram, D. J. *J. Am. Chem. Soc.* **1990**, *112*, 1254.
2. van Dienst, E.; Iwema Bakker, W. I.; Engbersen, J. F. J.; Verboom, W.; Reinhoudt, D. N. *Pure Appl. Chem.* **1993**, *65*, 387;
3. Timmerman, P.; Verboom, W.; van Veggel, F. C. J. M.; van Hoorn, W. P.; Reinhoudt, D. N. *Angew. Chem.* **1994**, *106*, 1313.
4. Rudkevich, D. M.; Verboom, W.; Reinhoudt, D. N. *J. Org. Chem.* **1994**, in press.
5. (a) Gutsche, C. D. *Calixarenes, Monographs in Supramolecular Chemistry*; Stoddart, J. F., Ed.; The

- Royal Society of Chemistry: London, 1989; Vol. 1; (b) Vicens, J.; Böhmer, V. *Calixarenes: A Versatile Class of Macrocyclic Compounds*; Topics in inclusion science; Kluwer Academic Press: Dordrecht, 1991; Vol. 3; (c) van Loon, J.-D.; Verboom, W.; Reinhoudt, D. N. *Org. Prep. Proc. Int.* **1992**, *24*, 437.
6. For recent references from our laboratory see: (a) Ostaszewski, R.; Stevens, T. W.; Verboom, W.; Reinhoudt, D. N.; Kaspersen, F. M. *Recl. Trav. Chim. Pays-Bas* **1991**, *110*, 294; (b) Cobben, P. L. H. M.; Egberink, R. J. M.; Bomer, J. G.; Bergveld, P.; Verboom, W.; Reinhoudt, D. N. *J. Am. Chem. Soc.* **1992**, *114*, 10573; (c) Brzozka, Z.; Lammerink, B.; Reinhoudt, D. N.; Ghidini, E.; Ungaro, R. *J. Chem. Soc., Perkin Trans. 2* **1993**, 1037; (d) Morzherin, Y.; Rudkevich, D. M.; Verboom, W.; Reinhoudt, D. N. *J. Org. Chem.* **1993**, *58*, 7602; (e) Iwema-Bakker, W. I.; Haas, M.; Khoo-Beattie, C.; Ostaszewski, R.; Franken, S. M.; den Hertog, Jr., H. J.; Verboom, W.; de Zeeuw, D.; Harkema, S.; Reinhoudt, D. N. *J. Am. Chem. Soc.* **1994**, *116*, 123.
  7. For recent references from our laboratory see: (a) van Loon, J.-D.; Janssen, R. G.; Verboom, W.; Reinhoudt, D. N. *Tetrahedron Lett.* **1992**, *33*, 5125; (b) van Loon, J.-D.; Heida, J. F.; Verboom, W.; Reinhoudt, D. N. *Recl. Trav. Chim. Pays-Bas* **1992**, *111*, 353; (c) Reichwein, A. M.; Verboom, W.; Harkema, S.; Spek, A. L.; Reinhoudt, D. N. *J. Chem. Soc., Perkin Trans. 2* **1994**, 1167.
  8. For leading references see: (a) Kuroda, Y.; Ogoshi, H. *Synlett* **1994**, 319; (b) Danks, I. P.; Sutherland, I. O.; Hong Yap, C. *J. Chem. Soc., Perkin Trans. 1* **1990**, 421; (c) Mackay, L. G.; Wylie, R. S.; Sanders, J. K. M. *J. Am. Chem. Soc.* **1994**, *116*, 3141; (d) Bonar-Law, R. P.; Sanders, J. K. M. *J. Chem. Soc., Chem. Commun.* **1991**, 574; (e) Nappa, M.; Valentine, J. S. *J. Am. Chem. Soc.* **1978**, *100*, 5075.
  9. Recently, Asfari et al. have prepared a free base *double* porphyrin double calix[4]arene in 0.4% yield. We were unable to reproduce the reported procedure. See: Asfari, Z.; Vicens, J.; Weiss, J. *Tetrahedron Lett.* **1993**, *34*, 627.
  10. Gutsche, C. D.; Iqbal, M.; Stewart, D. *J. Org. Chem.* **1986**, *51*, 742.
  11. Momenteau, M.; Mispelter, J.; Loock, B.; Bisagni, E. *J. Chem. Soc., Perkin Trans. 1* **1983**, 189.
  12. Momenteau, M. *Pure Appl. Chem.* **1986**, *58*, 1493.
  13. Crossley, M. J.; Field, L. D.; Forster, A. J.; Harding, M. M.; Sternhell, S. *J. Am. Chem. Soc.* **1987**, *109*, 341.
  14. Selected data for **1**: m.p. >330 °C; UV-VIS (CH<sub>2</sub>Cl<sub>2</sub>) λ<sub>max</sub> 426 nm; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.82, 8.67 (2 x s, 8H, pyrrole), 8.06 (d, J = 8.0 Hz, 4H, ArH), 7.69, 7.40 (2 x t, J = 8.0 Hz, 8H, ArH), 7.31 (d, J = 8.0 Hz, 4H, ArH), 6.79, 6.59, 6.50, 6.41 (4 x s, 16H, Calix), 6.52, 5.73 (2 x s, 4H, OH), 3.75, 2.94 (2 x d, J = 13.2 Hz, 8H, CH<sub>2</sub>-calix), 2.51, 2.48 (ABq, J = 13.2 Hz, 8H, CH<sub>2</sub>-calix), 3.8-3.7, 3.6-3.5, 3.1-3.0, 2.9-2.8 (4 x m, 16H, OCH<sub>2</sub>), 1.4-1.1 (m, 16H, CH<sub>2</sub>), 1.15, 1.11 (2 x s, 36H, *t*-Bu), 0.89 (s, 36H, *t*-Bu); FAB-MS *m/z* 2257.1 (M<sup>+</sup>, calc. 2257.0). Selected data for **2**: m.p. >330 °C; UV-VIS (CH<sub>2</sub>Cl<sub>2</sub>) λ<sub>max</sub> 424 nm; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.11, 8.76 (2 x s, 8H, pyrrole), 8.13 (d, J = 8.0 Hz, 4H, ArH), 8.03 (s, 4H, ArH), 7.88 (bs, 4H, NH), 7.88, 7.30 (2 x t, J = 8.0 Hz, 8H, ArH), 7.2-7.0 (m, 10H, Calix), 6.87 (t, J = 7.9 Hz, 2H, Calix), 6.69 (d, J = 7.9 Hz, 4H, Calix), 6.26 (s, 4H, Calix), 4.81 (s, 8H, CH<sub>2</sub>C(O)), 4.51, 4.39, 3.36, 3.06 (4 x d, J = 13.2 Hz, 16H, CH<sub>2</sub>-calix), 4.1-3.8 (m, 16H, OCH<sub>2</sub>), 2.2-2.0 (m, 16H, CH<sub>2</sub>), 1.10 (t, J = 7.2 Hz, 24H, CH<sub>3</sub>); FAB-MS *m/z* 2147.6 ([M+H]<sup>+</sup>, calc. 2146.9). Satisfactory elemental analyses were obtained for all new compounds synthesized.

(Received in UK 13 July 1994; accepted 29 July 1994)