

# On Graphs Satisfying a Local Ore-Type Condition

A. S. Asratian\*  
H. J. Broersma  
J. van den Heuvel  
H. J. Veldman

FACULTY OF APPLIED MATHEMATICS  
UNIVERSITY OF TWENTE  
P.O. BOX 217, 7500 AE ENSCHEDE,  
THE NETHERLANDS

## ABSTRACT

For an integer  $i$ , a graph is called an  $L_i$ -graph if, for each triple of vertices  $u, v, w$  with  $d(u, v) = 2$  and  $w \in N(u) \cap N(v)$ ,  $d(u) + d(v) \geq |N(u) \cup N(v) \cup N(w)| - i$ . Asratian and Khachatryan proved that connected  $L_0$ -graphs of order at least 3 are hamiltonian, thus improving Ore's Theorem. All  $K_{1,3}$ -free graphs are  $L_1$ -graphs, whence recognizing hamiltonian  $L_1$ -graphs is an NP-complete problem. The following results about  $L_1$ -graphs, unifying known results of Ore-type and known results on  $K_{1,3}$ -free graphs, are obtained. Set  $\mathcal{K} = \{G | K_{p,p+1} \subseteq G \subseteq K_p \vee \overline{K_{p+1}} \text{ for some } p \geq 2\}$  ( $\vee$  denotes join). If  $G$  is a 2-connected  $L_1$ -graph, then  $G$  is 1-tough unless  $G \in \mathcal{K}$ . Furthermore, if  $G$  is a connected  $L_1$ -graph of order at least 3 such that  $|N(u) \cap N(v)| \geq 2$  for every pair of vertices  $u, v$  with  $d(u, v) = 2$ , then  $G$  is hamiltonian unless  $G \in \mathcal{K}$ , and every pair of vertices  $x, y$  with  $d(x, y) \geq 3$  is connected by a Hamilton path. This result implies that of Asratian and Khachatryan. Finally, if  $G$  is a connected  $L_1$ -graph of even order, then  $G$  has a perfect matching. © 1996 John Wiley & Sons, Inc.

## 1. INTRODUCTION

We use Bondy and Murty [6] for terminology and notation not defined here and consider finite simple graphs only.

A classical result on hamiltonian graphs is the following.

---

\*On leave from Department of Mathematical Cybernetics, Yerevan State University, Yerevan, 375049, Republic of Armenia. Supported by the Netherlands Organization for Scientific Research (N.W.O.)

**Theorem 1** (Ore [11]). If  $G$  is a graph of order  $n \geq 3$  such that  $d(u) + d(v) \geq n$  for each pair of nonadjacent vertices  $u, v$ , then  $G$  is hamiltonian.

In Asratian<sup>1</sup> and Khachatryan [7], Theorem 1 was improved to a result of local nature, Theorem 2 below. For an integer  $i$ , we call a graph an  $L_i$ -graph ( $L$  for local) if, for each triple of vertices  $u, v, w$  with  $d(u, v) = 2$  and  $w \in N(u) \cap N(v)$ ,

$$d(u) + d(v) \geq |N(u) \cup N(v) \cup N(w)| - i,$$

or, equivalently (see [7]),

$$|N(u) \cap N(v)| \geq |N(w) \setminus (N(u) \cup N(v))| - i.$$

**Theorem 2** [7]. If  $G$  is a connected  $L_0$ -graph of order at least 3, then  $G$  is hamiltonian.

Clearly, Theorem 2 implies Theorem 1.

Almost all of the many existing generalizations of Theorem 1 only apply to graphs  $G$  with large edge density ( $|E(G)| \geq \text{constant} \cdot |V(G)|^2$ ) and small diameter ( $o(|V(G)|)$ ). An attractive feature of Theorem 2 is that it applies to infinite classes of graphs  $G$  with small edge density ( $\Delta(G) \leq \text{constant}$ ) and large diameter ( $\geq \text{constant} \cdot |V(G)|$ ) as well. One such class is provided in [7]. For future reference also, we here present a similar class. For positive integers  $p, q$ , define the graph  $G_{p,q}$  of order  $pq$  as follows: its vertex set is  $\bigcup_{i=1}^q V_i$ , where  $V_1, \dots, V_q$  are pairwise disjoint sets of cardinality  $p$ ; two vertices of  $G_{p,q}$  are adjacent if and only if they both belong to  $V_i \cup V_{i+1}$  for some  $i \in \{1, \dots, q-1\}$ , or to  $V_1 \cup V_q$ . Considering a fixed integer  $p \geq 2$ , we observe that  $G_{p,q}$ , being an  $L_{2-p}$ -graph, is hamiltonian by Theorem 2 unless  $p = 2$  and  $q = 1$ ; furthermore,  $G_{p,q}$  has maximum degree  $3p - 1$  for  $q \geq 3$ , and diameter  $\lfloor \frac{q}{2} \rfloor = \lfloor \frac{1}{2p} |V(G_{p,q})| \rfloor$  for  $q \geq 2$ .

We define the family  $\mathcal{K}$  of graphs by

$$\mathcal{K} = \{G | K_{p,p+1} \subseteq G \subseteq K_p \vee \overline{K_{p+1}} \text{ for some } p \geq 2\},$$

where  $\vee$  is the join operation. The class of extremal graphs for Theorem 1, i.e., nonhamiltonian graphs  $G$  such that  $d(u) + d(v) \geq |V(G)| - 1 \geq 2$  for each pair of nonadjacent vertices  $u, v$ , is  $\mathcal{K} \cup \{K_1 \vee (K_r + K_s) | r, s \geq 1\}$  (see, e.g., Skupień [13]). We point out here that the class of extremal graphs for Theorem 2, i.e., nonhamiltonian  $L_1$ -graphs of order at least 3, is far less restricted. If  $G$  and  $H$  are graphs, then  $G$  is called  $H$ -free if  $G$  has no induced subgraph isomorphic to  $H$ . The following observation was first made in Asratian and Khachatryan [2].

**Proposition 3** [2]. Every  $K_{1,3}$ -free graph is an  $L_1$ -graph.

**Proof.** Let  $u, v, w$  be vertices of a  $K_{1,3}$ -free graph  $G$  such that  $d(u, v) = 2$  and  $w \in N(u) \cap N(v)$ . Then  $|N(w) \setminus (N(u) \cup N(v))| \leq 2$  and  $|N(u) \cap N(v)| \geq 1$ , implying that  $G$  is an  $L_1$ -graph. ■

In Bertossi [4] it was shown that recognizing hamiltonian line graphs, and hence recognizing hamiltonian  $K_{1,3}$ -free graphs is an NP-complete problem. Hence the same is true for recognizing hamiltonian  $L_1$ -graphs, and there is little hope for a polynomial characterization of the extremal graphs for Theorem 2.

<sup>1</sup>In [7] the last name of the first author was transcribed as “Hasratian”.

The study of  $L_1$ -graphs in subsequent sections was motivated by the interesting fact that the class of  $L_1$ -graphs contains all  $K_{1,3}$ -free graphs as well as all graphs satisfying the hypothesis of Theorem 1 (even with  $n$  replaced by  $n - 1$ ). The nature of the investigated properties of  $L_1$ -graphs is reflected by the titles of Sections 2, 3, and 4. The proofs of the obtained results are postponed to Section 5.

## 2. TOUGHNESS OF $L_1$ -GRAPHS

Let  $\omega(G)$  denote the number of components of a graph  $G$ . A graph  $G$  is  $t$ -tough if  $|S| \geq t \cdot \omega(G - S)$  for every subset  $S$  of  $V(G)$  with  $\omega(G - S) > 1$ . Clearly, every hamiltonian graph is 1-tough. Hence the following result implies Theorem 1 (for  $n \geq 11$ ).

**Theorem 4** (Jung [8]). If  $G$  is a 1-tough graph of order  $n \geq 11$  such that  $d(u) + d(v) \geq n - 4$  for each pair of nonadjacent vertices  $u, v$ , then  $G$  is hamiltonian.

By analogy, one might expect that Theorem 2 could be strengthened to the assertion that 1-tough  $L_4$ -graphs of sufficiently large order are hamiltonian. However, our first result shows that the problem of recognizing hamiltonian graphs remains NP-complete even within the class of 1-tough  $L_1$ -graphs. (Recall that the problem is NP-complete for  $L_1$ -graphs, and hence for 2-connected  $L_1$ -graphs.)

**Theorem 5.** If  $G$  is a 2-connected  $L_1$ -graph, then either  $G$  is 1-tough or  $G \in \mathcal{K}$ .

By Proposition 3, Theorem 5 extends the case  $k = 2$  of the following result.

**Theorem 6** (Matthews and Sumner [10]). Every  $k$ -connected  $K_{1,3}$ -free graph is  $\frac{k}{2}$ -tough.

In view of Theorem 6 we note that there exist 1-tough  $L_1$ -graphs of arbitrary connectivity that are not  $(1 + \varepsilon)$ -tough for any  $\varepsilon > 0$ . For example, consider the graphs  $K_{p,p}$  and  $K_p \vee \overline{K_p}$ , and the graphs obtained from  $K_{p,p}$  and  $K_p \vee \overline{K_p}$  by deleting a perfect matching ( $p \geq 3$ ).

## 3. HAMILTONIAN PROPERTIES OF $L_1$ -GRAPHS

If  $u, v, w$  are vertices of an  $L_0$ -graph such that  $d(u, v) = 2$  and  $w \in N(u) \cap N(v)$ , then  $N(w) \setminus (N(u) \cup N(v)) \supseteq \{u, v\}$ , and hence  $|N(u) \cap N(v)| \geq |N(w) \setminus (N(u) \cup N(v))| \geq 2$ . Thus our next result implies Theorem 2.

**Theorem 7.** Let  $G$  be a connected  $L_1$ -graph of order at least 3 such that  $|N(u) \cap N(v)| \geq 2$  for every pair of vertices  $u, v$  with  $d(u, v) = 2$ . Then each of the following holds.

- (a) Either  $G$  is hamiltonian or  $G \in \mathcal{K}$ .
- (b) Every pair of vertices  $x, y$  with  $d(x, y) \geq 3$  is connected by a Hamilton path of  $G$ .

An immediate consequence of Theorem 7 (a) is the following.

**Corollary 8** (Asratian, Ambartsumian, and Sarkisian [1]). Let  $G$  be a connected  $L_1$ -graph such that  $|N(u) \cap N(v)| \geq 2$  for every pair of vertices  $u, v$  with  $d(u, v) = 2$ . Then  $G$  contains a Hamilton path.

The lower bound 3 on  $d(x, y)$  in Theorem 7 (b) cannot be relaxed. For example, consider for  $p \geq 2$  the graphs  $K_{p,p}$  and  $K_p \vee \overline{K_p}$ , and for  $p \geq 4$  the graphs obtained from  $K_{p,p}$  and  $K_p \overline{K_p}$

by deleting a perfect matching. Each of these graphs satisfies the hypothesis of Theorem 7, but contains pairs of vertices at distance 1 or 2 that are not connected by a Hamilton path.

By Proposition 3, Theorem 7 (a) has the following consequence also.

**Corollary 9** (see, e.g., Shi Ronghua [12]). Let  $G$  be a connected  $K_{1,3}$ -free graph of order at least 3 such that  $|N(u) \cap N(v)| \geq 2$  for every pair of vertices  $u, v$  with  $d(u, v) = 2$ . Then  $G$  is hamiltonian.

An example of a graph that is hamiltonian by Theorem 7, but not by Theorem 2 or Corollary 9, is the graph obtained from  $G_{3,q}$  ( $q \geq 3$ ) by deleting the edges of a cycle of length  $q$ , containing exactly one vertex of  $V_i$  for  $i = 1, \dots, q$ .

Although Theorem 7 implies Theorem 2, in Section 5 we also present a direct proof of Theorem 2 as a simpler alternative for the algorithmic proof in Asratian and Khachatryan [7].

#### 4. PERFECT MATCHINGS OF $L_1$ -GRAPHS

Our last result is the following.

**Theorem 10.** If  $G$  is a connected  $L_1$ -graph of even order, then  $G$  has a perfect matching.

The graph  $K_{p,p+2}$  ( $p \geq 1$ ) is a connected  $L_2$ -graph of even order without a perfect matching. Thus Theorem 10 is, in a sense, best possible.

**Corollary 11** (Las Vergnas [9], Sumner [14]). If  $G$  is a connected  $K_{1,3}$ -free graph of even order, then  $G$  has a perfect matching.

**Corollary 12** (see, e.g., Bondy and Chvátal [5]). If  $G$  is a graph of even order  $n \geq 2$  such that  $d(u) + d(v) \geq n - 1$  for each pair of nonadjacent vertices  $u, v$ , then  $G$  has a perfect matching.

#### 5. PROOFS

We successively present proofs of Theorems 5, 7, 2 and 10, but first introduce some additional notation.

Let  $G$  be a graph. For  $S \subseteq V(G)$ ,  $N_G(S)$ , or just  $N(S)$  if no confusion can arise, denotes the set of all vertices adjacent to at least one vertex of  $S$ . For  $v \in V(G)$ , we write  $N_G(v)$  instead of  $N_G(\{v\})$ .

Let  $C$  be a cycle of  $G$ . We denote by  $\vec{C}$  the cycle  $C$  with a given orientation, and by  $\overleftarrow{C}$  the cycle  $C$  with the reverse orientation. If  $u, v \in V(C)$ , then  $u\vec{C}v$  denotes the consecutive vertices of  $C$  from  $u$  to  $v$  in the direction specified by  $\vec{C}$ . The same vertices, in reverse order, are given by  $v\overleftarrow{C}u$ . We use  $u^+$  to denote the successor of  $u$  on  $\vec{C}$  and  $u^-$  to denote its predecessor.

Analogous notation is used with respect to paths instead of cycles.

In the proofs of Theorems 5 and 7 we will frequently use the following key lemma.

**Lemma 13.** Let  $G$  be an  $L_1$ -graph,  $v$  a vertex of  $G$  and  $W = \{w_1, \dots, w_k\}$  a subset of  $N(v)$  of cardinality  $k$ . Assume  $G$  contains an independent set  $U = \{u_1, \dots, u_k\}$  of cardinality  $k$  such that  $U \cap (N(v) \cup \{v\}) = \emptyset$  and, for  $i = 1, \dots, k$ ,  $u_i w_i \in E(G)$  and  $N(u_i) \cap (N(v) \setminus W) = \emptyset$ . Then  $N(w_i) \setminus (N(v) \cup \{v\}) \subseteq N(u_i) \cup U$  ( $i = 1, \dots, k$ ).

**Proof.** Under the hypothesis of the lemma, we have

$$N(u_i) \cap N(v) = N(u_i) \cap W \quad (i = 1, \dots, k), \quad (1)$$

and since  $U$  is an independent set,

$$N(w_i) \setminus (N(u_i) \cup N(v)) \supseteq (N(w_i) \cap U) \cup \{v\} \quad (i = 1, \dots, k). \quad (2)$$

Since  $G$  is an  $L_1$ -graph, it follows that

$$\begin{aligned} 0 &\leq \sum_{i=1}^k (|N(u_i) \cap N(v)| - |N(w_i) \setminus (N(u_i) \cup N(v))| + 1) \\ &= \sum_{i=1}^k |N(u_i) \cap N(v)| - \sum_{i=1}^k (|N(w_i) \setminus (N(u_i) \cup N(v))| - 1) \\ &\leq \sum_{i=1}^k |N(u_i) \cap W| - \sum_{i=1}^k |N(w_i) \cap U| = 0. \end{aligned} \quad (3)$$

(Note that both  $\sum_{i=1}^k |N(u_i) \cap W|$  and  $\sum_{i=1}^k |N(w_i) \cap U|$  represent the number of edges with one end in  $U$  and the other in  $W$ .) We conclude that equality holds throughout (2) and (3). In particular, (2) holds with equality, implying that

$$N(w_i) \setminus (N(u_i) \cup N(v) \cup \{v\}) = N(w_i) \cap U \subseteq U,$$

and hence

$$N(w_i) \setminus (N(v) \cup \{v\}) \subseteq N(u_i) \cup U \quad (i = 1, \dots, k). \quad \blacksquare$$

**Proof of Theorem 5.** Let  $G$  be a 2-connected  $L_1$ -graph and assume  $G$  is not 1-tough. Let  $X$  be a subset of  $V(G)$  of minimum cardinality for which  $\omega(G - X) > |X|$ . Since  $G$  is 2-connected,  $|X| \geq 2$ . Set  $l = |X|$  and  $m = \omega(G - X) - 1$ , so that  $m \geq l \geq 2$ . Let  $H_0, H_1, \dots, H_m$  be the components of  $G - X$ .

In order to prove that  $G \in \mathcal{K}$ , we first show that

$$\text{for every nonempty proper subset } S \text{ of } X, |\{i | N(S) \cap V(H_i) \neq \emptyset\}| \geq |S| + 2. \quad (4)$$

Suppose  $S \subseteq X$ ,  $\emptyset \neq S \neq X$  and  $|\{i | N(S) \cap V(H_i) \neq \emptyset\}| \leq |S| + 1$ . Set  $T = X \setminus S$ . Then  $\omega(G - T) \geq m + 1 - |S| \geq l + 1 - |S| = |T| + 1$ . This contradiction with the choice of  $X$  proves (4).

We next show that

$$\text{if } v \notin X \text{ and } N(v) \cap X \neq \emptyset, \text{ then } N(v) \supseteq X. \quad (5)$$

Suppose  $v \notin X$  and  $N(v) \cap X \neq \emptyset$ , but  $N(v) \not\supseteq X$ . Set  $W = N(v) \cap X$  and  $k = |W|$ . Then  $1 \leq k < l$ . Let  $w_1, \dots, w_k$  be the vertices of  $W$ . By (4) and Hall's Theorem (see Bondy and Murty [6, p. 72]),  $N(W) \setminus X$  contains a subset  $U = \{u_1, \dots, u_k\}$  of cardinality  $k$  such that no two vertices of  $U \cup \{v\}$  are in the same component of  $G - X$  and  $u_1 w_1, \dots, u_k w_k \in$

$E(G)$ . By Lemma 13, we have  $N(w_i) \setminus (N(v) \cup \{v\}) \subseteq N(u_i) \cup U (i = 1, \dots, k)$ . But then  $\{|i|N(W) \cap V(H_i) \neq \emptyset\} \leq k + 1 = |W| + 1$ . This contradiction with (4) proves (5).

Let  $x$  be a vertex in  $X$  and  $y_i$  a vertex of  $H_i$  with  $N(y_i) \cap X \neq \emptyset (i = 0, 1, \dots, m)$ . Set  $Y = \{y_0, y_1, \dots, y_m\}$ . By (5),  $N(y_i) \supseteq X$  for all  $i$ , implying that  $N(x) \supseteq Y$ . Since  $G$  is an  $L_1$ -graph, we obtain

$$\begin{aligned} 0 &\leq |N(y_i) \cap N(y_j)| - |N(x) \setminus (N(y_i) \cup N(y_j))| + 1 \\ &= |X| - |N(x) \setminus (N(y_i) \cup N(y_j))| + 1 \\ &\leq |X| - |Y| + 1 = l - m \leq 0 \quad (i \neq j). \end{aligned} \tag{6}$$

Thus equality holds throughout (6). Hence  $m = l$  and  $N(x) \setminus (N(y_i) \cup N(y_j)) = Y$  whenever  $i \neq j$ . Consider a vertex  $y_h$  in  $Y$ . We have  $|X| \geq 2$  and hence  $|Y| \geq 3$ , so there exist distinct vertices  $y_i, y_j$  with  $y_h \neq y_i, y_j$ . Since  $N(x) \setminus (N(y_i) \cup N(y_j)) = Y$ , we obtain  $N(x) \cap V(H_h) = \{y_h\}$ . Since  $G$  is 2-connected, it follows that  $V(H_i) = \{y_i\}$  for all  $i$ , whence  $G \in \mathcal{K}$ . ■

**Proof of Theorem 7.** Let  $G$  satisfy the hypothesis of the theorem. Since  $|N(u) \cap N(v)| \geq 2$  whenever  $d(u, v) = 2$ ,

$$G \text{ is 2-connected.} \tag{7}$$

- (a) Assuming  $G$  is nonhamiltonian, let  $\vec{C}$  be a longest cycle of  $G$  and  $v$  a vertex in  $V(G) \setminus V(C)$  with  $N(v) \cap V(C) \neq \emptyset$ . Set  $W = N(v) \cap V(C)$  and  $k = |W|$ . Let  $w_1, \dots, w_k$  be the vertices of  $W$ , occurring on  $\vec{C}$  in the order of their indices. Set  $u_i = w_i^+ (i = 1, \dots, k)$  and  $U = \{u_1, \dots, u_k\}$ .

The choice of  $C$  implies that  $U \cap (N(v) \cup \{v\}) = \emptyset$ ,  $U$  is an independent set, and

$$\begin{aligned} N(u_i) \cap (N(v) \setminus W) &= N(u_i) \cap N(v) \cap (V(G) \setminus V(C)) = \emptyset \\ &(i = 1, \dots, k). \end{aligned} \tag{8}$$

Hence by Lemma 13,

$$N(w_i) \setminus (N(v) \cup \{v\}) \subseteq N(u_i) \cup U \quad (i = 1, \dots, k). \tag{9}$$

Noting that  $k \geq 2$  by (8) and the fact that  $|N(u_1) \cap N(v)| \geq 2$ , we now prove by contradiction that

$$u_i = w_{i+1}^- \quad (i = 1, \dots, k; \text{ indices mod } k). \tag{10}$$

Assume without loss of generality that  $u_1 \neq w_2^-$ , whence  $w_2^- \notin U$ . Then by (9),  $w_2^- \in N(u_2)$ . Since  $C$  is a longest cycle,  $w_2^- w_3^- \notin E(G)$ . Hence  $u_2 \neq w_3^-$ . Repetition of this argument shows that  $u_i \neq w_{i+1}^-$  and  $u_i w_i^- \in E(G)$  for all  $i \in \{1, \dots, k\}$ . By assumption,  $N(u_1) \cap N(v)$  contains a vertex  $x \neq w_1$ . By (8),  $x \in V(C)$ , say that  $x = w_i$ . But then the cycle  $w_1 v w_i u_1 \vec{C} w_i^- u_i \vec{C} w_1$  is longer than  $C$ . This contradiction proves (10).

Since  $C$  is a longest cycle, there exists no path joining two vertices of  $U \cup \{v\}$  with all internal vertices in  $V(G) \setminus V(C)$ . Hence by (10),  $\omega(G - W) > |W|$ . By (7) and Theorem 5, it follows that  $G \in \mathcal{K}$ .

- (b) Let  $x$  and  $y$  be vertices of  $G$  with  $d(x, y) \geq 3$  and let  $\vec{P}$  be a longest  $(x, y)$ -path. Assuming  $P$  is not a Hamilton path, let  $v$  be a vertex in  $V(G) \setminus V(P)$  with  $N(v) \cap V(P) \neq \emptyset$ . Set  $W = N(v) \cap V(P)$  and  $k = |W|$ . As in the proof of (a), we have  $k \geq 2$ . Let  $w_1, \dots, w_k$  be the vertices of  $W$ , occurring on  $\vec{P}$  in the order of their indices. Since  $d(x, y) \geq 3$ ,  $w_1 \neq x$  or  $w_k \neq y$ . Assume without loss of generality that  $w_k \neq y$ . Set  $u_i = w_i^+$  ( $i = 1, \dots, k$ ) and  $U = \{u_1, \dots, u_k\}$ .

Since  $P$  is a longest  $(x, y)$ -path, Lemma 13 can be applied to obtain

$$N(w_i) \setminus (N(v) \cup \{v\}) \subseteq N(u_i) \cup U \quad (i = 1, \dots, k). \quad (11)$$

We now establish the following claims.

$$\text{If } i < j \text{ and } u_j w_j^- \in E(G), \text{ then } u_i w_j \notin E(G). \quad (12)$$

Assuming the contrary, the path  $x \vec{P} w_i v w_j u_i \vec{P} w_j^- u_j \vec{P} y$  contradicts the choice of  $P$ .

$$w_1 = x. \quad (13)$$

Assuming  $w_1 \neq x$ , we have  $u_1 w_1^- \in E(G)$  by (11). As in the proof of (10), we obtain  $u_i w_i^- \in E(G)$  for all  $i \in \{1, \dots, k\}$  and  $u_i w_j \in E(G)$  for some  $j \in \{2, \dots, k\}$ , contradicting (12).

$$u_i = w_{i+1}^- \quad (i = 1, \dots, k - 1). \quad (14)$$

Assuming the contrary, set  $r = \min\{i \mid u_i \neq w_{i+1}^-\}$ . As in the proof of (10), we obtain  $u_i w_i^- \in E(G)$  for all  $i \in \{r + 1, \dots, k\}$ . Hence by (12),  $u_i w_j \notin E(G)$  whenever  $i \leq r$  and  $j \geq r + 1$ . By Lemma 13, it follows that  $N(w_i) \setminus (N(v) \cup \{v\}) \subseteq N(u_i) \cup \{u_1, \dots, u_r\}$  ( $i = 1, \dots, r$ ). Hence  $u_{r+1} w_i \notin E(G)$  for  $i \leq r$ , implying that  $\emptyset \neq (N(u_{r+1}) \cap N(v)) \setminus \{w_{r+1}\} \subseteq \{w_{r+2}, \dots, w_k\}$ , contradicting (12).

$$\text{For every longest } (x, y)\text{-path } Q, V(G) \setminus V(Q) \text{ is an independent set.} \quad (15)$$

It suffices to show that  $N(v) \subseteq V(P)$ . Suppose  $v$  has a neighbor  $v_1 \in V(G) \setminus V(P)$ . The choice of  $P$  implies  $N(v_1) \cap (U \cup W) = \emptyset = N(v_1) \cap N(w_1) \cap (V(G) \setminus (V(P) \cup \{v\}))$ . In particular,  $d(v_1, w_1) = 2$  and hence  $|N(v_1) \cap N(w_1)| \geq 2$ . Using (14) and the assumption  $d(x, y) \geq 3$ , we conclude that  $v_1$  and  $w_1$  have a common neighbor  $z$  on  $u_k^+ \vec{P} y^-$ . By (11),  $u_1 z \in E(G)$ . Repeating the above arguments with  $P$  and  $v_1$  instead of  $P$  and  $v$ , we obtain  $v_1 y \in E(G)$  (since  $v_1 x \notin E(G)$ ), and  $v_1 z^{++} \in E(G)$ . Now the path  $x u_1 z \vec{P} w_2 v v_1 z^{++} \vec{P} y$  contradicts the choice of  $P$ .

$$N(u_i) \subseteq V(P) \quad (i = 1, \dots, k - 1). \quad (16)$$

Assuming  $N(u_i) \not\subseteq V(P)$  for some  $i \in \{1, \dots, k - 1\}$ , the path  $x \vec{P} w_i v w_{i+1} \vec{P} y$  contradicts (15).

The above observations justify the following conclusions.

If some longest  $(x, y)$ -path does not contain the vertex  $z$ , then either

$$zx \in E(G) \text{ or } zy \in E(G). \quad (17)$$

If  $\vec{Q}$  is any longest  $(x, y)$ -path,  $z \notin V(Q), q \in V(Q)$  and  $zq \in E(G)$ ,  
 then the vertices of  $x\vec{Q}q$  (if  $zx \in E(G)$ ) or  $q\vec{Q}y$  (if  $zy \in E(G)$ )  
 are alternately neighbors and nonneighbors of  $z$ . (18)

Henceforth additionally assume  $P$  and  $v$  are chosen in such a way that

$$d(v) \text{ is as large as possible.} \tag{19}$$

If  $u_i x \in E(G)$  for all  $i \in \{1, \dots, k - 1\}$ , then, considering the path  $x\vec{P}w_i v w_{i+1} \vec{P}y$ , (18) and (19) imply  $u_i$  has no neighbor on  $u_k \vec{P}y$  ( $i = 1, \dots, k - 1$ ). Together with (16) this implies  $\omega(G - W) > |W|$ . By (7) and Theorem 5 we conclude that  $G \in \mathcal{K}$ , contradicting the fact that  $G$  has diameter at least 3. Hence, for some  $i \in \{2, \dots, k - 1\}$ ,  $u_i$  is not adjacent to  $x$ . By (17), we obtain

$$u_i y \in E(G) \text{ for some } i \in \{2, \dots, k - 1\}. \tag{20}$$

Let  $r = \min\{i \in \{2, \dots, k - 1\} | u_i y \in E(G)\}$  and  $s = \max\{i \in \{1, \dots, k - 1\} | u_i x \in E(G)\}$ . We first show

$$r > s. \tag{21}$$

Assuming the contrary, consider the vertex  $w_s$ . Clearly, (18) implies  $u_s w_j \in E(G)$  for all  $j \in \{1, \dots, s\}$ . If  $j \in \{1, \dots, s\}$  and  $u_j x \in E(G)$ , then, considering the path  $x\vec{P}w_j u_s \vec{P}w_{j+1} v w_{s+1} \vec{P}y$  and using (18) again, we obtain  $u_j w_s \in E(G)$ . Hence  $N(x) \cap U \subseteq N(w_s)$ . Clearly, (18) implies  $N(y) \cap \{u_r, \dots, u_{s-1}\} \subseteq N(w_s)$  and  $u_r w_j \in E(G)$  for all  $j \in \{r + 1, \dots, k\}$ . If  $j \in \{s, \dots, k\}$  and  $u_j y \in E(G)$ , then, considering the path  $x\vec{P}w_r v w_j \vec{P}u_r u_j^+ \vec{P}y$  and using (18) again, we obtain  $u_j w_{r+1} \in E(G)$  and hence  $u_j w_s \in E(G)$ . Hence  $N(y) \cap U \subseteq N(w_s)$ . We conclude that  $U \subseteq N(w_s)$ . Hence  $|N(w_s) \setminus (N(u_r) \cup N(v))| \geq k + 1$ , while  $|N(u_r) \cap N(v)| \leq k - 1$ . This contradiction with the fact that  $G$  is an  $L_1$ -graph completes the proof of (21).

Let  $j \in \{r, \dots, k\}$ . By (17) and (21),  $u_j y \in E(G)$  and by (18),  $u_j w_k \in E(G)$ . Suppose  $u_j w_r \notin E(G)$ . Then, by (18),  $u_j w_i \notin E(G)$  for all  $i \in \{1, \dots, r\}$ . Hence  $|N(u_j) \cap N(v)| \leq k - r$ , while  $|N(w_k) \setminus (N(u_j) \cup N(v))| \geq k - r + 2$ , a contradiction. Thus

$$u_j w_r \in E(G) \text{ for all } j \in \{r, \dots, k\}. \tag{22}$$

Now consider the path  $x\vec{P}w_r v w_{r+1} \vec{P}y$ , and let  $p = \min\{i \in \{2, \dots, r\} | u_r w_i \in E(G)\}$ ,  $j \in \{p - 1, \dots, r - 1\}$ . By (17) and (21),  $u_j x \in E(G)$  and by (18),  $u_j w_p \in E(G)$ . Suppose  $u_j w_r \notin E(G)$ . Then, by (18),  $u_j w_i \notin E(G)$  for all  $i \in \{r, \dots, k\}$ . Hence  $|N(u_j) \cap N(u_r)| \leq r - p$ , while  $|N(w_p) \setminus (N(u_j) \cup N(u_r))| \geq r - p + 3$ , a contradiction. Thus

$$u_j w_r \in E(G) \quad \text{for all } j \in \{p - 1, \dots, r - 1\}. \tag{23}$$

By (22) and (23),  $|N(w_r) \setminus (N(u_r) \cup N(v))| \geq k - p + 3$ , while  $|N(u_r) \cap N(v)| \leq k - p + 1$ , our final contradiction. ■



An independent algorithmic proof of Theorem 7 (a), similar to the proof of Theorem 2 given in Asratian and Khachatryan [7], will appear in Asratian and Sarkisian [3].

We now use the arguments in the proof of Theorem 7 (a) to obtain a short direct proof of Theorem 2, as announced in Section 3.

**Proof of Theorem 2.** Let  $G$  be a connected  $L_0$ -graph with  $|V(G)| \geq 3$ . Assuming  $G$  is nonhamiltonian, define  $\vec{C}$ ,  $v$ ,  $W$ ,  $k$ ,  $w_1, \dots, w_k$ ,  $u_1, \dots, u_k$ ,  $U$  as in the proof of Theorem 7 (a). By the choice of  $C$ , all conditions in Lemma 13 are satisfied. Hence (1) and (2) hold. Since  $G$  is an  $L_0$ -graph, we obtain, instead of (3),

$$\begin{aligned} 0 &\leq \sum_{i=1}^k (|N(u_i) \cap N(v)| - |N(w_i) \setminus (N(u_i) \cup N(v))|) \\ &= \sum_{i=1}^k |N(u_i) \cap N(v)| - \sum_{i=1}^k |N(w_i) \setminus (N(u_i) \cup N(v))| \\ &\leq \sum_{i=1}^k |N(u_i) \cap W| - \sum_{i=1}^k (|N(w_i) \cap U| + 1) = -k < 0, \end{aligned}$$

an immediate contradiction. ■

**Proof of Theorem 10** (by induction). Let  $G$  be a connected  $L_1$ -graph of even order. If  $|V(G)| = 2$ , then clearly  $G$  has a perfect matching. Now assume  $|V(G)| > 2$  and every connected  $L_1$ -graph of even order smaller than  $|V(G)|$  has a perfect matching. If  $G$  is a block, then by Theorem 5, the number of components, and hence certainly the number of odd components of  $G - S$  does not exceed  $|S|$ , and we are done by Tutte's Theorem (see Bondy and Murty [6, p. 76]). Now assume  $G$  contains a cut vertex  $w$ . Let  $G_1$  and  $G_2$  be distinct components of  $G - w$ . For  $i = 1, 2$ , let  $u_i$  be a neighbor of  $w$  in  $G_i$ . Since  $|N(u_1) \cap N(u_2)| = 1$  and  $G$  is an  $L_1$ -graph, we have  $N(w) \setminus (N(u_1) \cup N(u_2)) = \{u_1, u_2\}$ . In other words, every vertex in  $N(w) \setminus \{u_1, u_2\}$  is adjacent to either  $u_1$  or  $u_2$ . It follows that  $G_1$  and  $G_2$  are the only components of  $G - w$  and, since  $u_i$  is an arbitrary neighbor of  $w$  in  $G_i$ ,

$$G[N(w) \cap V(G_i)] \text{ is complete } (i = 1, 2). \tag{24}$$

Since  $|V(G)|$  is even, exactly one of the graphs  $G_1$  and  $G_2$ ,  $G_1$  say, has odd order. Set  $H = G[V(G_1) \cup \{w\}]$ . We now show that  $G_2$  and  $H$  are  $L_1$ -graphs.

Let  $x, y$ , and  $z$  be vertices of  $G_2$  such that  $d_{G_2}(x, y) = 2$  and  $z \in N_{G_2}(x) \cap N_{G_2}(y)$ . By (24),  $w \notin N_G(x) \cap N_G(y)$ , implying that  $N_{G_2}(x) \cap N_{G_2}(y) = N_G(x) \cap N_G(y)$ . Furthermore,  $N_{G_2}(z) \setminus (N_{G_2}(x) \cup N_{G_2}(y)) \subseteq N_G(z) \setminus (N_G(x) \cup N_G(y))$ . Since  $G$  is an  $L_1$ -graph, it follows that  $G_2$  is an  $L_1$ -graph.

A similar argument shows that  $H$  is an  $L_1$ -graph.

Since, moreover, the graphs  $G_2$  and  $H$  have even order smaller than  $|V(G)|$ , each of them has a perfect matching. The union of the two matchings is a perfect matching of  $G$ . ■

## References

- [1] A. S. Asratian, O. A. Ambartsumian, and G. V. Sarkisian, Some local condition for the hamiltonicity and pancyclicity of a graph, *Doclady Academ. Nauk Armenian SSR (Russian)* **19** (1) (1990), 19–22.
- [2] A. S. Asratian and N. K. Khachatryan, personal communication.
- [3] A. S. Asratian and G. V. Sarkisian, Some hamiltonian properties of graphs with local Ore's type conditions, in preparation.
- [4] A. A. Bertossi, The edge hamiltonian path problem is NP-complete, *Information Processing Lett.* **13** (1981), 157–159.
- [5] J. A. Bondy and V. Chvátal, A method in graph theory, *Discrete Math.* **15** (1976), 111–135.
- [6] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*, Macmillan, London and Elsevier, New York (1976).
- [7] A. S. Hasratian and N. K. Khachatryan, Some localization theorems on hamiltonian circuits, *J. Combinatorial Theory B* **49** (1990), 287–294.
- [8] H. A. Jung, On maximal circuits in finite graphs, *Ann. Discrete Math.* **3** (1978), 129–144.
- [9] M. Las Vergnas, A note on matchings in graphs. *Cahiers du Centre d'Etudes de Recherche Operationelle* **17** (1975), 257–260.
- [10] M. M. Matthews and D. P. Sumner, Hamiltonian results in  $K_{1,3}$ -free graphs, *J. Graph Theory* **8** (1984), 139–146.
- [11] O. Ore, Note on hamiltonian circuits, *Amer. Math. Monthly* **67** (1960), 55.
- [12] Shi Ronghua, 2-Neighborhoods and hamiltonian conditions, *J. Graph Theory* **16** (1992), 267–271.
- [13] Z. Skupién, An improvement of Jung's condition for hamiltonicity, *30. Internationales Wissenschaftliches Kolloquium, TH Ilmenau* (1985), Heft **5**, 111–113.
- [14] D. P. Sumner, Graphs with 1-factors, *Proc. Amer. Math. Soc.* **42** (1974), 8–12.

Received October 18, 1994